

# Can a Benign Ovarian Tumour develop a Malignant Endometrial Neoplasm? Sertoliform Ovarian Cystoadenoma getting together with a Low Malignancy Degree Endometrioid Adenocarcinoma G2

Jose Martinez-Mas <sup>1</sup>, Andres Bueno Crespo <sup>2</sup>, Pedro J. Garcia Laencina <sup>3</sup>, Juan Pedro Martínez Cendán <sup>4</sup>, Alejandra Isaac Montero <sup>5</sup>, Asensio B. Garcia Nicolas <sup>6</sup>, Manuel Remezal-Solano <sup>7</sup>

1. Obstetrics and Gynecology Department, Santa Lucía University Hospital, Cartagena, Murcia, Spain 2. Informatics Department, San Antonio Catholic University of Murcia (Spain) 3. Informatics Department, University Center for Defence, Murcia 4. Obstetrics and Gynecology Department, Catholic University of Murcia (UCAM), Murcia, Spain, Cartagena, ESP 5. Pathologic Anatomy, University Hospital Santa Lucia 6. Obstetrics and Gynecology, University Hospital Santa Lucia 7. Obstetrics and Gynecology Department, Catholic University of Murcia (UCAM), Murcia, Spain

✉ **Corresponding author:** Jose Martinez-Mas, jomartmas@gmail.com

**Categories:** Obstetrics/Gynecology

**Keywords:** sertoliform cystadenoma, endometrioid adenocarcinoma, estrogenic overproduction, induced endometrial cancer, postmenopausal

## How to cite this poster

Martinez-Mas J, Bueno Crespo A, Garcia Laencina P J, et al. (2015) Can a Benign Ovarian Tumour develop a Malignant Endometrial Neoplasm? Sertoliform Ovarian Cystoadenoma getting together with a Low Malignancy Degree Endometrioid Adenocarcinoma G2. Cureus 7(9): e.

## Abstract

### Introduction:

Sertoli cells tumours are neoplasms with a very low incidence, being sertoliform cystadenomae extremely unusual. Its first description was made on 1982 by Young and Roth, with very few cases published after them. This is a neoplasm only described on postmenopausal women, with an average age of presentation of 68 year old, and they could suffer virilization symptoms in different degrees. Histologically, it is a well differentiated neoplasm, with a low malignancy degree, and good prognosis if it is confined to ovaries. In other way, the finding of this ovarian tumour getting together with an endometrioid adencarcinoma is an extremely rare case.

## Open Access

Published 09/09/2015

## Copyright

© Copyright 2015

Martinez-Mas et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Distributed under

Creative Commons CC-BY 3.0

### Medical Case:

A 66 year old woman, who attended our office relating postmenopausal metrorrhage, is presented. The patient had a medical history of diabetes mellitus type II, arterial hypertension, dyslipidemia, liver steatosis, glaucoma and appendectomy. She got three deliveries and had her menopause at the age of 53.

An outpatient hysteroscopy was carried on, finding three endometrial polyps, two of them appeared as normal, and the bigger one had a glandular appearance with atypical vascularization, taking biopsy of it that was informed as endometrioid type adenocarcinoma with mucus-secreting pattern.

Ultrasound scanning found an adnexal left 2.5 cm nodule with heterogeneous ultrasonic pattern.

Magnetic Resonance found that adnexal left nodule, with heterogeneous signal pattern and significant contrast agent captation, suggestive of malignancy.

Serum Ca125 detected was 5 UI/ml.

An abdominal total hysterectomy with double adnexectomy, was carried on, with intraoperative biopsy, informed as a borderline tumour. Due to this finding, the operation was completed with omentectomy, lymphadenectomy and peritoneal washing.

Last histological analysis identifies the ovarian tumour as sertoliform adenoma from rete ovarii, with hepatocyte multifocal metaplasia. This tumour is suspected of being estrogenic functioning one, probably was the inductor of low degree and moderately differentiated G2 endometrioid adenocarcinoma, that infiltrated less than 50% of myometrial wall found on Hysterectomy specimen. There was no lymphatic or omental metastases. FIGO staging IA, TNM staging T1aN0M0.

## Conclusions

Sertoliform cystoadenoma is a very rare finding, with only few cases reported. It could be an estrogenic functioning tumour and due to this characteristic could induce endometrial hyperplasia, and at last, could be the reason for appearance of low malignancy degree endometrial neoplasms.