Acute Myeloid Leukemia with Occult Systemic Mastocytosis or Atypical Mast Cells Post-induction

Salahuddin Khan ¹, Zain Abid ¹, Humza F. Siddiqui ¹, Saima Zahoor ², Ghulam Haider ²

¹. Oncology, Jinnah Postgraduate Medical Center, Karachi, PAK ². Oncology, Jinnah Postgraduate Medical Centre, Karachi, PAK

Corresponding author: Zain Abid, dr.zainabid@gmail.com

Abstract

Systemic mastocytosis (SM) is a state of disease that is related to the clonal, neoplastic proliferation of mast cells. Patients who present with SM-Acute Myeloid Leukemia (AML) often have the worst outcome. We present a case of an 18-year-old female who was diagnosed with AML (FLT3 (Fms like tyrosine kinase 3) and PML-RARA (promyelocytic leukemia-retinoic acid receptor alpha) translocation-negative) and after initial treatment with a standard induction regimen of cytarabine and daunorubicin (3+7 regimen), her bone marrow showed blast cells less than 5% and dense aggregates/sheets of atypical/immature mast cells with immunohistochemical stain CD117+ve and toluidine blue positive in mast cell aggregates. Mastocytosis is a clonal neoplastic proliferation of mast cells that accumulate in one or more organ system. Therefore, it is essential to diagnose systemic mastocytosis, particularly in patients of hematological neoplasms.

Categories: Internal Medicine, Oncology

Keywords: systemic mastocytosis, acute myeloid leukemia, associated clonal hematologic non mast cell lineage (ahnmd)

Introduction

Systemic mastocytosis (SM) is an abnormal growth and accumulation of mast cells in extracutaneous organs, bone marrow being the most common site, with bone marrow biopsy and aspiration used in making the diagnosis [1]. Approximately 40% of patients with SM present with an associated clonal hematological non-mast cell lineage disorder [2-3]. Systemic mastocytosis with an associated, clonal, hematologic non-mast cell lineage disorder (SM-AHNMD) is the second most common type of mastocytosis [4].

a. These non-mast cell lineage disorders may include myelodysplastic syndrome (MDS), myeloproliferative neoplasm (MPN), acute myeloid leukemia (AML), chronic myelogenous leukemia, plasma cell myeloma, non-Hodgkin lymphoma, and unclassifiable myelogenous malignancies [5].

b. An associated AML has the worst prognosis [6].

We present a case of AML (M3), which after treatment showed atypical/immature mast cells in the post-induction marrow biopsy. The clinical course and prognosis of SM are variable and can range from plodding to aggressive.

Case Presentation

An 18-year-old female, married, presented with a history of low-grade fever, dizziness, and lethargy for three weeks. On examination, she was pale and her spleen was palpable. Her complete blood count revealed hemoglobin (Hb) of 11.2 gm/dl, total leukocyte count (TLC) of 52,300/mm³, and platelet count of 8000/mm³; her differential leukocyte count showed promyelocytes 79%. In Figure 1, the bone marrow aspiration showed blasts and abnormal promyelocytes 85%, and Auer rods were positive. The histochemical stain was positive for Sudan black, whereas erythropoiesis and myelopoiesis were depressed. The bone marrow trephine biopsy revealed sheets of blasts and abnormal promyelocytes consistent with acute promyelocytic leukemia (FAB AML M3 (French-American-British acute myeloid leukemia type 3)).

Received 11/21/2018
Review began 12/11/2018
Review ended 12/12/2018
Published 12/14/2018

© Copyright 2018

Khan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Khan S, Abid Z, Siddiqui H F, et al. (December 14, 2018) Acute Myeloid Leukemia with Occult Systemic Mastocytosis or Atypical Mast Cells Post-induction. Cureus 10(12): e3734. DOI 10.7759/cureus.3734

ARMED FORCES BONE MARROW TRANSPLANT CENTRE
 CMH Medical Complex, Rawalpindi (46000) Pakistan
 Tel: +92-51-9270076, Fax: +92-51-9271860, Email: afbmtc@outlook.com, Web: www.afbmtc.org

Report No.	Received on: 27/06/2018	Reported on: 28/06/2018	Sex: FEMALE
P No.	Rank:	ID No.	Age: 18 YEARS
Name:	Unit:	Contact No.	

CLINICAL INFORMATION
 This young female presented with history of low grade fever, dizziness and lethargy. Peripheral blood film from Ayub Medical Complex revealed blast cells. On examination she is pale and her spleen tip is palpable.

BLOOD COMPLETE PICTURE

Test	Result	Reference Range
Total Leukocytes (WBCs)	52.3 x 10 ⁹ /l	(Male 4.0 – 10.0) (Female 4.0 – 10.0)
Total Red Cells (RBCs)	3.80 x 10 ¹² /l	(Male 4.5 – 6.3) (Female 3.8 – 5.2)
Hemoglobin	11.2 g/dl	(Male 14 – 18) (Female 11.7 – 15.2)
Hematocrit (PCV)	0.33 l/l	(Male 0.39 – 0.52) (Female 0.35 – 0.46)
MCV	87.9 fl	(Male 77 – 91) (Female 77 – 91)
MCH	29.5 pg	(Male 26 – 32) (Female 26 – 32)
MCHC	33.5 g/dl	(Male 32 – 36) (Female 32 – 36)
Platelet Counts	06 x 10 ⁹ /l	(Male 150 – 400) (Female 150 – 400)
Absolute Neutrophils Count	- x 10 ⁹ /l	

DIFFERENTIAL

Test	Result	Reference Range
Neutrophils	08	40 – 75 %
Lymphocytes	06	20 – 45 %
Monocytes	-	02 – 10 %
Eosinophils	02	01 – 06 %
Basophils	-	
Promyelocytes	79	
Blasts cells	05	
ESR in mm at the end of 1 st hr	76	(Male 0–9) (Female 15)
RETICULOCYTE COUNT	0.5%	(0.5 – 2%)

BONE MARROW ASPIRATION

Site (s)	PSIS
Aspiration	Easy
Consistency of bone	Normal
Cellularity	Hypercellular packed marrow
Erythropoiesis	Depressed
Myelopoiesis	Depressed
M.E. Ratio	-
Blasts + Abnormal promyelocytes	>85%, Auer Rods +
Megakaryocytes	-
Lymphocytes	-
Plasma cells	-
Abnormal cells	-
Iron	-

MORPHOLOGY

Test	Result
Anisocytosis	+
Microcytosis	-
Poikilocytosis	+
Macrocytosis	-
Hypochromia	-
Spherocytes	-
RBC fragmentation	-
Target cells	-
Nucleated RBC's	04/100 WBC's
Others:	-

CYTOCHEMICAL STAIN
 Sudan Black- Positive +++

OPINION
ACUTE PROMYELOCYTIC LEUKAEMIA (FAB AML-M3)
 Bone marrow trephine biopsy report to follow

ADVISED

1. PCR for PML-RARα
2. FLT3 ITD and NPM1
3. Coagulation profile (PT, APTT and D-dimers)

Reported By: Maj Gen Tariq Mahmood Satti
 Brig Qamar Un Nisa Chaudhry
 Lt Col Nighat Shahbaz

Li Col Mehreen Ali Khan
 Maj Jahanzeb-Ur-Rehman
 Maj Ghassan Umair

FIGURE 1: Bone marrow aspiration (pre-induction chemotherapy)

Molecular genetics, including the polymerase chain reaction (PCR) for AML gene markers and PCR for FLT3 ITD (Fms like tyrosine kinase 3-internal tandem duplication) mutations, were negative. The TP53 gene mutation by fluorescence in situ hybridization (FISH) was negative. Immunophenotyping for acute leukemia was done, but the report was insufficient in its description. Cytogenetics showed that a culture of lymphocytes at 37°C failed to yield any growth. Promyelocytic leukemia-retinoic acid receptor alpha (PML-RARA) translocation by PCR was done, which was not detected.

We gave induction chemotherapy (3+7 regimen). On Day 16, her complete blood count (CBC) showed improvement with hemoglobin (Hb) 11.4 gm/dl, white blood cell (WBC) 5,100/mm³, and platelet count 240,000/mm³. The bone trephine procedure was done on post-induction Day 28, in which the bone marrow aspirate showed tri-lineage hematopoiesis, with a prominent increase in erythroid precursors and an increase in mast cells that constituted around 7% of the total nucleated non-erythroid population. These mast cells were spindle-shaped with centric to eccentric nuclei with clumped chromatin and absent nucleoli. They were also present in aggregates with erythroid precursors and prominent in the spicules. Blast cells were less than 5%. Bone trephine (hematoxylin and eosin (H&E)) section showed overall cellularity of around 55% to 60%. Cellular areas exhibited tri-lineage hematopoiesis alternating with areas of dense aggregates/sheets of atypical/immature mast cells. Immunohistochemical stains of mast cell aggregates were CD117 positive. Toluidine blue in the mast cell aggregates was metachromatic positive. Overall, the post-induction bone marrow (BM)/trephine biopsy showed a clearance of blast cells and increased mastocytes, which was suggestive of systemic mastocytosis with an associated hematologic neoplasm (SM-

AHN, systemic mastocytosis with associated acute myeloid leukemia (SM-AML)) according to the World Health Organization (WHO) 2016 classification of hematolymphoid neoplasms.

But clinically, according to updated WHO diagnostic criteria for systemic mastocytosis 2017, we cannot confirm the diagnosis of SM-AHN, as the mast cells were 7% in our patient while the criteria are >25%, and investigations like KIT-D816V or other activating point mutations of KIT and serum tryptase levels were also not available in our country. We started consolidation chemotherapy of the patient with high-dose cytarabine.

Discussion

The diagnosis of SM-AHNMD may be difficult to establish because the associated malignancy may mask the histological and cytological features of system mastocytosis (SM). The diagnosis can only be made when there is clear morphologic evidence of both SM with multifocal tissue infiltrates and AHNMD [7]. Malignant mast cells may abnormally express CD2 and/or CD25, which may be detected by immunohistochemistry or flow cytometry, which was not done in our case because of an affordability issue. Activating c-kit mutations are considered the hallmark of neoplastic mast cells [8], which was not available in our country. In SM-AHNMD, AHNMD may be diagnosed before, simultaneously with/after the diagnosis of SM. In some cases, the diagnosis of SM is missed/masked at the time of diagnosis, mainly due to the excess number of blasts and the tendency of mast cells to localize within the stroma of bone marrow particles [9]. However, the persistence of mastocytosis at the time of post-induction BM evaluation with a reduction in blast percentage helps us to diagnose SM at that time [10-14].

Conclusions

The diagnosis of SM is difficult to establish, as the associated malignancy may obscure the morphological features of SM. It is essential to diagnose SM, particularly in AHNMD, which may be difficult to diagnose during the course of the disease. However, a reduction in blast cell percentage at the time of a post-induction marrow evaluation helps in diagnosis.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Valent P, Akin C, Sperr WR, et al.: Diagnosis and treatment of systemic mastocytosis: state of the art . Br J Haematol. 2003, 122:695-717.
2. Wang SA, Hutchinson L, Tang G, et al.: Systemic mastocytosis with associated clonal hematological non-mast cell lineage disease: Clinical significance and comparison of chromosomal abnormalities in SM and AHNMD components. Am J Hematol. 2013, 88:219-224.
3. Zhrebker L, Cooper B, Krause JR: Systemic mastocytosis with associated acute myelogenous leukemia . Proc (Baylor Univ Med Cent). 2014, 27:22-24. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3862125/>.
4. Swerdlow SH, Campo E, Harris NL, et al.: WHO Classification of Tumors of Haematopoietic & Lymphoid Tissue. IARC, Lyon, France; 2008.
5. Pullarkat VA, Bueso-Ramos C, Lai R, et al.: Systemic mastocytosis with associated clonal hematological non-mast-cell lineage disease, analysis of clinicopathologic features and activating c-kit mutations. Am J Hematol. 2003, 73:12-17.
6. Pardanani A, Lim KH, Lasho TL, Finke C, McClure RF, Li C-Y, Tefferi A: Prognostically relevant breakdown of 123 patients with systemic mastocytosis associated with other myeloid malignancies. Blood. 2009, 114:3769. [10.1182/blood-2009-05-220145](https://doi.org/10.1182/blood-2009-05-220145)
7. Stoecker MM, Wang E: Systemic mastocytosis with associated clonal hematologic nonmast cell lineage disease: a clinicopathologic review. Arch Pathol Lab Med. 2012, 136:832-838.
8. Feger F, Dumas AR, Leriche L, Valent P, Arock M: Kit and c-kit mutations in mastocytosis: a short overview with special reference to novel molecular & diagnostic concepts. Int Arch Allergy Immuno. 2002, 127:110-114. [10.1159/000048179](https://doi.org/10.1159/000048179)
9. Sabattini E, Bacci F, Sagromoso C, Pileri SA: WHO classification of tumours of haematopoietic and lymphoid tissues in 2008: an overview. Pathologica. 2010, 102:83-87.
10. Bernd HW, Sotlar K, Lorenzen J, Osieka R, Fabry U, Valent P: Acute myeloid leukemia with t(8;21) associate systemic mastocytosis. Report of an unusual case review of the literature. J Clin Pathol. 2004, 57:324-328.
11. Mahadeo KM, Wolgast L, McMahon C, Cole PD: System mastocytosis in a child with acute myeloid leukemia . Pediatr Blood Cancer. 2011, 57:684-687.
12. Pullarkat ST, Pullarkat V, Kroft SH, et al.: Systemic mastocytosis associated with acute myeloid leukemia . J Hematopathol. 2009, 2:27-33.

13. Horny HP, Sotlar K, Sperr WR, Valent P: Systemic mastocytosis with associated clonal haematological non-mast cell lineage diseases: a histopathological challenge. *J Clin Pathol.* 2005, 57:604-608. [10.1136/jcp.2003.014860](https://doi.org/10.1136/jcp.2003.014860)
14. Valent P, Horny HP, Escribano L, et al.: Diagnostic criteria and classification of mastocytosis: a consensus proposal. *Leuk Res.* 2001, 25:603-625. [10.1016/S0145-2126\(01\)00038-8](https://doi.org/10.1016/S0145-2126(01)00038-8)