

Role of Ultrasound in the Diagnosis of Chronic Kidney Disease and its Correlation with Serum Creatinine Level

Shakeel Ahmed ¹, Sanobar Bughio ², Maria Hassan ², Sajan Lal ², Muhammad Ali ²

¹. Urology, Sindh Institute of Urology and Transplantation, Karachi, PAK ². Radiology, Dr. Ziauddin Hospital, Karachi, PAK

Corresponding author: Sanobar Bughio, drsanobarbughio@gmail.com

Abstract

Objective: We aimed to study the role of ultrasound in the diagnosis of chronic kidney disease (CKD) and its correlation with serum creatinine level.

Materials and methods: This was a retrospective cross sectional study conducted in the ultrasound department of Dr. Ziauddin Hospital Clifton campus, Karachi from April 6, 2017 to October 6, 2017 for a period of six months. A total of 200 patients with CKD and glomerular filtration rate (GFR) determined to be < 60 ml/min were included in this study. Blood tests were gathered from the chosen patients, serum creatinine estimation was done for those patients and they underwent ultrasonography on the same day to assess echogenicity, parenchymal thickness, cortical thickness, and longitudinal length. This information was noted in the pro forma.

Results: The average age of the patients was 54.62 ± 13.3 years. Mean serum creatinine was significant among echogenicity grades ($p=0.0005$). Mean parenchymal thickness was also significant among echogenicity grades ($p=0.0005$). Mean longitudinal length was also significant among echogenicity grades ($p=0.0005$). Mean corticomedullary distinction was also significant among echogenicity grades ($p=0.0005$). A statistically significant highly positive correlation was observed between serum creatinine and cortical echogenicity grading ($r=0.915 P = 0.0005$).

Conclusion: The best sonographic parameter that correlates with serum creatinine is renal cortical echogenicity and its grading in comparison to longitudinal length, parenchymal thickness, and cortical thickness in patients of CKD. Since renal cortical echogenicity has the advantage of being irreversible in comparison to serum creatinine levels, it can be used as a parameter of renal function.

Categories: Radiology, Nephrology

Keywords: chronic kidney disease, renal echogenicity, serum creatinine

Received 01/30/2019

Review began 03/11/2019

Review ended 03/12/2019

Published 03/12/2019

© Copyright 2019

Ahmed et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

A deranged creatinine level over a period of few months to years is termed chronic kidney disease (CKD). CKD is based on the extent of kidney damage, calculated through decreased glomerular filtration rate (GFR) (i.e. < 60 ml/min per 1.7 m^2) for more than three months [1, 2].

Ultrasonography is a noninvasive and inexpensive investigation modality with sufficient anatomical details necessary to diagnose renal diseases without exposing the patient to radiation or contrast and hence has replaced standard radiography in our country and abroad [3-5]. All these factors promote early detection and prediction of deranged renal function tests necessary for making a therapeutic decision.

Sonography identifies renal length, thickness, and echogenicity of renal parenchyma apart from its importance in detailing a dilated collecting system [6]. These details assist in identifying the extent of renal parenchymal damage and the possibility of its reversibility [7, 8], and the decision to perform a renal biopsy [9]. According to a study, abnormal sonographic findings were seen in 67% of cases of CKD [10].

Due to the presence of collagen, echogenicity is increased in interstitial fibrosis and glomerulosclerosis [11], but this has never been recognized. Increase in echogenicity may also increase interstitial inflammation. The human eye can also assess echogenicity but it is unreliable [12, 13]. In a small group of adults, renal parenchymal echogenicity can be reliably quantitated and established within a normal range [14]. It was found that there is significant correlation between renal length or cortical echogenicity with glomerular sclerosis or tubular atrophy [15].

Renal morphology can be determined by a number of means that include measuring renal length and volume and renal cortical thickness. Renal function can also be evaluated through renal length and cortical

How to cite this article

Ahmed S, Bughio S, Hassan M, et al. (March 12, 2019) Role of Ultrasound in the Diagnosis of Chronic Kidney Disease and its Correlation with Serum Creatinine Level. Cureus 11(3): e4241. DOI 10.7759/cureus.4241

thickness, and important clinical decisions can be made on its basis. Therefore serial sonographic evaluations are done to find out the progression of renal disease or its normality [16]. Although renal parenchymal volume is quite an accurate measurement in patients with end stage renal disease, measurement of renal longitudinal length is sufficient in normal patients [17].

Hence, ultrasound is a good modality to ascertain renal insufficiency and progression of disease. The aim of our study was to correlate renal echogenicity with serum creatinine levels and to investigate the significance of renal echogenicity in identifying the progression of CKD as well as the use of sonographic imaging in the grading of CKD.

Materials And Methods

This was a retrospective cross-sectional study conducted in the ultrasound department of Dr. Ziauddin Hospital Clifton campus, Karachi from April 6, 2017 to October 6, 2017 for a period of six months. All patients referred for an ultrasound of the kidneys, whose creatinine was checked on the same day on which the ultrasound was performed, were considered. A total of 200 patients were included in this study.

New patients presenting for CKD workup, patients known to have CKD as per operational definition, CKD stages 3/4/5 and GFR determined to be < 60 ml/min as calculated by the Modification of Diet in Renal Disease (MDRD) equation, and patients of age above 30 years (male and female) were included in the study. Known patients of acute kidney injury, kidney transplant patients, patients on hemodialysis, patients on peritoneal dialysis, patients with fatty liver, chronic liver disease and solitary kidney were excluded from the study.

Ultrasound of the kidneys and liver was performed using the standard B Mode grey scale ultrasound with sector curved array transducer of 3.5-5 MHz. The parenchymal echogenicity of both the liver and kidney was assessed by applying low tissue harmonic and speckle reduction imaging to reduce the interobserver bias. The gain and time gain compensation were adjusted manually. The longitudinal length was measured in a section visually estimated to represent the largest longitudinal section. The width and thickness were measured in a section perpendicular to the longitudinal axis of the kidney as assessed from the longitudinal image. It is not necessary to keep the ultrasound probe perpendicular to the skin. However, the level of this transverse section was placed quite close to the hilum of the kidney but at the same time free of the pelvis.

Statistical data analysis was done using the Statistical Package for the Social Sciences (SPSS version 20) (IBM Corp., Armonk, NY). Mean and SD were calculated for age; mean parenchymal thickness, mean longitudinal size, frequency and percentages were calculated for gender and grade of echogenicity. Statistical analysis was calculated using one way analysis of variance (ANOVA). The relationship between serum creatinine and sonographic parameters was assessed by correlation coefficient analysis. P values less than 0.05 were considered statistically significant.

Results

A total of 200 patients with CKD and GFR determined to be < 60 ml/min as calculated by the MDRD equation were included in this study. Twenty percent of the patients were below and equal to 40 years of age, 42.5% were between 41 and 60 years, and 37.5% were above 60 years of age. The average age of the patients was 54.62 ± 13.3 years; the average serum creatinine, parenchymal thickness, longitudinal length, and cortical thickness are shown in Table 1.

Variables	Mean \pm SD	95% Confidence Interval for Mean		Median (IQR)
		Lower Bound	Upper Bound	
Age (years)	54.62 \pm 13.30	52.77	56.47	56 (21)
Serum creatinine (mg/dl)	2.19 \pm 1.08	2.046	2.337	1.9 (1.3)
Parenchymal thickness	4.69 \pm 0.82	4.578	4.806	4.5 (0.9)
Longitudinal length	9.87 \pm 0.94	9.745	10.008	10 (0.8)
Cortical thickness	0.94 \pm 0.253	0.902	0.973	1.1 (0.3)

TABLE 1: Descriptive Statistics of the Characteristics of the Patients [n=200]

Out of the 200 patients, 60% were male and 40% were female. The ultrasound findings of the renal parenchymal changes and its grading, Grade 0 is shown in Figure 1.



FIGURE 1: Grade-0 Echogenicity of Renal Parenchyma Less Than That of the Liver with Maintained Corticomedullary Distinction

Sixty (30%) patients had Grade 0 parenchymal changes, 60 (30%) patients had Grade 1 parenchymal changes, 44 (22%) patients had Grade 2 parenchymal changes, 24 (12%) patients had Grade 3 parenchymal changes, and 12 (6%) patients had Grade 4 parenchymal changes.

The mean serum creatinine was 1.252 ± 0.050 mg/dl (95%CI: 1.239 to 1.265) for Grade 0, 1.853 ± 0.129 mg/dl (95%CI: 1.82 to 1.887) for Grade 1, 2.568 ± 0.651 mg/dl (95%CI: 2.370 to 2.766) for Grade 2, 3.275 ± 0.352 mg/dl (95%CI: 3.126 to 3.424) for Grade 3, and 5.033 ± 0.528 mg/dl (95%CI: 4.698 to 5.369) for Grade 4.

Mean serum creatinine was significant among echogenicity grades [ANOVA F-Value= 367.726; p=0.0005] (Table 2).

Echogenicity Grades [Grading based on ultrasound features]	Serum Creatinine (mg/dl)						F Value	P-Value	
	No. of Patients	Mean	SD	95% Confidence Interval for Mean		Lower Bound	Upper Bound		
				Lower Bound	Upper Bound				
Grade 0	60	1.252	0.050	1.239	1.265				
Grade 1	60	1.853	0.129	1.820	1.887				
Grade 2	44	2.568	0.651	2.370	2.766	367.726	0.0005		
Grade 3	24	3.275	0.352	3.126	3.424				
Grade 4	12	5.033	0.528	4.698	5.369				
Total	200	2.191	1.0431	2.046	2.337				

TABLE 2: Comparison of Serum Creatinine with Renal Cortical Echogenicity

Mean parenchymal thickness was also significant among echogenicity grades (ANOVA F-value= 31.628; p=0.0005) (Table 3).

Echogenicity Grades [Grading based on ultrasound features]	Parenchymal Thickness (cm)						F Value	P-Value		
	No. of Patients	Mean	SD	95% Confidence Interval for Mean						
				Lower Bound	Upper Bound					
Grade 0	60	4.397	0.101	4.371	4.423					
Grade 1	60	4.582	0.905	4.348	4.816					
Grade 2	44	5.048	0.920	4.768	5.328	31.628	0.0005			
Grade 3	24	4.433	0.481	4.230	4.637					
Grade 4	12	5.933	1.029	5.280	6.587					
Total	200	4.692	0.820	4.578	4.806					

TABLE 3: Comparison of Parenchymal Thickness with Renal Cortical Echogenicity

Mean longitudinal length was also significant among echogenicity grades (ANOVA F-value= 66.004; p=0.0005) (Table 4).

Echogenicity Grades [Grading based on ultrasound features]	Longitudinal Length (cm)						F Value	P-Value		
	No. of Patients	Mean	SD	95% Confidence Interval for Mean						
				Lower Bound	Upper Bound					
Grade 0	60	10.000	0.0000	10.000	10.000					
Grade 1	60	10.272	0.8114	10.062	10.481					
Grade 2	44	10.348	0.7457	10.121	10.574	66.004	0.0005			
Grade 3	24	8.758	0.4698	8.551	8.965					
Grade 4	12	7.792	0.8107	7.277	8.307					
Total	200	9.876	0.9450	9.745	10.008					

TABLE 4: Comparison of Longitudinal Length with Renal Cortical Echogenicity

Mean cortical thickness was also significant among echogenicity grades (ANOVA F-value= 477.83; p=0.0005) (Table 5).

Echogenicity Grades [Grading based on ultrasound features]	Cortical Thickness (cm)						F Value	P-Value		
	No. of Patients	Mean	SD	95% Confidence Interval for Mean						
				Lower Bound	Upper Bound					
Grade 0	60	1.100	0.0000	1.100	1.100					
Grade 1	60	1.093	0.0756	1.074	1.113					
Grade 2	44	0.900	0.1034	0.869	0.931		477.83	0.0005		
Grade 3	24	0.462	0.1135	0.415	0.510					
Grade 4	12	0.433	0.0888	0.377	0.490					
Total	200	0.938	0.2531	0.902	0.973					

TABLE 5: Comparison of Cortical Thickness with Renal Cortical Echogenicity

A statistically significant highly positive correlation was observed between serum creatinine and cortical echogenicity grading ($r=0.915$ $P = 0.0005$) as shown in Table 6. There was also a statistically significant positive correlation between mean parenchymal thickness and renal echogenicity ($r=0.336$; $P = 0.005$). There was also a statistically significant negative correlation between longitudinal length, cortical thickness with renal echogenicity ($r= -0.513$; $P = 0.005$) and ($r= -0.869$; $P = 0.0005$), respectively (Table 6).

A statistically significant negative correlation was also observed between longitudinal size and serum creatinine ($r= -0.505$; $P = 0.0005$); a statistically significant negative correlation was observed between cortical thickness and serum creatinine ($r= -0.845$; $P = 0.0005$) and a statistically significant positive correlation was observed between parenchymal thickness and serum creatinine ($r=0.413$; $P = 0.0005$) (Table 6).

		Parenchymal Thickness	Longitudinal Length	Cortical Thickness	Echogenicity Grade
Serum creatinine (mg/dl)	Pearson correlation	0.413**	-0.505**	-0.845**	0.915**
	P-Value	0.0005	0.0005	0.0005	0.0005
Parenchymal thickness	Pearson correlation		0.092	-0.240**	0.336**
	P-Value		0.194	0.001	0.005
Longitudinal length	Pearson correlation			0.634**	-0.513**
	P-Value			0.0005	0.0005
Cortical thickness	Pearson correlation				-0.869**
	P-Value				0.0005

TABLE 6: Statistical Correlation between Serum Creatinine and Mean Parenchymal Thickness, Mean Longitudinal Size, Mean Cortical Thickness and Echogenicity grade

** Correlation is significant at the 0.01 level (2-tailed).

Discussion

The term chronic kidney disease means progressive damage to kidneys caused by structural or functional abnormalities of the kidney that can get worse over time. When the damage gets worse, the kidneys stop working, with or without decreasing GFR, and it is manifested by either pathological abnormalities or changes in markers of kidney damage or abnormalities in the imaging tests [18].

This study determines the functional capacity of the kidneys in CKD and determination of GFR using serum creatinine. Sonography is the ideal imaging modality as it is easily available and affordable to provide real-time information on the renal measurements and echogenicity.

Ultrasonographic findings such as longitudinal length, echogenicity, parenchymal and cortical thickness can be affected by chronic kidney disease [19, 20]. The GFR and stage of the disease can be ascertained by the endogenous serum creatinine level [21].

According to O'Neill, the useful upper limit of the normal range for kidney length is said to be 12 cm [20], whereas in our study the mean longitudinal length was 9.7 cm. According to Fiorini and Barozzi, renal length under 8 cm is definitely reduced and should be attributed to chronic renal failure [22]. As renal length decreases with decreasing renal function, renal length has traditionally been considered a surrogate marker of renal function. Hence for progression of disease process, estimation of renal length should be preferred to renal volume.

The mean serum creatinine in our study was 1.25 mg/dl for Grade 0, 1.85 mg/dl for Grade 1, 2.5 mg/dl for Grade 2, 3.27 mg/dl for Grade 3, and 5.03 mg/dl for Grade 4. Mean serum creatinine was significant among echogenicity grades [ANOVA F-Value= 367.726; p=0.0005]. Our study showed a statistically significant correlation between serum creatinine and the grade of echogenicity ($p=0.0005$). This value was similarly seen in a study by Siddappa et al. who also noted a statistically significant correlation between these parameters ($p=0.004$) [23]. Studies by Ibnaieye et al. and Singh A et al. [24] showed similar values ($p=0.001$).

This study showed that the mean parenchymal thickness was significant among echogenicity grades (ANOVA F-value= 31.628; p=0.0005). Similar findings were also obtained in the study by Siddappa et al. There was a statistically significant positive correlation observed between renal echogenicity grading and parenchymal thickness ($p = 0.009$) [25]. As the echogenicity increased, there was a decrease in the mean parenchymal thickness.

This study showed that the mean cortical thickness was significant among echogenicity grades (ANOVA F-value= 477.83; p=0.0005). In our study the mean cortical thickness was found to be 9.3 mm ($p=0.0005$). Similar results were shown in the study by Singh A et al., where mean cortical thickness was found to be 8.5 mm [24]. As the echogenicity increased, there was a decrease in mean cortical thickness.

The limitation of this study is that this study group included more patients with Grade 1 and Grade 2 in comparison to patients with Grade 3 and Grade 4 CKD. This most likely occurred as this study was conducted in a tertiary care hospital, where most cases were treated with renal replacement therapies like hemodialysis, peritoneal dialysis, and renal transplantation, and such patients were excluded from this study.

As serum creatinine increases, the renal cortical echogenicity is increased. Since changes in renal echogenicity are irreversible, a sonological grading of CKD can be carried out, allowing the severity of CKD to be assessed.

Conclusions

The best sonographic parameter that correlates with serum creatinine is renal cortical echogenicity and its grading in comparison to longitudinal length, parenchymal thickness and cortical thickness in patients of CKD. Since renal cortical echogenicity has the advantage of being irreversible in comparison to serum creatinine levels, it can be used as a parameter of renal function.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Levey AS, Coresh J: Chronic kidney disease . The Lancet. 2012, 379:165-80. [10.1016/S0140-6736\(11\)60178-5](https://doi.org/10.1016/S0140-6736(11)60178-5)
- Akbari A, Clase CM, Acott P, et al.: Canadian Society of Nephrology commentary on the KDIGO clinical practice guideline for CKD evaluation and management. Am J Kidney Dis. 2015, 65:177-205.

[10.1053/j.ajkd.2014.10.013](https://doi.org/10.1053/j.ajkd.2014.10.013)

3. Insana MF, Hall TJ, Fishback JL: Identifying acoustic scattering sources in normal renal parenchyma from the anisotropy in acoustic properties. *Ultrasound Med Biol.* 1991, 17:613-26. [10.1016/0301-5629\(91\)90032-R](https://doi.org/10.1016/0301-5629(91)90032-R)
4. Rafique M: Value of routine renal and abdominal ultrasonography in patients undergoing prostatectomy . *Int Urol Nephrol.* 2006, 38:153-6. [10.1007/s11255-005-3830-0](https://doi.org/10.1007/s11255-005-3830-0)
5. Rosansky SJ: Renal function trajectory is more important than chronic kidney disease stage for managing patients with chronic kidney disease. *Am J Nephrol.* 2012, 36:1-0. [10.1159/000339327](https://doi.org/10.1159/000339327)
6. Maojoud O, Ahid S, Cherrah Y: The cost-utility of treating anemia with continuous erythropoietin receptor activator or epoetin versus routine blood transfusions among chronic hemodialysis patients. *Int J Nephrol Renovasc Dis.* 2016, 2016:35-43. [10.2147/IJNRD.S96027](https://doi.org/10.2147/IJNRD.S96027)
7. Rosenfield AT, Siegel NJ: Renal parenchymal disease: histopathologic-sonographic correlation . *Am J Roentgenol.* 1981, 137:793-8. [10.2214/ajr.137.4.793](https://doi.org/10.2214/ajr.137.4.793)
8. Rosenfield AT, Taylor KJ, Crade M, DeGraaf CS: Anatomy and pathology of the kidney by gray scale ultrasound. *Radiology.* 1978, 128:737-44. [10.1148/128.3.737](https://doi.org/10.1148/128.3.737)
9. Levey AS, Becker C, Inker LA: Glomerular filtration rate and albuminuria for detection and staging of acute and chronic kidney disease in adults: a systematic review. *JAMA.* 2015, 313:837-46. [10.1001/jama.2015.0602](https://doi.org/10.1001/jama.2015.0602)
10. Päivänsalo M, Huttunen K, Suramo I: Ultrasonographic findings in renal parenchymal diseases . *Scand J Urol Nephrol.* 1985, 19:119-23. [10.3109/00365598509180238](https://doi.org/10.3109/00365598509180238)
11. Hricak H, Cruz C, Romanski R, et al.: Renal parenchymal disease: sonographic-histologic correlation . *Radiology.* 1982, 144:141-7. [10.1148/radiology.144.1.7089245](https://doi.org/10.1148/radiology.144.1.7089245)
12. Lamont AC, Graebe AC, Pelmore JM, Thompson JR: Ultrasound assessment of renal cortical brightness in infants: is naked eye evaluation reliable? . *Invest Radiol.* 1990, 25:250-3.
13. Eggert P, Debus F, Kreller-Laugwitz G, Oppermann HC: Densitometric measurement of renal echogenicity in infants and naked eye evaluation: a comparison. *Pediatr Radiol.* 1991, 21:111-3.
14. Manley JA, O'Neill WC: How echogenic is echogenic? Quantitative acoustics of the renal cortex . *Am J Kidney Dis.* 2001, 37:706-11. [10.1016/S0272-6386\(01\)80118-9](https://doi.org/10.1016/S0272-6386(01)80118-9)
15. Galbraith LE, Ronksley PE, Barnieh LJ, et al.: The see kidney disease targeted screening program for CKD . *Clin J Am Soc Nephrol.* 2016, 11:964-972. [10.2215/CJN.11961115](https://doi.org/10.2215/CJN.11961115)
16. Cheong B, Muthupillai R, Rubin MF, Flamm SD: Normal values for renal length and volume as measured by magnetic resonance imaging. *Clin J Am Soc Nephrol.* 2007, 2:38-45. [10.2215/CJN.00930306](https://doi.org/10.2215/CJN.00930306)
17. Mazzotta L, Sarteschi LM, Carlini A, Antonelli A: Comparison of renal ultrasonographic and functional biometry in healthy patients and in patients with chronic renal failure. [Article in Italian]. *Arch Ital Urol Androl.* 2002, 74:206-9.
18. Bailie GR, Uhlig K, Levey AS: Clinical practice guidelines in nephrology: evaluation, classification, and stratification of chronic kidney disease. *Pharmacotherapy.* 2005, 25:491-502. [10.1592/phco.25.4.491.61034](https://doi.org/10.1592/phco.25.4.491.61034)
19. O'Neill WC: Atlas of Renal Ultrasonography . WB Saunders Company, Philadelphia, USA; 2001.
20. O'Neill WC: Sonographic evaluation of renal failure . *Am J Kidney Dis.* 2000, 35:1021-38. [10.1016/S0272-6386\(00\)70036-9](https://doi.org/10.1016/S0272-6386(00)70036-9)
21. Burtis CA, Ashwood ER: Tietz Textbook of Clinical Chemistry. American Association for Clinical Chemistry, 1997.
22. Fiorini F, Barozzi L: The role of ultrasonography in the study of medical nephropathy . *J Ultrasound.* 2007, 10:161-7. [10.1016/j.jus.2007.09.001](https://doi.org/10.1016/j.jus.2007.09.001)
23. Siddappa JK, Singla S, Mohammed Al Ameen SC, Kumar N: Correlation of ultrasonographic parameters with serum creatinine in chronic kidney disease. *J Clin Imaging Sci.* 2013, 3:1-6.
24. Singh A, Gupta K, Chander R, Vira M: Sonographic grading of renal cortical echogenicity and raised serum creatinine in patients with chronic kidney disease. *J Evolution Med Dent Sci.* 2016, 5:2279-2286. [10.1426/jemds/2016/550](https://doi.org/10.1426/jemds/2016/550)
25. Platt JF, Rubin JM, Bowerman RA, Marn CS: The inability to detect kidney disease on the basis of echogenicity. *Am J Roentgenol.* 1988, 151:317-9. [10.2214/ajr.151.2.317](https://doi.org/10.2214/ajr.151.2.317)