Comparison of the Efficacy of Tramadol and Paroxetine in the Management of Premature Ejaculation

Muhammad Fazal Ur Rehman¹, Ali Imran Zaidi², Tanveer Ul haq², Shoaib Rafique², Farman Ali³

1. Urology, Bakhtawar Amin Medical and Dental College, Multan, PAK 2. Urology, Multan Institute of Kidney Diseases, Multan, PAK 3. Internal Medicine, District Headquarter Hospital, Khanewal, PAK

Corresponding author: Farman Ali, chfarmanali358@gmail.com

Abstract

Objective

The goal of this study was to compare the efficacy of tramadol and paroxetine in the treatment of primary premature ejaculation (PE).

Study design

This study was a randomized controlled trial performed in the outpatient department of Nishtar Hospital, Multan, from January 2017 to January 2018.

Methodology

One hundred six patients were diagnosed with PE and included in the study. The patients were categorized into two groups receiving either tramadol or paroxetine through a lottery randomization method. The main variables were baseline PE, baseline satisfaction after intercourse, baseline intravaginal ejaculatory latency time (IELT), ejaculation control, difficulty in ejaculation, and after-treatment satisfaction with sexual intercourse and IELT. We used IBM SPSS Statistics for Windows, Version 23.0 (Armonk, NY: IBM Corp.) for data analysis, and $p \leq 0.05$ was considered statistically significant.

Results

Ejaculation control, difficulty in ejaculation, and distress due to ejaculation in patients in the tramadol group was noted as 24.5%, 7.5%, and 7.5%, respectively. Ejaculation control, difficulty in ejaculation, and distress due to ejaculation in the paroxetine group was noted as 49.1%, 17%, and 24.5%, respectively. The differences were statistically significant within the groups at baseline and after treatment of PE ($p < 0.001$).

Conclusion

Tramadol is an effective and useful drug as compared to paroxetine for the treatment of PE. Tramadol can be used as an alternative to other medications for the treatment of lifelong PE.

Categories: Urology

Keywords: tramadol, paroxetine, premature ejaculation, latency time.

Introduction

In mammals, a trend has been observed with regards to ejaculating quickly for the purpose of reproducing with multiple partners in order to share genetic material. In *Homo sapiens*, having multiple sexual intercourses is not only limited to reproductive purposes but also has important effects on quality of life. As males reach puberty, they start to engage in sexual intercourse, which alters their ejaculation time and perceived ejaculation control with the passage of time. Therefore, premature ejaculation (PE) should not be considered or defined as pathological [1].

Delay in ejaculation with every intercourse can lead to negative consequences such as frustration, distress, and avoidance of sexual contact [2]. Ejaculation within one minute of vaginal penetration in every intercourse since an individual's first sexual episode is considered lifelong PE, which leads to a reduction in latency time, and more often, PE occurs when ejaculation is acquired in three minutes or sooner [3].

The management of PE remains a challenge for clinicians, patients, and their partners. The prevalence of PE is around 20% to 40% worldwide, and its diagnosis is often complicated [4]. The definition of PE frequently relies on ill-defined subjective descriptions, and as little as 5% of individuals fulfill the definition of PE according to the International Society of Sexual Medicine (ISSM) criteria [5]. Determining PE represents a

Received 03/30/2020
Review began 04/17/2020
Review ended 09/26/2020
Published 09/29/2020

© Copyright 2020

Ur Rehman et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Ur Rehman M, Imran Zaidi A, Ul Haq T, et al. (September 29, 2020) Comparison of the Efficacy of Tramadol and Paroxetine in the Management of Premature Ejaculation. Cureus 12(9): e10725. DOI 10.7759/cureus.10725

challenge in that PE can be found in a person with frequent sexual interaction and rapid ejaculation or a person with infrequent sexual interaction and a single episode of PE.

Psychological or relationship problems, sexual performance anxiety, hyperthyroidism, acute or chronic prostatitis, diabetes mellitus, hypertension, and high body mass index (BMI) are causes of acquired PE [6]. However, lifelong PE is related to a complex interplay of central and peripheral dopaminergic, endocrinological, serotonergic, epigenetic, oxytocinergic, and genetic factors [7]. In the treatment of PE, tramadol and sildenafil are not recommended. However, paroxetine and dapoxetine (selective serotonin reuptake inhibitors [SSRIs]), are recommended by the ISSM guidelines [8,9]. Tramadol could improve intravaginal ejaculatory latency time (IELT) and promote satisfaction level, but ideal dosing remains to be determined. Tramadol is well tolerated at a dose of 25 to 50 mg [10,11].

Therefore, we conducted this randomized controlled trial to compare the efficacy of tramadol and paroxetine in the treatment of primary PE.

Materials And Methods

This randomized controlled trial was conducted at the outpatient department of Nishtar Hospital, Multan, from January 2017 to January 2018. All patients were diagnosed based on history, physical examination, and a dedicated validated questionnaire. Proper approval was obtained from the ethics committee and institutional review board. Potent married men aged 30 to 40 years who had been in a relationship for at least six months with uncontrolled ejaculation within one minute of vaginal intromission were included in the study. Patients with low libido, erectile dysfunction, psychiatric disorders, or psychological conditions such as depression, alcohol abuse, use of sedative drugs, or having organic illnesses like hyperthyroidism, asthma, hypothyroidism, diabetes, and cardiac arrhythmias which would limit the use of SSRIs were excluded from the study. Patients with any history of previous use of treatments for PE within the three months from the presentation were excluded as well. IELT was defined as the time from vaginal intromission to intravaginal ejaculation. The confidence interval was set at 95%, and the power of the study at 80%. P1 (i.e., control over ejaculation in the tramadol group from baseline to post-treatment) was 8.4%. P2 (i.e., control over ejaculation in the tramadol group from baseline to post-treatment) was 22%. A total of 106 patients were included in this study. Fifty-three patients were enrolled in the tramadol group and 53 in the paroxetine group.

Patients who fulfilled the inclusion criteria were randomly assigned into two groups: Group A and Group B.

Patients in Group A received 50 mg of tramadol two hours before planned intercourse. Group B patients received 20 mg of paroxetine four hours before planned intercourse once a day, thrice weekly for eight weeks. Informed consent was obtained before inclusion in the study. All patients were monitored via follow-up for eight weeks from the start of treatment on weekly basis.

Data were entered and analyzed using IBM SPSS Statistics for Windows, Version 23.0 (Armonk, NY: IBM Corp.). Mean and standard deviation, as well as frequency and percentages, were calculated for numerical data like age, duration of disease, gender, socioeconomic status, and IELT (IELT was noted on a stopwatch). A p-value ≤ 0.05 was considered statistically significant.

Results

A total of 106 patients were included in the study; 53 patients were randomized to the tramadol group and 53 to the paroxetine group. The mean age of patients in the tramadol group was 31.8 ± 3.8 years, and the mean age of patients in the paroxetine group was 32.2 ± 4.1 years, respectively. The mean baseline IELT for patients in the tramadol group versus paroxetine group was 44.93 ± 3.95 seconds and 43.74 ± 2.14 seconds, respectively, with no statistically significant difference ($p=0.057$). The mean treatment IELT for patients in the tramadol and paroxetine groups was 136.91 ± 5.61 seconds versus 95.42 ± 5.53 seconds, respectively, denoting a statistically significant difference ($p=0.000$). The differences were statistically significant within the groups' baseline and treatment IELT ($p<0.001$; Table 1).

Variables	Tramadol n=53	Paroxetine n=53	P-value
Age (years)	31.8±3.8	32.2±4.1	0.426
Baseline IELT (seconds)	44.93±3.95	43.74±2.14	0.057
Treatment IELT (seconds)	136.91±5.61	95.42±5.53	0.000
Paired sample test p-value	0.000	0.000	--

TABLE 1: Intravaginal ejaculatory latency time in both groups

The mean baseline PE score of patients in the tramadol and paroxetine groups was 9.68±2.23 versus 9.15±1.05, respectively, with no significant difference (p=0.117). Ejaculation control, difficulty in ejaculation, and distress due to ejaculation in the tramadol group were noted in 26 (49.1%), 4 (7.5%), and 2 (3.8%), and 9 (17%) patients, respectively. Ejaculation control, difficulty in ejaculation, and distress due to ejaculation in the paroxetine group were noted in 33 (62.3%), 20 (37.7%), 7 (13.2%), and 20 (37.7%) patients respectively.

Meanwhile, the mean post-treatment PE score in the tramadol and paroxetine groups was 14.01±1.31 versus 9.82±1.96, respectively, a statistically significant difference (p=0.000). Ejaculation control, difficulty in ejaculation, and distress due to ejaculation in the tramadol group were noted in 13 (24.5%), 4 (7.5%), and 4 (7.5%) patients, respectively. Ejaculation control, difficulty in ejaculation, and distress due to ejaculation in the paroxetine group were noted in 26 (49.1%), 9 (17%), and 13 (24.5%), respectively. The differences were statistically significant between the groups' baseline and post-treatment PE (p<0.001; Table 2).

Variables	Tramadol n=53	Paroxetine n=53	P-value
Baseline PE	9.68±2.23	9.15±1.05	0.117
Treatment PE	14.01±1.31	9.82±1.96	0.000
Paired sample test p-value	0.000	0.036	--
Baseline satisfaction with sexual intercourse			
Control over ejaculation	n=26 (49.1%)	n=33 (62.3%)	0.171
Difficulty	n=2 (3.8%)	n=7 (13.2%)	0.081
Ejaculation-related distress	n=9 (17%)	n=20 (37.7%)	0.017
After treatment satisfaction with sexual intercourse			
Control over ejaculation	n=13 (24.5%)	n=26 (49.1%)	0.009
Difficulty	n=4 (7.5%)	n=9 (17%)	0.139
Ejaculation-related distress	n=4 (7.5%)	n=13 (24.5%)	0.017

TABLE 2: Premature ejaculation profile score in both groups

Discussion

In our study, we observed a marked improvement in PE with the use of tramadol compared to paroxetine. A similar study was conducted by Hamidi-Madani et al., reporting that tramadol is an effective and safe alternative to paroxetine for the treatment of PE [12]. That study included three groups of patients who received tramadol, paroxetine, or placebo. A similar study was conducted by Zhang et al. and concluded that paroxetine is more effective and gave long-lasting results when used for the purpose of PE, but paroxetine in combination is more effective [13].

A study conducted by Wu et al. concluded that tramadol and paroxetine had no significant difference in terms of mean IELT; it also suggested that tramadol can be used as a treatment option for PE [14]. Safarinejad and Hosseini reported that tramadol can be used as PE treatment option and that it can increase the IELT when used for six weeks or more [15]. Both studies have findings that are consistent with ours.

Another study conducted by Kaynar et al. reported that similar findings regarding the use of tramadol [16]. Compared to placebo, after eight weeks of treatment of PE with tramadol, a significant increase in mean IELT was noted. In another study by Alghobary et al., controversial results were seen that at sixth-week treatment both paroxetine and tramadol increase the IELT 11 fold and seven fold respectively, but at 12-week treatment, tramadol decreases the IELT by five fold but paroxetine increases it 22 folds [17].

Bar-Or et al. compared two different doses of tramadol (62 mg and 89 mg) versus placebo for 12 weeks and observed an increase in IELT in all groups, with tramadol 62 mg being more effective, given that a 4.2-fold significant increase from baseline was observed [18]. Ozcan et al. reported controversial findings suggesting that paroxetine increased all four indicators of PE [19].

Patrick et al. reported increase in all four items of PE when tramadol was used as an alternative to all other medications [20]. Salem et al. reported that "on demand" use of tramadol for six weeks increased mean IELT [21]. Collectively, these results allow medical practitioners to consider the use of tramadol for PE treatment.

Limitations

Assessment of all variables was dependent on self-estimation of patient using stopwatch which has some degree of bias regarding the exact time. Further studies are needed to collect observations with some strong and more reliable methods.

Conclusions

Tramadol is effective and useful compared to paroxetine in the treatment of PE. Tramadol can be used as an alternative to other medications for the treatment of lifelong PE.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. Bakhtawar Amin Medical and Dental College issued approval 28/URO/BMDC-2018. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Puppo V, Puppo G: Comprehensive review of the anatomy and physiology of male ejaculation: premature ejaculation is not a disease. *Clin Anat*. 2016, 29:111-119. [10.1002/ca.22655](#)
2. Serefoglu EC, McMahon CG, Waldinger MD, et al.: An evidence-based unified definition of lifelong and acquired premature ejaculation: report of the second International Society for Sexual Medicine Ad Hoc Committee for the Definition of Premature Ejaculation. *J Sex Med*. 2014, 2:41-59. [10.1002/sm2.27](#)
3. McCabe MP, Sharlip ID, Atalla E, et al.: Definitions of sexual dysfunctions in women and men: a consensus statement from the fourth International Consultation on Sexual Medicine 2015. *J Sex Med*. 2015, 13:135-143. [10.1016/j.jsxm.2015.12.019](#)
4. Kalejaiye O, Almekaty K, Blecher G, Minhas S: Premature ejaculation: challenging new and the old concepts. *F1000Res*. 2017, 6:2084. [10.12688/f1000research.12150.1](#)
5. Althof SE, McMahon CG, Waldinger MD, et al.: An update of the International Society of Sexual Medicine's Guidelines for the Diagnosis and Treatment of Premature Ejaculation (PE). *Sex Med*. 2014, 2:60-90. [10.1002/sm2.28](#)
6. McMahon CG, Lee G, Park JK, Adaikan PG: Premature ejaculation and erectile dysfunction prevalence and attitudes in the Asia-Pacific region. *J Sex Med*. 2012, 9:454-465. [10.1111/j.1743-6109.2011.02507.x](#)
7. Waldinger MD, McIntosh J, Schweitzer DH: A five-nation survey to assess the distribution of the intravaginal ejaculatory latency time among the general male population. *J Sex Med*. 2009, 6:2888-2895. [10.1111/j.1743-6109.2009.01392.x](#)
8. Kirby EW, Carson CC, Coward RM: Tramadol for the management of premature ejaculation: a timely systematic review. *Int J Impot Res*. 2015, 27:121-127. [10.1038/ijir.2015.7](#)
9. Montorsi F, Adaikan G, Becher E, et al.: Summary of the recommendations on sexual dysfunctions in men. *J Sex Med*. 2010, 7:3572-3588. [10.1111/j.1743-6109.2010.02062.x](#)
10. Gameel TA, Tawfik AM, Abou-Farha MO, Bastawisy MG, El-Bendary MA, El-Gamasy AEN: On-demand use of tramadol, sildenafil, paroxetine and local anaesthetics for the management of premature ejaculation: a randomised placebo-controlled clinical trial. *Arab J Urol*. 2013, 11:392-397. [10.1016/j.aju.2013.05.003](#)
11. Janssen PK, van Schaik R, Zwinderman AH, Olivier B, Waldinger MD: The 5-HT_{1A} receptor C(1019)G polymorphism influences the intravaginal ejaculation latency time in Dutch Caucasian men with lifelong premature ejaculation. *Pharmacol Biochem Behav*. 2014, 121:184-188. [10.1016/j.pbb.2014.01.004](#)
12. Hamidi-Madani A, Motiee R, Mokhtari G, Nasseh H, Esmaeili S, Kazemnezhad E: The efficacy and safety of on-demand tramadol and paroxetine use in treatment of life long premature ejaculation: a randomized

- double-blind placebo-controlled clinical trial. *J Reprod Infertil*. 2018, 19:10-15.
13. Zhang D, Cheng Y, Wu K: Paroxetine in the treatment of premature ejaculation: a systematic review and meta-analysis. *BMC Urol*. 2019, 19:2.
14. Wu T, Yue X, Duan X, Luo D, Cheng Y, Tian Y, Wang K: Efficacy and safety of tramadol for premature ejaculation: a systematic review and meta-analysis. *Urology*. 2012, 80:618-624. [10.1016/j.urology.2012.05.035](https://doi.org/10.1016/j.urology.2012.05.035)
15. Safarinejad MR, Hosseini SY: Safety and efficacy of tramadol in the treatment of premature ejaculation: a double-blind, placebo-controlled, fixed-dose, randomized study. *J Clin Psychopharmacol*. 2006, 26:27-31. [10.1097/01.jcp.0000195110.79027.3f](https://doi.org/10.1097/01.jcp.0000195110.79027.3f)
16. Kaynar M, Kilic O, Yurdakul T: On-demand tramadol hydrochloride use in premature ejaculation treatment. *Urology*. 2012, 79:145-149. [10.1016/j.urology.2011.09.031](https://doi.org/10.1016/j.urology.2011.09.031)
17. Alghobary M, El-Bayoumy Y, Mostafa Y, Mahmoud EHM, Amr M: Evaluation of tramadol on demand vs. daily paroxetine as a long-term treatment of lifelong premature ejaculation. *J Sex Med*. 2010, 7:2860-2867. [10.1111/j.1743-6109.2010.01789.x](https://doi.org/10.1111/j.1743-6109.2010.01789.x)
18. Bar-Or D, Salottolo KM, Orlando A, Winkler JV, Tramadol ODT Study Group: A randomized double-blind, placebo-controlled multicenter study to evaluate the efficacy and safety of two doses of the tramadol orally disintegrating tablet for the treatment of premature ejaculation within less than 2 minutes. *Eur Urol*. 2012, 61:736-743. [10.1016/j.eururo.2011.08.039](https://doi.org/10.1016/j.eururo.2011.08.039)
19. Ozcan L, Polat EC, Otunctemur A, Ozbek E: Duloxetine, dual serotonin and norepinephrine reuptake inhibitor, versus paroxetine, selective serotonin reuptake inhibitor, in the treatment for premature ejaculation. *Int Urol Nephrol*. 2015, 47:283-287. [10.1007/s11255-014-0905-9](https://doi.org/10.1007/s11255-014-0905-9)
20. Patrick DL, Althof SE, Pryor JL, et al.: Original research—ejaculatory disorders: premature ejaculation: an observational study of men and their partners. *J Sex Med*. 2005, 2:358-367. [10.1111/j.1743-6109.2005.20353.x](https://doi.org/10.1111/j.1743-6109.2005.20353.x)
21. Salem AM, Kamel II, Rashed LA, GamalEl Din SF: Effects of paroxetine on intravaginal ejaculatory latency time in Egyptian patients with lifelong premature ejaculation as a function of serotonin transporter polymorphism. *Int J Impot Res*. 2017, 29:7-11. [10.1038/ijir.2016.36](https://doi.org/10.1038/ijir.2016.36).