A Complicated Case of Vertebral Osteomyelitis by *Serratia Marcescens*

Abdul Rana¹, Noor Ul Ann Rabbani¹, Samuel Wood¹, Colin McCorkle¹, Christine Gilkerson¹

1. Internal Medicine, Marshall University Joan C. Edwards School of Medicine, Huntington, USA

Corresponding author: Abdul Rana, ranaa@marshall.edu

Abstract

Osteomyelitis is an infection of the bone and bone marrow that can be caused by an exogenous source or by hematogenous spread. The hematogenous spread of bacteria usually results in bacteremia with subsequent infection of the bone. The most commonly associated pathogen with this type of infection is *Staphylococcus aureus*, with other gram-negative organisms, such as *Pseudomonas aeruginosa*, also frequently encountered. The microorganism *Serratia marcescens* is a rare and infrequently encountered cause of this condition known to cause nosocomial infections. This organism can be notoriously difficult to treat, with resistance to many commonly used antibiotics. The case presented is one of vertebral osteomyelitis in an intravenous drug user caused by *Serratia marcescens* with subsequent treatment and management of the condition. This case allows for investigation into the continued management of intravenous drug user infections, with the isolation and treatment of less commonly encountered pathogens.

Categories: Internal Medicine, Pathology, Infectious Disease

Keywords: vertebral osteomyelitis, *serratia marcescens*, hematogenous, intravenous drug abuse

Introduction

Osteomyelitis is an infection localized to the bone. The most common pathogens associated with this infection are *Staphylococcus aureus* and aerobic gram-negative bacilli. The causes of osteomyelitis can be divided into pyogenic and non-pyogenic, and the most common organisms in each group are *Staphylococcus aureus* and *Mycobacterium tuberculosis*, respectively [1]. Osteomyelitis can also be divided based on the mode of spread into the hematogenous and non-hematogenous routes. *Serratia* species, although well-known, are not common pathogens associated with either classification of osteomyelitis. *Serratia* is a gram-negative, motile, facultative anaerobic bacillus of the Enterobacteriaceae group. These bacteria are known to cause a spectrum of clinical diseases in humans most often related to indwelling catheters or respiratory infections but can include other infections such as osteomyelitis, meningitis, or septic arthritis. *Serratia marcescens* has been a well-known cause of severe infections in humans since the 1960s [2]. Currently, it accounts for approximately 2% of nosocomial infections [3]. We present a case of vertebral osteomyelitis caused by *Serratia marcescens* in an immunocompetent intravenous (IV) drug user and the subsequent treatment and management of his condition.

Case Presentation

A 29-year-old male with a history of IV drug use, without any known comorbidities presented to the hospital with complaints of neck and back pain for three weeks. He denied any fever or chills, any radiation of pain, or any numbness or tingling. He was seen earlier in the emergency room during that week, given a shot of IV ketorolac for pain, and then discharged. The patient's physical exam was unremarkable for any focal neurological deficits. His illness did not subside, the pain worsened, and he subsequently presented later in the week with worsening neck and back pain. He also complained of an abscess on his right leg, located above the lateral malleolus with clean edges that he first noticed two to three weeks ago. At the time of presentation, he was alert and oriented, and vitals were within normal limits. The computed tomography (CT) scan revealed osteomyelitis at the cervical vertebra (C4-C5) level and prevertebral fluid collection. Neurosurgery was consulted, and magnetic resonance imaging (MRI) was recommended, which revealed C4-C5 osteomyelitis with an epidural abscess. The patient was started on IV vancomycin and admitted to the floor. During his hospital stay, neurosurgery performed an anterior corpectomy and posterior cervical instrumented fusion. The procedure was complicated by a cerebrospinal fluid (CSF) leak, and intraoperative cultures grew *Serratia marcescens*, as shown in Figure 1. The patient's hospital course was complicated by the CSF leak and drainage from the surgical wound, for which he had to be taken back to the operating room (OR) for an anterior washout. A lumbar drain was placed that day for drainage. Infectious disease recommended adding aztreonam 2 g IV every eight hours for six weeks. Vancomycin was discontinued after the cultures did not grow methicillin-resistant *Staphylococcus aureus* (MRSA). Blood cultures were also negative during his hospital stay. The patient completed six weeks of treatment with IV aztreonam and was discharged home after completion of the antibiotics course. He was followed up in the clinic where his condition has significantly improved and he did not have any active complaints during that visit.

Received 05/29/2020
Review began 06/16/2020
Review ended 06/28/2020
Published 07/04/2020

© Copyright 2020

Rana et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Rana A, Rabbani N, Wood S, et al. (July 04, 2020) A Complicated Case of Vertebral Osteomyelitis by *Serratia Marcescens*. Cureus 12(7): e9002. DOI 10.7759/cureus.9002

FIGURE 1: Cultures isolated from the epidural abscess grew *Serratia marcescens* on blood agar

Medium-sized colonies appearing buff in color. Bacteria produce red pigment.

Discussion

Serratia is an infrequent cause of osteomyelitis that is usually seen in immunocompromised patients [3]. *Serratia marcescens* is a motile, gram-negative bacillus from the Enterobacteriaceae family, with natural red pigment. It has been associated with nosocomial opportunistic infection and intravenous drug abuse (IVDA) [2,4]. Although frequently known for its pigment production, one study found the non-pigment producing strains to be most often isolated in the hospital setting, mainly infecting immunocompromised hosts [5]. *Serratia marcescens* was discovered by an Italian pharmacist in 1819 and produces beta-lactamase that can make it challenging to treat. Some of the risk factors associated with this infection are due to instrumentation or catheterization. *Serratia* has various virulent features that include hemolysin production, swarming, and biofilm production, which are regulated by RssAB signaling and temperature [4]. Although this particular species of *Serratia* is a rare cause of osteomyelitis, it is not unheard of even in a patient with no predisposing factors [5]. Vertebral osteomyelitis is a rare form of osteomyelitis, accounting for 2%-7% of all cases, and has an incidence rate of 2.4 cases per 100,000 [1]. Historically, the classic pathogen associated with pyogenic vertebral osteomyelitis is *Staphylococcus aureus*, both the methicillin-resistant and methicillin-sensitive strains.

Our patient likely developed hematogenous osteomyelitis by direct inoculation associated with IV drug use, which then subsequently spread to the C4-C5 area of the spine [2]. He had a history of IVDA, which predisposed him to this particular infection. Bacterial cultures are essential to form a diagnosis. *Serratia* resides in water, making hospital sinks a potential source, and has also been known to be spread by healthcare workers [5]. There is widespread antibiotic resistance to *Serratia marcescens* due to the production of beta-lactamase [3,6]. The drug of choice should be tailored according to the site of infection and the specificity and sensitivity of the organism to a particular antibiotic. *Serratia* has been susceptible to numerous antibiotics such as third and fourth generation cephalosporins, aztreonam, carbapenem, and trimethoprim-sulfamethoxazole. *Serratia* has been found to be resistant to colistin and cephalothin [7].

The fact that this patient had a history of Intravenous drug abuse predisposed him to vertebral osteomyelitis, but the causative organism being *Serratia* is not entirely unknown but somewhat unusual.

There is limited data regarding the prevalence of community-acquired *Serratia* infections. Still, there have been well-documented reservoirs of *Serratia* in the hospital setting, along with outbreaks of *Serratia* related to a contaminated source [8]. A study from Blossom et al. found *Serratia* sources in saline solutions and pre-filled heparin syringes [9]. A survey from Bopp DJ et al. found a *Serratia* outbreak related to contaminated IV magnesium sulfate [10]. These studies could suggest a similar possible growth to our presented case, possibly that the current patient had a source of *Serratia* on the IV syringe for drug abuse.

Conclusions

The case presented offers a unique opportunity to explore the relationship between IV drug use and osteomyelitis with a rare pathogen. This study could be extended further by analyzing other, related infectious sources of *Serratia marcescens* to look for shared epidemiological features.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Martins HF, Raposo A, Baptista I, Almeida J: *Serratia marcescens* osteomyelitis in Cushing's disease. *BMJ Case Rep.* 2015, 2015:bcr2015212872. [10.1136/bcr-2015-212872](https://doi.org/10.1136/bcr-2015-212872)
2. Khanna A, Khanna M, Aggarwal A: *Serratia marcescens*- a rare opportunistic nosocomial pathogen and measures to limit its spread in hospitalized patients. *J Clin Diagn Res.* 2013, 7:243-246. [10.7860/JCDR/2013/5010.2737](https://doi.org/10.7860/JCDR/2013/5010.2737)
3. Yu VL: *Serratia marcescens*: historical perspective and clinical review. *N Engl J Med.* 1979, 300:887-893. [10.1056/NEJM197904193001604](https://doi.org/10.1056/NEJM197904193001604)
4. Lin CS, Horng JT, Yang CH, et al.: RssAB-FlhDC-ShlBA as a major pathogenesis pathway in *Serratia marcescens*. *Infect Immun.* 2010, 78:4870-4881. DOI: [10.1128/IAI.00661-10](https://doi.org/10.1128/IAI.00661-10)
5. Hadid H, Usman M, Thapa S: Severe osteomyelitis and septic arthritis due to *Serratia marcescens* in an immunocompetent patient. *Case Rep Infect Dis.* 2015, 2015:347652. [10.1155/2015/347652](https://doi.org/10.1155/2015/347652)
6. Empel J, Baraniak A, Literacka E, et al.: Molecular survey of beta-lactamases conferring resistance to newer beta-lactams in Enterobacteriaceae isolates from Polish hospitals. *Antimicrob Agents Chemother.* 2008, 52:2449-2454. [10.1128/AAC.00043-08](https://doi.org/10.1128/AAC.00043-08)
7. Dowzicky MJ, Park CH: Update on antimicrobial susceptibility rates among gram-negative and gram-positive organisms in the United States: results from the Tigecycline Evaluation and Surveillance Trial (TEST) 2005 to 2007. *Clin Ther.* 2008, 30:2040-2050. [10.1016/j.clinthera.2008.11.006](https://doi.org/10.1016/j.clinthera.2008.11.006)
8. Corrah TW, Enoch DA, Aliyu SH, Lever AM: Bacteraemia and subsequent vertebral osteomyelitis: a retrospective review of 125 patients. *QJM.* 2011, 104:201-207. [10.1093/qjmed/hcq178](https://doi.org/10.1093/qjmed/hcq178)
9. Blossom D, Noble-Wang J, Su J, et al.: *Serratia* in Prefilled Syringes Investigation Team Group. Multistate outbreak of *Serratia marcescens* bloodstream infections caused by contamination of prefilled heparin and isotonic sodium chloride solution syringes. *Arch Intern Med.* 2009, 169:1705-1711. [10.1001/archinternmed.2009.290](https://doi.org/10.1001/archinternmed.2009.290)
10. Bopp DJ, Jernigan DB, Kazakova S, Bresnitz EA, Tan CG, McDonald LC: A multistate outbreak of *Serratia marcescens* bloodstream infection associated with contaminated intravenous magnesium sulfate from a compounding pharmacy. *Clin Infect Dis.* 2007, 45:527-533.