Pattern and Outcome of Thoracic Injuries in a Busy Tertiary Care Unit

Misauq Mazcuri¹, Tanveer Ahmad¹, Ambreen Abid¹, Pratikshya Thapaliya¹, Mansab Ali², Nadir Ali¹

1. Thoracic Surgery, Jinnah Postgraduate Medical Centre, Karachi, PAK 2. General Surgery, Jinnah Postgraduate Medical Centre, Karachi, PAK

Corresponding author: Misauq Mazcuri, misauq229@gmail.com

Abstract

Introduction: Thoracic traumas are one of the most commonly encountered injuries in the emergency room. They range from blunt chest injuries due to road traffic accidents to penetrating chest injuries. Immediate medical and surgical interventions are essential to improve the outcome. This study was conducted to assess the pattern of thoracic trauma presenting to the emergency room, their outcome and factors contributing to it.

Methods: This prospective, observational, cross-sectional study was conducted in the Department of Thoracic Surgery, Jinnah Post Graduate Medical Center, Karachi from January 1 until July 31, 2020, with institutional ethical approval. Patients age ≥ 12 years presenting with traumatic thoracic injury with or without associated injuries were included. Characteristics of their injuries and management outcome were studied. All data was processed through Statistical Package for Social Sciences (SPSS) Statistics version 22 (IBM Corp., Armonk, NY, USA).

Results: A total of 199 patients were included; 154 (77.4%) patients were male and 45 (22.6%) patients were female. The most common age group presenting with trauma was the middle age (30-60 years), which included 101 (50.8%) patients. Out of the total, 126 (63.3%) had blunt chest injuries and 73 (36.6%) had penetrating chest injuries. Road traffic accidents were the most common cause of blunt chest injuries seen in 83 (65.8%) patients, whereas gunshot was the most common mode of penetrating chest injuries encountered in 41 (56.2%) cases. Tube thoracostomies were performed in 166 (83.4%) patients and thoracotomies in seven (3.51%) patients. Out of the total, 57 (28.6%) patients required mechanical ventilation and it was associated with blunt trauma, hemothorax, rib fracture, abdominal and head injuries ($p \leq 0.05$). Mortality was seen in 22 (11.1%), which was associated with hemothorax, head injuries, mechanical ventilation and severe blood loss ($p \leq 0.05$).

Conclusion: Traumatic thoracic injuries are a preventable cause of mortality. Blunt chest injuries are more common than penetrating chest injuries. Proper implementation of public safety measures ensures less frequent and severe outcomes. Emergency department team and specialized thoracic surgeons must come together to manage these critical patients with utmost care.

Review began 10/15/2020

Review ended 10/20/2020

Published 10/26/2020

© Copyright 2020

Mazcuri et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Categories: Cardiac/Thoracic/Vascular Surgery, Pulmonology, Trauma

Keywords: thoracic trauma, blunt chest injuries, penetrating chest injuries, hemothorax, tube thoracostomies, mortality

Introduction

Trauma is among the most common reasons for hospitalization across the world. As many as two-thirds of patients of trauma have chest trauma [1]. Thoracic trauma may range from a rib fracture to life-threatening injury involving the heart or great vessels. After head injury, thoracic trauma is the second leading cause of death [1]. In the United States, 35% of all trauma-related mortalities are as a result of thoracic trauma alone or in association with other non-thoracic injuries [2].

Blunt chest injuries (BCI) are more common than penetrating and account for 20-25% of all thoracic trauma-related mortalities [2]. One of the most important causes of blunt chest injuries is road traffic accidents (RTAs), especially in developing countries [3]. RTAs constitute almost 80% of all cases of BCI. Other causes include a fall from height, assault, and occupational injuries [2]. BCI is critical and can lead to acute respiratory failure and death due to airway obstruction and lung injury [4]. Penetrating chest injuries (PCIs), although less common, can be acutely fatal. Immediate intervention is critical in PCI in order to prevent mortality. PCIs are mostly a result of assault including stab wounds, gunshot, and occupational mishaps [5].

In the Pakistani population, traumatic thoracic injuries are more common among the younger age group (21-50 years). The commonest thoracic injuries include rib fractures, with head injuries contributing the common associated non-thoracic injuries [6]. Poor outcome in patients with thoracic trauma is predicated by advanced age [2], hemothorax [7], need for mechanical ventilation (MV) [7], and associated non-thoracic

How to cite this article

Mazcuri M, Ahmad T, Abid A, et al. (October 26, 2020) Pattern and Outcome of Thoracic Injuries in a Busy Tertiary Care Unit. Cureus 12(10): e11181. DOI 10.7759/cureus.11181

injuries [8,9].

The study is conceived to quantify the burden of thoracic injuries in a large tertiary care hospital and evaluate the factors responsible for morbidity and mortality in thoracic injuries.

Materials And Methods

A prospective, observational, cross-sectional study was conducted in the Department of Thoracic Surgery, Jinnah Post Graduate Medical Center, Karachi. The duration of this study was from January 1 until July 31, 2020. The study was conducted after ethical approval from the Institutional Review Board. Informed consent was obtained from all study participants and their guardians.

Patients age 12 years or above, presenting with traumatic injury to the thorax with or without other associated injuries, were included in the study. All patients were assessed and managed according to the Advanced Trauma Life Support (ATLS) protocol. It was followed by radiological evaluation such as X-ray, Focused Assessment with Sonography in Trauma (FAST), or computed tomography (CT) scan of the brain as indicated. In the case of associated non-thoracic injuries, a multidisciplinary approach was followed. Patients were stabilized within the emergency room. Tube thoracostomies and thoracotomies were performed as indicated. Mechanical ventilation was started if required.

Patient data were collected on a questionnaire comprised of demographic information of the patient, characteristics of injury, primary diagnosis, associated thoracic and non-thoracic injuries, medical and surgical interventions, and outcome of the injury. All data was processed through Statistical Package for Social Sciences (SPSS) Statistics version 22 (IBM Corp., Armonk, NY, USA). For categorical data, frequencies and percentages were calculated. Chi-square was applied to determine statistical correlation. For continuous data, mean and standard deviation (SD) was calculated. Independent T-test was applied to determine statistical correlation. P-value ≤ 0.05 was taken as significant.

Results

A total of 199 patients with thoracic trauma were included in this study, 154 (77.4%) were males and 45 (22.6%) were females. The mean age was 39.45 ± 16.46 years (range: 12-90 years). There were 62 (31.1%) young participants (age less than 30 years), 101 (50.8%) middle aged participants (age 30-60 years), and 36 (18.1%) elderly (age more than 60 years).

In our study there were 126 (63.3%) patients with blunt traumas and 73 (36.6%) patients with penetrating traumas. Among penetrating trauma, gunshot injuries (n=41; 56.2%) were the most common, followed by stab wounds (n=28, 38.3%) and occupational injuries (n=4, 5.4%) as shown in Figure 1.

Among blunt traumas, RTAs (n=83; 65.8%) were frequently identified, followed by fall from height (n=36, 28.5%), assault (n=4, 3.2%) and occupational injuries (n=3, 2.4%) as shown in Figure 2.

Motor bike accidents were the most frequent mode of RTA (n=38; 45.7%). There were 21 (25.3%) car-related accidents including car collision, fall from car, and hit by car, and 13 (15.6%) rikshaw related accidents. There were 11 (13.2%) bus-related accidents including bus collision, fall from bus, and hit by bus. Characteristics of all patient injuries including mode of injuries, diagnosis, and associated thoracic and non-thoracic injuries are summarized in Table 1.

Characteristics of Injuries	Frequency (%)
MODE OF INJURY	
Blunt	126 (63.3%)
Penetrating	73 (36.6%)
DIAGNOSIS AT THE TIME OF ADMISSION	
Hemothorax	73 (36.7%)
Hemopneumothorax	57 (28.8%)
Pneumothorax	40 (20.1%)
Multiple rib fracture	15 (7.5%)
Miscellaneous	14 (7.0%)
Flail chest	12 (6.0%)
Tension Pneumothorax	2 (1.0%)
ASSOCIATED THORACIC INJURIES	
Rib fracture	117 (58.8%)
Lung contusion	67 (33.7%)
Subcutaneous emphysema	32 (16.1%)
ASSOCIATED NON THORACIC INJURIES	
Bone fractures	102 (51.2%)
Abdominal injuries	32 (16.1%)
Head injuries	23 (11.6%)
Maxillofacial injuries	9 (4.5%)

TABLE 1: Characteristics of Injuries (N=199)

Among bone fractures, femur (n=39) and clavicle (n=37) were the most common fractured bones. Among 23 patients with associated head injury, 16 (8.0%) patients had traumatic brain injury (TBI) including diffuse axonal injury (DAI) and extra dural hemorrhage (EDH). Of 32 (16.1%) patients with abdominal injuries, liver (n=16, 50%) and spleen (n=12, 37.5%) were most commonly involved. The mean time interval between arrival at the ER and medical/surgical intervention was 15.9 ± 12.5 minutes (range: 5-55 minutes).

Surgical intervention included tube thoracostomy in 166 (83.4%), wound repair and debridement in 23 (11.6%), laparotomy in 21 (10.5%), craniotomy in 10 (5.0%), thoracotomy in seven (3.51%), removal of foreign body/bullet in two (1.0%) and conservative management in 24 (12.1%) patients. Out of 166 patients (83.4%) with tube thoracostomy, 119 (71.7%) patients did not require any other surgical intervention and 47 (28.3%) required more than one surgical procedure due to associated non-thoracic injuries. In 16 (69.5%) patients with wound repair, tube thoracostomy was performed. Seven patients required thoracotomy; their indications are summarized in Table 2.

S. No.	Indication	Intraoperative Findings	Mode of Trauma
1.	Left Diaphragmatic hernia with multiple rib fracture	Large defect (10x10 cm) with herniated stomach and spleen	Blunt
2.	Unstable sternum with Hemopneumothorax	Sternal body fracture with multiple rib fracture	
3.		Ruptured Left ventricle and pericardium	
4.	>1500 ml blood loss stat 250ml/hr for more 3 hours	Superior vena cava laceration	Penetrating
5.		Intercostal vessel bleeding	
6.		Intercostal vessel bleeding	
7.		Superior vena cava laceration with lung laceration	

TABLE 2: Indications of thoracotomy (n=7)

The mean blood loss was 169.4 ± 256.5 ml (range: 0-1600 ml). Transfusion of blood products was required in 129 (64.8%) patients. Average numbers of blood products transfused were 2.73 ± 1.72 .

Mechanical ventilation was needed in 57 (28.6%) patients, 44 patients had blunt trauma ($p=0.01$), 22 with hemothorax ($p<0.001$), rib fractures were seen in 40 ($p=0.03$), 20 patients with abdominal injuries, and 22 with head injury ($p<0.001$). In patients on ventilatory support, surgical interventions included tube thoracostomy in 56 ($p<0.001$), thoracotomy in five ($p=0.03$), other non-thoracic procedures in 30 ($p<0.001$). MV was also related with those having four or more rib fractures ($p=0.02$).

The mean duration of hospital stay was 4.4 ± 2.5 days (range: 1-14 days). The mortality rate in our study was 11.1% ($n=22$). Among these patients, 19 (86.4%) had associated non-thoracic injuries and only three (13.6%) with thoracic injury without any other associated non-thoracic injury ($p>0.05$). Out of three, one had flail chest with acute respiratory distress syndrome (ARDS) and two patients had massive hemothorax for which they underwent thoracotomy.

Mortality was seen in 22 (11.05%) patients. The predictors of death include higher mean blood loss (365.9 ± 460.6 ml), MV ($p=0.001$), hemothorax, and associated head injury ($p\leq 0.05$). Characteristics of the patients stratified according to their outcome are presented in Table 3.

Variables	Outcome		P value
	Discharge (n=177)	Death (n=22)	
Mode of injury			
Penetrating trauma	69 (38.9%)	4 (18.2%)	0.05
Blunt trauma	108 (61.0%)	18 (81.8%)	
Diagnosis at the time of admission			
Hemothorax	59 (33.3%)	14 (63.6%)	0.005
Hemopneumothorax	52 (29.4%)	5 (22.7%)	0.51
Pneumothorax	38 (20.9%)	2 (13.6%)	0.17
Flail chest	10 (5.9%)	2 (9.1%)	0.52
Multiple rib fracture	15 (8.5 %)	0	NA
Tension Pneumothorax	1 (0.5%)	1 (4.5%)	0.07
Associated thoracic injuries			
Rib fracture	100 (56.5%)	17 (77.3%)	0.06
Lung contusion	62 (35.0%)	5 (22.7%)	0.19
Subcutaneous emphysema	30 (16.9%)	2 (9.1%)	0.34
Associated non-thoracic injuries			
Bone fractures	85 (48.0%)	14 (63.6%)	0.07
Abdominal injuries	24 (13.5%)	6 (27.3%)	0.09
Head injury	13 (7.3%)	10 (45.5%)	<0.001
Maxillofacial injury	8 (4.5%)	1 (4.5%)	0.99
Surgical procedures			
Tube thoracostomy	145 (81.9%)	21 (95.5%)	0.10
Thoracotomy	5 (2.8%)	2 (9.09%)	0.13
Others	53 (29.9%)	10 (45.5%)	0.14
Mechanical ventilation	36 (20.3%)	21 (95.5%)	<0.001
Mean blood loss (ml)	144.9 ± 207.9	365.9 ± 460.6	<0.001

TABLE 3: Stratification of patient characteristics according to their outcome (N=199)

Discussion

Traumatic thoracic injuries are commonly blunt in nature and affect middle-aged men (30-60 years). Males are more prone as they are mobile and engaged in high-risk activities [4]. The majority of BCIs are a result of RTAs; other common causes include fall from height, assault, and occupational injuries [2-4]. Similar male predilection and involvement of young to middle-aged individuals has also been reported in older Pakistani reports [6,10,11]. In a country like Pakistan, where most of the population behind the wheel is a middle-aged male, the socioeconomic implications of RTAs and subsequent critical injuries are devastating. Blunt traumas may be fatal in case of associated lung injuries or polytrauma [4,8,9]. Among RTAs, motorbike accidents were most frequently encountered. In low-to-middle-income countries, motorbike accidents are a severe problem due to various reasons such as inexperienced bikers, disregard for safety measures, over speeding, overloading bikes with multiple passengers, and road conditions such as narrow, non-maintained, and poorly illuminated at night [12,13]. Penetrating trauma, in this study, was most commonly as a result of gunshot (56.2%) and stab (38.3%) wound. Most international data has reported stab wounds to be more common than gunshot injuries (93.7% vs. 5.94%) [3,14]. In some local studies, stab wounds have been more frequently encountered and in others gunshot wounds [10,11].

We reported hemothorax, hemopneumothorax, and pneumothorax as the common primary thoracic injuries and rib fractures and lung contusions as the common associated thoracic injuries. Hemothorax was related to more adverse outcomes such as MV ($p=0.001$) and mortality ($p=0.005$) also reported in other studies [7]. Hemothorax leads to hypovolemic shock pertaining to decrease organ perfusion and further assisted ventilation [15]. Hemothorax in blunt trauma may be due to rib fractures, laceration in lung parenchyma, and injuries to major thoracic and intercostal arteries [2], as is the case in our study.

Rib fractures were associated with MV only ($p=0.03$). With an increase in number of rib fractures the need for MV also increases. We saw rib fracture of four and more to be associated with MV ($p=0.02$). Shulzhenko et al. stated that increased number of rib fractures is associated with increase in mechanical ventilation time [16]. Although we did not find any correlation of rib fractures with mortality ($p=0.06$), Yadollahi et al. in their work reported more than thrice the risk of mortality in blunt trauma patients with rib fractures. Their mortality risk was not associated with hemothorax ($p\geq 0.05$) [4] as in our study. Literature has reported varying frequencies of hemothorax, hemopneumothorax, pneumothorax, rib fractures, and lung contusions in patients with thoracic trauma [6,11,17,18]. The frequencies may vary drastically depending on the extent of the injury.

In non-thoracic associated injuries bone fractures - especially femur and clavicle - and injury to abdominal viscera - liver and spleen - was common in our study. Bone fractures were the most frequently encountered associated injury. It did not affect the outcome and was not associated with MV ($p=0.40$) or mortality ($p=0.07$). Abdominal injuries were associated with MV ($p=0.001$); and head injuries, although, not as common, were associated with both MV and mortality ($p=0.001$). Associated extremity and visceral injuries have been commonly reported in the literature [10,17,19]. Association of non-thoracic injuries has been a predictor of mortality in these patients as explained in the literature [20,21]. There is twice the risk of mortality in patients with head injuries associated with thoracic trauma [18].

Tube thoracostomy was utilized for management of most of our patients (83.4%). In the literature, tube thoracostomy was performed in 45-89% patients [3,6,17,21]. Very few of our patients required thoracotomy (3.5%). Literature also reports the need for thoracotomy in a minority of patients (5-9%) [3,6,16,20]. Only a small proportion of cases were managed conservatively (12.1%); the proportion of patients being managed conservatively was higher in the literature (33-38%) [3]. Both tube thoracostomy and thoracotomy were related to MV ($p=0.001$ and 0.03 respectively).

We managed all of our flail chest patients ($n=12$; 6%) non-surgically as according to The Eastern Association for the Surgery of Trauma (EAST) guidelines. Surgery should only be reserved for patients who cannot be weaned off from the ventilator [22]. Nine patients with flail chest needed MV, out of which one died secondary to ARDS.

El-Menyar et al. proposed that thoracotomy increased the risk of death by 2.3 times but the study failed to provide any significant statistical correlation. The same results were seen in our study ($p=0.13$) [18]. The overall mortality rate in our study was 11.1%, similar to a previous report of 12% [6]. With improved risk stratification, mortality can further be reduced. Literature has reported a mortality rate of 1-22% for traumatic thoracic injuries [3,10,11,17,18,20,21,23].

This is the first study to present the pattern and outcome of traumatic thoracic injuries from Jinnah Postgraduate Medical Center, Karachi. This study is unique in reporting for the first time the data from the largest tertiary care institute in south Pakistan. It has helped us document the burden of thoracic trauma in a large multidisciplinary setup even in the duration in which it was studied. It identifies the outcome of thoracic injuries and factors which determine the outcome. However, this study has its limitations too. It cannot generalize the situation of the entire country as it was based on one institution only. Its mortality report is only in-hospital and it did not include post-discharge morbidity or mortality.

Conclusions

Traumatic thoracic injuries are a preventable cause of mortality in adults. In our population, RTAs, gunshots, and stabs remain a common cause. Tube thoracostomy is frequently needed. Thoracotomy and mechanical ventilation for managing chest injuries are not uncommon. Severity of injury and adverse outcome is determined by the presence of hemothorax, associated abdominal, and head injuries.

In the case of an event, the emergency department team and the team of specialized thoracic surgeons must come together to manage these critical patients with utmost care and vigilance. Favourable outcome can be attained with proper care and management.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. Institutional Review Board Committee, JPMC, Karachi issued approval No.F.2-81/2019-GENL/23145/JPMC. With reference to your letter

on the subject noted above and to say that Institutional Review Board has approved your subject proposal.

Animal subjects: All authors have confirmed that this study did not involve animal subjects or tissue.

Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Ludwig C, Koryllos A: Management of chest trauma. *J Thorac Dis.* 2017, 9:172-177. [10.21037/jtd.2017.03.52](#)
2. Edgecombe L, Sigmon DF, Galuska MA, et al.: Thoracic Trauma. StatPearls, Treasure Island; 2020.
3. Allassal M, Elrakhawy H, Saffan M, et al.: Chest trauma: a tertiary center experience. *SciFed J Surg.* 2017, 1:2.
4. Yadollahi M, Arabi AH, Mahmoudi A, Zamani M, Farahmand M: Blunt thoracic injury mortality and clinical presentation. *Trauma Monthly.* 2018, 23:13079.
5. Jain A, Burns B: Penetrating Chest Trauma. StatPearls, Treasure Island; 2020.
6. Saaqi M, Shah SA: Thoracic trauma: presentation and management outcome. *J Coll Physicians Surg Pak.* 2008, 18:230-233.
7. Tsai YM, Lin KH, Huang TW, Chen CY, Hong ZJ, Hsu SD: Outcomes of patients with blunt chest trauma encountered at emergency department and possible risk factors affecting mortality. *J Med Sci.* 2017, 37:97. [10.4103/jmedsci.jmedsci_123_16](#)
8. Sharma P, Jakhmola CK: Presentation and outcome of thoracic injuries at a tertiary care service hospital. *Int J Sci Stud.* 2018, 5:126-130.
9. Ekpe EE, Eyo C: Determinants of mortality in chest trauma patients. *Niger J Surg.* 2014, 20:30-4.
10. Tariq UM, Faruque A, Ansari H, et al.: Changes in the patterns, presentation and management of penetrating chest trauma patients at a level II trauma centre in southern Pakistan over the last two decades. *Interact Cardiovasc Thorac Surg.* 2011, 12:24-7. [10.1510/icvts.2010.242750](#)
11. Dongel I, Coskun A, Ozbay S, Bayram M, Atli B: Management of thoracic trauma in emergency service: analysis of 1139 cases. *Pak J Med Sci.* 2013, 29:58. [10.12669/pjms.291.2704](#)
12. Manna N, Mallik S, Mandal PK, et al.: Epidemiological factors of road traffic accidents: a study in a tertiary care setting in India. *J Pak Med Stud.* 2013, 3:48-53.
13. Khan K, Jamil M, Memon IA, Idrees Z: Pattern of injuries in motorbike accidents. *J Pak Orthop Assoc.* 2018, 30:123-127.
14. Aghaei Afshar M, Mangeli F, Nakhaei A: Evaluation of injuries caused by penetrating chest traumas in patients referred to the emergency room. *Indian J Surg.* 2015, 77:191-4. [10.1007/s12262-012-0757-4](#)
15. Taghavi S, Askari R: Hypovolemic Shock. StatPearls, Treasure Island; 2020.
16. Shulzhenko NO, Zens TJ, Beems MV, et al.: Number of rib fractures thresholds independently predict worse outcomes in older patients with blunt trauma. *Surgery.* 2017, 161:1083-9. [10.1016/j.surg.2016.10.018](#)
17. Al-Koudmani I, Darwish B, Al-Kateb K, Taifour Y: Chest trauma experience over eleven-year period at al-mouassat university teaching hospital-Damascus: a retrospective review of 888 cases. *J Cardiothorac Surg.* 2012, 7:35. [10.1186/1749-8090-7-35](#)
18. El-Menyar A, Abdelrahman H, Al-Hassani A, et al.: Clinical presentation and time-based mortality in patients with chest injuries associated with road traffic accidents. *Arch Trauma Res.* 2016, 5:31888. [10.5812/atr.31888](#)
19. Kumar BA, Chakravarthy GR, Bharath A: Blunt trauma chest: a study on clinical pattern. *J Dent Med Sci.* 2017, 16:1-7. [10.9790/0853-1603070107](#)
20. Lema MK, Chalya PL, Mabula JB, Mahalu W: Pattern and outcome of chest injuries at Bugando Medical Centre in Northwestern Tanzania. *J Cardiothorac Surg.* 2011, 6:7. [10.1186/1749-8090-6-7](#)
21. Çakmak M, Kandemir MN: Study of 433 operated cases of thoracic trauma. *Indian J Surg.* 2016, 78:477-81. [10.1007/s12262-015-1414-5](#)
22. Simon B, Ebert J, Bokhari F, et al.: Management of pulmonary contusion and flail chest: an Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg.* 2012, 73:351-61. [10.1097/TA.0b013e31827019fd](#)
23. Huber S, Biberthaler P, Delhey P, et al.: Predictors of poor outcomes after significant chest trauma in multiply injured patients: a retrospective analysis from the German Trauma Registry (Trauma Register DGU®). *Scand J Trauma Resus Emerg Med.* 2014, 22:52. [10.1186/s13049-014-0052-4](#)