Bilateral Renal Lymphangiectasia Associated With Polycythemia: A Rare Case Report

Tahereh Orouji Jokar¹, Ikechukwu Chidobem², Nazia Khan²

1. Hematology and Oncology, Prime Healthcare, Saint Clare's Hospital, Denville, USA 2. Internal Medicine, Prime Healthcare, Saint Clare's Hospital, Denville, USA

Corresponding author: Tahereh Orouji Jokar, oroujit@gmail.com

Abstract

Renal lymphangiectasia is a rare benign mesenchymal tumor of unclear etiology resulting from dilatation of perinephric lymphatic channels and formation of cystic masses. Polycythemia is a rarely associated finding with only five cases reported in the literature. We report a case of bilateral renal lymphangiectasia associated with polycythemia in a 38-year-old man who was managed conservatively with pain control. There are no clear guidelines for the management of renal lymphangiectasia; although most patients can be treated conservatively, some cases, whose diagnosis is unclear or develop complications, require definitive surgical excision.

Categories: Oncology, Hematology

Keywords: lymphangiectasia, renal lymphangiectasia, cystic retroperitoneal mass, perinephric lymphangiectasia, bilateral lymphangiectasia, polycythemia

Introduction

Renal Lymphangiectasia, is a rare disorder of unclear etiology, with arguments in the literature both for and against an acquired versus congenital pathogenesis [1]. It is a bilateral cystic infiltration of the perirenal and parapelvic space, secondary to a miscommunication between the renal and retroperitoneal lymphatics, which results in dilatation of perinephric lymphatic channels and can lead to the formation of a unilocular or multilocular cystic mass [2]. Here, we present a case of bilateral renal lymphangiectasia associated with polycythemia.

Case Presentation

A 38-year-old male presented with right flank pain for two days. He had a past medical history significant for poorly controlled type 2 diabetes mellitus. The patient was a never-smoker and did not drink alcohol. His physical examination was positive for tenderness in the right lower abdominal quadrant and right flank, but was otherwise normal. He was not obese. Laboratory investigations revealed polycythemia (hemoglobin, hematocrit, and red blood cell count levels at 18.9 g/dl, 57.2% and 6.57 million/uL, respectively). Leukocyte count and platelet count were normal. Kidney function test was normal and urinalysis was positive for glucosuria, but was otherwise unremarkable. His polycythemia prompted molecular analysis of the patient's blood, which was negative for Janus kinase 2 (JAK2), V617F, Exon 12, calreticulin gene (CALR), and MPL mutations, thereby making polycythemia vera an unlikely diagnosis. Erythropoietin and lactate dehydrogenase levels were also within the normal range.

A computed tomography (CT) scan of the abdomen/pelvis showed bilaterally enlarged kidneys with multiple bilateral parapelvic cysts, as well as infiltrating soft tissues bilaterally (left greater than the right), infiltrating nodal tissues along the roof of the mesentery extending into the right lower quadrant, and infiltrating nodal tissues within the retro-peritoneum surrounding the aorta and inferior vena cava, as well as an enhancing splenic nodule (Figures 1-2). A follow up full body fluorodeoxyglucose (FDG)-positron emission tomography (PET) scan showed a soft tissue nodule within the spleen, bilaterally enlarged kidneys with multiple bilateral parapelvic cysts, as well as bilateral pararenal hypodense soft tissues, and infiltration of soft tissues within the retroperitoneum as well as in the periaortic region (Figure 3). The patient was managed conservatively with pain control. A CT-guided aspiration and core biopsy of the left parapelvic cystic mass and left perinephric soft tissue mass was done, and the patient was discharged to outpatient follow up. The pathology report later showed no evidence of malignancy. The cystic fluid cell count and cultures were also unremarkable.

Review began 06/07/2021

Review ended 06/23/2021

Published 07/03/2021

© Copyright 2021

Orouji Jokar et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Orouji Jokar T, Chidobem I, Khan N (July 03, 2021) Bilateral Renal Lymphangiectasia Associated With Polycythemia: A Rare Case Report. Cureus 13(7): e16130. DOI 10.7759/cureus.16130

FIGURE 1: CT findings of lymphangiectasia: arrows pointing to perinephric lymphatic infiltration

FIGURE 2: CT findings of perirenal lymphangiectasia: arrows pointing to bilateral perinephric lymphangiectasia with more pronounced changes on the left

FIGURE 3: Positron emission tomography scan: arrow highlighting left perinephric infiltration with lymphatic tissue

Discussion

Renal lymphangiectasia is a rare disorder, which is usually bilateral, and unrelated to age or gender [3]. Patients with renal lymphangiectasia are usually asymptomatic but may present with abdominal distension, flank pain, abdominal pain, hypertension, hematuria, or renal vein thrombosis [1,3]. Complications described in previous reports include hemorrhage, infection, and deterioration in renal function [2,4]. Polycythemia is a very rare association of the condition, with only five cases reported in the literature [3].

The differential diagnosis of this condition includes autosomal dominant polycystic kidney disease, although in this condition the renal parenchyma is diffusely abnormal; tumors such as liposarcoma, leiomyosarcoma, fibrosarcoma, malignant teratomas, and multilocular cystic nephroma, which are predominantly cystic or

necrotic, but these malignancies usually contain substantial solid components; abscess and urinoma may also produce a similar appearance, but almost all these conditions can be differentiated with the help of clinical history, normal biochemical parameters and typical imaging findings of perirenal and parapelvic involvement sparing the renal parenchyma [2,5].

The diagnosis of renal lymphangiectasia is usually made confidently on imaging (ultrasonography, CT, or MRI) due to its characteristic cystic patterns, however, as in our case, aspiration and cytological examination of the fluid may be performed if there is suspicion for malignancy [1-4,6].

There are no clear guidelines as to when conservative management, percutaneous drainage, or surgery should be used; however, treatment is generally conservative for asymptomatic patients. Complicated cases are treated with nephrectomy, percutaneous drainage, and marsupialization, however, there is a risk of an increase in the size of cysts in the contralateral kidney when a nephrectomy is performed in patients with bilateral involvement [1,3,4,5].

Conclusions

Renal lymphangiectasia is a rare disorder, usually asymptomatic. Diagnosis is primarily radiologic, however, other modalities may be necessary to rule out confounding etiologies. Treatment varies based on symptomatic presentation and can range from observation to radical surgical resection.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Bansal K, Sureka B, Pargewar S, Arora A: Renal lymphangiectasia: one disease, many names! . Indian J Nephrol. 2016, 26:57-8. [10.4103/0971-4065.155730](https://doi.org/10.4103/0971-4065.155730)
2. Parazols F, Fournier L, Bouvier N, Pelage JP, Lobbedez T, Hurault de Ligny B: Renal lymphangiectasia, a rare case of perirenal infiltration [Article in French]. Nephrol Ther. 2015, 11:118-20. [10.1016/j.nephro.2014.11.005](https://doi.org/10.1016/j.nephro.2014.11.005)
3. Chaabouni A, Rebai N, Fourati M, et al.: Cystic lymphangioma of the kidney: diagnosis and management . Int J Surg Case Rep. 2012, 3:587-9. [10.1016/j.ijscr.2012.01.013](https://doi.org/10.1016/j.ijscr.2012.01.013)
4. Blanc M, Schmutz G, Belzile F, Sabbagh R: Renal lymphangiectasia presenting with hypertension and polycythemia. Can Urol Assoc J. 2014, 8:E163-6. [10.5489/cuaj.1596](https://doi.org/10.5489/cuaj.1596)
5. Umapathy S, Alavandar E, Renganathan R, Thambidurai S, Arunachalam VK: Renal lymphangiectasia: an unusual mimicker of cystic renal disease - a case series and literature review. Cureus. 2020, 12:e10849. [10.7759/cureus.10849](https://doi.org/10.7759/cureus.10849)
6. Jeon TG, Kong DH, Park HJ, Kim S, Park WY, Kim SD, Ha HK: Perirenal lymphangiomatosis. World J Mens Health. 2014, 32:116-9. [10.5534/wjmh.2014.32.2.116](https://doi.org/10.5534/wjmh.2014.32.2.116)