

Causes of the Failure of Biological Therapy at a Tertiary Center: A Cross-Sectional Retrospective Study

Review began 08/05/2021

Review ended 09/19/2021

Published 09/24/2021

© Copyright 2021

Fatani et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Arwa Z. Fatani¹, Nada A. Bugshan¹, Hanan M. AlSayyad¹, Mayar A. Shafei¹, Nada M. Hariri¹, Laila T. Alrashid¹, Ahlam Y. Lasker¹, Badreyah A. Aldauig¹, Suzan M. Attar²

1. Faculty of Medicine, King Abdulaziz University, Jeddah, SAU 2. Department of Internal Medicine, King Abdulaziz University, Jeddah, SAU

Corresponding author: Arwa Z. Fatani, arwazuhaif@gmail.com

Abstract

Introduction

Rheumatoid arthritis (RA) is one of the most commonly encountered autoimmune diseases. Treatment generally includes disease-modifying anti-rheumatic drugs (DMARDs) and/or biological therapy. However, a significant proportion of the patients do not respond to treatment either as a (primary failure) or lose efficacy over time (secondary failure). Several factors are assumed to influence these conditions.

Objectives

To estimate the prevalence of failure of biological therapy in patients with RA and its causes.

Methods

A total of 335 RA patients who were diagnosed at a tertiary center in Jeddah, Saudi Arabia, and had a failure after receiving biological therapy were included in this study. Several variables were considered; patient's socio-demographic data, comorbid conditions, types of biological therapy, the duration of using biological therapy in months, number of biological therapies, allergic reactions, disease activity, and treatment duration.

Results

Overall the prevalence of failure to biological therapy was 58%; 77% primary failure and 23% secondary failure. Patients with negative rheumatoid factor (RF) ($p=0.006$), using low-dose steroids, and with a longer disease duration had a significant failure of biological therapy ($p=0.023$).

Conclusion

A high percentage of RA patients had a failure of biological therapy. A multicentric trial is recommended to look for additional factors.

Categories: Internal Medicine, Rheumatology

Keywords: rheumatoid arthritis, biological therapy, refractory disease, failures, tnf inhibitors, rituximab

Introduction

The principal rheumatoid arthritis (RA) treatments are disease-modifying anti-rheumatic drugs (DMARDs) and/or biological DMARDs (bDMARDs) [1]. The introduction of bDMARDs designed to inhibit particular cellular or molecular targets specifically involved in the pathogenesis of the disease has significantly enhanced the outcome of RA therapy, resulting in clinical recovery or decreased activity of the disease. However, patients may also discontinue or delay care due to unfavored side effects. A randomized controlled trial in Greece showed that a considerable number of patients discontinued anti-tumor necrosis factor (TNF) therapy in 2019 either due to primary failure, secondary lack of response, or intolerance. Primary failure was observed in a considerable percentage of RA patients who don't respond to bDMARDs. Treatments that are possibly a result of non-TNF inflammatory pathways may be dominant in individual patients. In a previous study, it was found that starting DMARDs at a younger age, a high baseline disease activity score (DAS)-28 score, poorer early response within the first six months of treatment with bDMARDs (estimated by delta-DAS-28), and the presence of erosions were correlated with multi-refractoriness. However, anti-TNF agents are not effective in all patients. About 30% of patients treated with a TNF inhibitor failed to achieve an improvement of 20% in American College of Rheumatology criteria (ACR20; primary failure or inefficacy), and more patients lose efficacy during therapy (secondary failure or acquired therapeutic resistance) or experience adverse events following treatment with a TNF inhibitor. In addition, others may demonstrate primary response initially and then develop over time (secondary failure) as a decline of efficacy. There is a

How to cite this article

Fatani A Z, Bugshan N A, AlSayyad H M, et al. (September 24, 2021) Causes of the Failure of Biological Therapy at a Tertiary Center: A Cross-Sectional Retrospective Study. Cureus 13(9): e18253. DOI 10.7759/cureus.18253

previous study that documented the frequency of patients who do not achieve even the weakest response to the standard dosage of anti-TNF agents, which ranges between 28% and 58%, namely, ACR20 (20% improvement) response. Efficacy and adverse effects were the major factors for the discontinuation. The other factors are genetic mutations such as FAS-L and Caspase-9 in the apoptosis-related genes. There are other contributing factors to secondary failure of bDMARDs, such as a longer period of illness lasting more than two years, smoking, and small bowel involvement [2]. An additional cause that influences TNF- α inhibitors' secondary failure is anti-drug antibodies development [3]. Methotrexate (MTX) discontinuation after the initiation of bDMARDs, adherence to treatment, or variations in the pharmacokinetics of TNF- α inhibitors. Still, there aren't enough studies in the literature. Therefore, we aimed to determine retrospectively the causes of bDMARDs failure in RA patients in a tertiary center.

Materials And Methods

A retrospective study was conducted from April 2015 to January 2019 at a tertiary center. The ethical committee approved this study. Our sample size comprised 335 patients (51 (11.9%) of male patients) and (284 (66.2%) of female) patients who had either primary or secondary failure to RA treatment. Primary failure is generally defined as no clinical response within the initial treatment while secondary failure is defined as loss of effectiveness of the drug after initial remission. The data included admitted patients and those in the database. Those who failed to respond to bDMARDs were not able to progress to a better prognosis and the desired treatment plan. While patients experiencing infection, pregnancy, or had recent cancer <5 years were excluded. The medical records were obtained, collected, and evaluated using a dedicated data extraction sheet along with examining the patient in the clinic. These records consisted of the patient's sociodemographic data based on their age, gender, nationality, body mass index (BMI), weight, height, smoking, disease duration, family history of RA, and a history of abortions and pregnancies. The comorbid conditions were diabetes mellitus, tuberculosis (TB), hypertension, liver cirrhosis, hyperlipidemia, cardiovascular diseases, cancer, lymphoma, chronic obstructive pulmonary disease (COPD), asthma. The medications were methotrexate, steroids, aspirin, non-steroidal anti-inflammatory drugs (NSAIDs), and traditional DMARDs. The use of bDMARDs included adalimumab, etanercept, infliximab, rituximab, and tocilizumab. The records also included the duration of using bDMARDs in months, the name of the first bDMARDs, the number of current bDMARDs, and the name of the second bDMARDs. Also, the study measured local and general allergic reactions and the lab results were rheumatoid factor (RF) positive or negative, RF level, anti-cyclic citrullinated peptide (CCP) level, erythrocyte sedimentation rate (ESR) level, c-reactive protein (CRP) level, and antinuclear antibodies (ANA) level. Disease activity was measured by the DAS-28 score and the result was measured as well. Data entry was done using Microsoft Excel 2016 (Microsoft Corporation, Redmond, WA) and statistical analysis was performed by using the Statistical Package for the Social Sciences (SPSS) software, version 21 (IBM Corp., Armonk, NY). Categorical variables, including primary variables, were described using frequencies. Continuous variables for normally distributed were described using means and standards. A univariate analysis was conducted for categorical variables using the chi-square test to check for all the possible causes. A test with a p-value of <0.05 was considered significant. All information in this study was confidential, with no access to data other than to those authorized.

Results

Twenty-six percent (26%) of studied patients had an age ranging from (40 to <50 years), 84.8% were females with 2.5% who had a previous pregnancy and 7.4% who had a previous abortion. Of these patients, 65.1% were of Saudi nationality, 15.2% were current smokers, and 32.2% were obese. Most of the patients (89.6%) were on monotherapy biological treatment, 19.1% had a family history of RA, and 43% had RF (Table 1).

Variable		No. (%)
Age	<40	98 (9.3)
	40 - <50	87 (26)
	50 - <60	762 (22.7)
	≥60	74 (22.1)
Gender	Male	51 (15.2)
	Female	284 (84.8)
For females	Pregnancy	
	Yes	7 (2.5)
	No	277 (97.5)
	Previous abortion	
	Yes	21 (7.4)
	No	263 (92.6)
Nationality	Saudi	218 (65.1)
	Non- Saudi	117 (34.9)
Smoking	Yes	51 (15.2)
	No	274 (81.8)
	Ex-smoker	10 (3)
BMI categories	Underweight	6 (1.8)
	Normal	56 (16.7)
	Over	74 (22.1)
	Obesity	108 (32.2)
	Severe obesity	91 (27.2)
Current biological therapy	Monotherapy	300 (89.6)
	Two types	21 (6.3)
	Three types	14 (4.2)
Family history of RA	Yes	64 (19.1)
	No	271 (80.9)
RF	Yes	144 (43)
	No	182 (54.3)
	Not applicable	9 (2.7)

TABLE 1: Distribution of studied patients according to their characters, smoking, BMI categories, currently used biological therapy, and a family history of RF

Ex-smoker; no longer smoking, RF; rheumatoid factor, RA; rheumatoid arthritis

Table 2 shows that 4.8% of patients had infertility, 12.2% had TB and most of it was of the pulmonary type (67.4%). More than half of the patients (59.4%) had comorbidities, where the most common was vasculitis (39.4%), hypertension (HTN; 29%), diabetes mellitus (DM; 28.1%), and hyperlipidemia (23.9%).

Variable	No. (%)	
Infertility	Yes	16 (4.8)
	No	319 (95.2)
TB	Yes	41 (12.2)
	No	294 (87.8)
Old TB	Yes	30 (9)
	No	305 (91)
Latent TB	Yes	20 (6)
	No	315 (94)
TB type	Bone	5 (11.6)
	Abdomen	5 (11.6)
	Potts	4 (9.3)
	Pulmonary	29 (67.4)
Co-morbidity	Yes	199 (59.4)
	No	136 (40.6)
Co-morbid disease	DM	94 (28.1)
	HTN	97 (29)
	CVD	31 (9.3)
	Hyperlipidemia	80 (23.9)
	Hypoalbuminemia	47 (14)
	Renal impairment	46 (13.7)
	Previous lung diseases	64 (19.1)
	Asthma	29 (8.7)
	COPD	18 (5.4)
	Lymphoma	19 (5.7)
	Vasculitis	132 (39.4)
	Osteoporosis	40 (11.9)
	SLE	41 (12.2)
	Psoriasis	30 (9)
	Cirrhosis	1 (0.3)
	Fatty liver	58 (17.3)
	Cancer:	34 (10.1)
	Cancer thyroid	1 (2.9)
	Cancer lung	13 (38.2)
	Multiplemyeloma	7 (20.6)
Cancer breast	4 (11.8)	
Cancer bladder	4 (11.8)	
Cancer colon	5 (14.7)	

TABLE 2: Distribution of studied patients according to the presence of infertility, TB, and

comorbidities

TB; tuberculosis, DM; diabetes mellitus, HTN; hypertension, CVD; cardiovascular diseases, COPD; chronic obstructive lung disease, SLE; systemic lupus erythematosus

Table 3 demonstrated that 9.3% of patients had an allergic reaction to taken biological therapy and the most common allergic drug was rituximab (41.9%). Of the patients, 49.6% had anti-CCP, 57.6% had anti-mutated citrullinated vimentin (MCV), 51.3% had high ESR, 70.4% had a CRP level >3, and 31.4% had a double-stranded DNA (dsDNA) level of 0-200. Most of the patients (60%) had a positive ANA, 83.6% had disease activity, and 48.7% had a DAS-28 of 3.2-5.1, 46.6% had an aspartate aminotransferase (AST) of 10-40, 49% had an alanine transaminase (ALT) of 7-56, and 19.4% had an acetylsalicylic acid (ASA) of 65. Most of the patients were using NSAIDs (80.6%), 56.7% were using a steroid dose of <5 mg, 80% were using Methtrix, and 78.5% were using Methotrexate (MTX). The mean disease duration (months), HB, platelet, and creatine levels, and MTX dose were 38.91 ± 70.16 months, 11.78 ± 1.73, 324.24 ± 100.71, 88.65 ± 389.67, and 8.68 ± 4.41, respectively.

Variable	No. (%)	
Allergic reaction to taken biological therapy	Yes	31 (9.3)
	No	304 (90.7)
If yes, what is the allergic drug?	Rituximab	13 (41.9)
	Infliximab	6 (19.4)
	Adalimumab (Humira)	4 (12.9)
	Etanercept (Enbrel)	4 (12.9)
	Tocilizumab	4 (12.9)
Anti-CCP	Yes	166 (49.6)
	No	169 (50.4)
Anti-MCV	Yes	193 (57.6)
	No	141 (42.1)
	Not applicable	1 (0.3)
ESR	High	172 (51.3)
	Normal	163 (48.7)
CRP	>3	236 (70.4)
	≤ 3	98 (29.3)
	Not applicable	1 (0.3)
dsDNA	0-200	41.5 (31.4)
	201-300	30.1 (30.1)
	301-800	26.9 (26.9)
	>800	1.5 (1.5)
ANA	Positive	201 (60)
	Negative	134 (40)
Activity	Yes	280 (83.6)
	No	55 (16.4)
DAS-28	≤ 2.6	2 (0.6)
	2.6-3.2	114 (34)
	3.2-5.1	163 (48.7)

	≥ 5.1	56 (16.7)
	≤ 2.6	2 (0.6)
AST	<10	93 (27.8)
	10-40	156 (46.6)
	>40	86 (25.7)
ALT	<7	93(27.8)
	7-56	164 (49)
	>56	78 (23.3)
ASA	Yes	65 (19.4)
	No	270 (80.6)
NSAIDs	Yes	270 (80.6)
	No	65 (19.4)
Steroid dose	<5mg	190 (56.7)
	5-10mg	133 (39.7)
	>10mg	12 (3.6)
Used drugs	Methotrexate	263 (78.5)
	Antimalarial	107 (31.9)
	Methtrx	268 (80)
	Sulfa	60 (17.9)
	Avara	49 (14.6)
	Immurane	47 (14)
Mean and SD of study variables	Disease duration (months)	38.91 ± 70.16
	HB level	11.78 ± 1.73
	Platelet level	324.24 ± 100.71
	Creatine level	88.65 ± 389.67
	MTX dose	8.68± 4.41

TABLE 3: Distribution of studied patients according to the allergic reaction to taken biological therapy, clinical and laboratory data, and drugs used

Anti-CCP; anti-cyclic citrullinated peptide, Anti-MCV; anti-mutated citrullinated vimentin, ESR; erythrocyte sedimentation rate, CRP; C-reactive protein, dsDNA; double-stranded DNA, ANA; antinuclear antibodies, DAS-28; disease activity score, AST; aspartate aminotransferase, ALT; alanine transaminase, NSAIDs; nonsteroidal anti-inflammatory drugs, HB level; hemoglobin level, MTX dose; methotrexate dose, ASA; acetylsalicylic acid

Of the studied patients, 58.5% failed to respond to biological therapy. In contrast, only 41.5% responded to biological therapy.

Figure 1 shows that most of the patients were using Humira (34%) and rituximab (49.3%) as the first biological therapy and Humira (22.7%) and rituximab (34%) as the second biological therapy.




FIGURE 1: Percentage distribution of patients according to used first and second biological therapy

Table 4 shows that patients of Saudi nationality and those who did not have RF had a significantly higher percentage of those who had a failure of biological therapy ($p < 0.05$). On the other hand, a non-significant difference was found between failure of biological therapy and other patient characteristics such as smoking, BMI categories, and currently used biological therapy.

Variable	Biological failure		χ^2	P-value	
	Present No. (%)	Absent No. (%)			
Age	<40	37 (37.8)	61 (62.2)	1.75	0.627
	40 - <50	41 (47.1)	46 (52.9)		
	50 - <60	31 (40.8)	45 (59.2)		
	≥ 60	30 (40.5)	44 (59.5)		
Gender	Male	22 (43.1)	29 (56.9)	0.06	10.796
	Female	117 (41.2)	167 (58.8)		
Nationality	Saudi	80 (36.7)	38 (63.3)	5.91	0.015
	Non- Saudi	59 (50.4)	58 (49.6)		
Smoking	Yes	25 (49)	26 (51)	3.08	0.214
	No	108 (39.4)	166 (60.6)		
	Ex-smoker	6 (60)	4 (40)		
BMI categories	Underweight	0 (0.0)	6 (100)	5.14	0.273
	Normal	22 (39.3)	34 (60.7)		
	Over	29 (3.2)	45 (60.8)		
	Obesity	48 (44.4)	60 (55.6)		
	Severe obesity	40 (44)	51 (956)		
Family history of RA	Yes	32 (50)	32 (50)	2.35	0.125
	No	107 (39.5)	164 (60.5)		
RF	Yes	74 (51.4)	70 (48.6)	10.18	0.001
	No	65 (34)	126 (66)		

TABLE 4: Relation to failure of biological therapy and patients' characteristics, smoking, BMI categories, currently used biological therapy, and family history of RF

RA; rheumatoid arthritis, RF; rheumatoid factor, BMI; body mass index

Table 5 shows that patients with vasculitis had a significantly higher percent of those who had a failure of biological therapy ($p < 0.05$). On the other hand, a non-significant difference was found between the failure of biological therapy and infertility, TB, and other comorbidities (cancer ($\chi^2=1.3$, p -value=0.254), COPD ($\chi^2=1.55$, p -value=0.213), asthma ($\chi^2=3.83$, p -value=0.05), previous lung diseases ($\chi^2=0.47$, p -value=0.47), renal impairment ($\chi^2=0.38$, p -value=0.538), hypoalbuminemia ($\chi^2=0.63$, p -value=0.424), hyperlipidemia ($\chi^2=0.22$, p -value=0.639), CVD ($\chi^2=2.5$, p -value=0.113), DM ($\chi^2=0.54$, p -value=0.459), osteoporosis ($\chi^2=0.67$, p -value=0.411), SLE ($\chi^2=0.46$, p -value=0.496), and psoriasis ($\chi^2=0.04$, p -value=0.63) ($p > 0.05$).

Variable		Biological failure		χ^2	P-Value
		Absent No. (%)	Present No. (%)		
TB	Yes	23 (56.1)	18 (43.9)	0.11	0.73
	No	173 (58.8)	121 (41.2)		
Comorbidity	Yes	116 (58.3)	83 (41.7)	0.009	0.92
	No	80 (58.8)	56 (41.2)		

TABLE 5: Relation to failure of biological therapy and presence of infertility, TB, and comorbidities

TB; tuberculosis

Table 6 shows that patients who had an allergic reaction to taken biological therapy, those who used NSAIDs and who had a steroid dose <5 mg, who were not using antimalarial drugs, sulfa, or Immurane, and those with a longer disease duration had a significantly higher percentage of those who had a failure of biological therapy ($p < 0.05$). On the other hand, a non-significant difference was found between the failure of biological therapy and other clinical and laboratory data and other drugs used (Methotrexate (DMARD) ($\chi^2 = 0.05$, p -value = 0.813), Methtrix (MTX) ($\chi^2 = 0.37$, p -value = 0.542), and Avara ($\chi^2 = 2.14$, p -value = 0.143)) ($p > 0.05$).

Variable		P-value		χ^2	P-value
		Absent No. (%)	Present No. (%)		
Allergic reaction to taken biological therapy	Yes	31 (100)	0 (0.0)	24.22	< 0.001
	No	165 (54.3)	139 (45.7)		
Anti-CCP	Yes	96 (57.8)	70 (42.2)	0.06	0.803
	No	100 (59.2)	69 (40.8)		
Anti-MCV	Yes	11 (57.5)	82 (42.5)	1.67	0.433
	No	85 (60.3)	56 (39.7)		
	Not applicable	0 (0.0)	1 (100)		
ESR	High	97 (56.4)	75 (43.6)	0.65	0.42
	Normal	99 (60.7)	64 (39.3)		
CRP	> 3	135 (57.2)	101 (42.8)	2.13	0.343
	≤ 3	61 (82.2)	37 (37.8)		
	Not applicable	0 (0.0)	1 (100)		
dsDNA	0-200	75 (45)	64 (46)	2.1	0.53
	201-300	61 (60.4)	40 (39.6)		
	301-800	57 (63.3)	33 (36.7)		
	> 800	3 (60)	2 (40)		
ANA	Positive	114 (56.7)	87 (43.3)	0.66	0.415
	Negative	82 (61.2)	52 (88.8)		
Activity	Yes	164 (58.6)	116 (41.4)	0.003	0.957
	No	32 (58.2)	23 (41.8)		
	≤ 2.6	2 (100)	0 (0.0)		

DAS-28	2.6-3.2	70 (61.4)	441 (38.6)	5.35	0.148
	3.2-5.1	98 (60.1)	65 (39.9)		
	≥ 5.1	26 (46.4)	30 (53.6)		
NSAIDs	Yes	169 (62.6)	101 (37.4)	9.56	0.002
	No	27 (41.5)	38 (58.5)		
	< 5mg	129 (67.9)	61 (32.1)		
Steroid dose	5-10mg	64 (48.1)	69 (51.9)	20.88	< 0.001
	> 10mg	2 (18.2)	9 (81.8)		
	Not applicable	1 (100)	0 (0.0)		
Quantitative variables	Duration	47.55± 85.27	26.55± 36.48	3.74*	< 0.001
	HB level	11.73 ± 1.72	11.85± 1.75	0.85*	0.393
	Platelet level	330.59±114.47	315.29 ±76.7	0.87*	0.38
	Creatine level	96.93± 50.8	76.98± 75.93	1.04*	0.298
	MTX Dose	8.58 ± 4.58	8.82 ± 4.18	0.81*	0.215

TABLE 6: Relation of failure of biological therapy and allergic reaction to taken biological therapy, clinical and laboratory data, and drugs used

* Mann-Whitney test

Anti-CCP; Anti-cyclic citrullinated peptide, Anti-MCV; Anti-mutated citrullinated vimentin, ESR; erythrocyte sedimentation rate, CRP; C-reactive protein, dsDNA; double-stranded DNA, ANA; antinuclear antibodies, DAS-28; disease activity score, NSAIDs; nonsteroidal anti-inflammatory drugs, HB level; hemoglobin level, MTX; methotrexate

By binary logistic regression analysis, using a steroid dose <5 mg was an independent predictor (risk factor) for the failure of biological therapy among studied patients ($p < 0.05$) as shown in Table 7.

Variable	P-value	Odds ratio
Nationality	0.143	1.48
RF	0.64	0.65
Vasculitis	4.66	1.23
Allergic reaction to taken biological therapy	0.998	5.04
NSAID	0.165	1.58
Steroid dose	0.023	1.79
Duration	0.177	0.99

TABLE 7: Binary logistic regression analysis of risk factors of failure to biological therapy

NSAIDs; nonsteroidal anti-inflammatory drugs, RF; rheumatoid factor

Discussion

This study estimated that more than half of RA patients had primary failure compared to secondary failure of biological therapy (bDMARD). More than half of the patients had comorbid conditions similar to previously conducted studies [4-6]. Patients who were RF negative had a significantly higher percentage of failure of biological therapy, contrary to a study that included 400 patients receiving bDMARD in the city of Bogotá, Colombia, which demonstrated that RF-negative patients have frequent remission and lower levels

of disability compared to RF-positive patients treated with anti-TNF alpha agents [7]. Furthermore, an observational study was done between January 2000 and August 2019, which reported that the later that bDMARD was initiated and longer disease duration are prone to have multi-refractory diseases, as they present with advanced disease courses. Thus, early intervention with biological therapy is recommended in order to establish beneficial treatment outcomes [5]. Our data demonstrated that the higher the dose of steroids, the better the outcome, as most of our patients who received >10 mg/d had less failure of biological therapy (18.2%). Marije F. Bakker et al. showed that the inclusion of prednisone 10 mg/d from the start of an MTX-based, tight-control strategy slows erosive joint damage and further enhances clinical efficacy [8]. A non-significant relation was found between the failure of biological therapy and smoking compared to multiple studies that proved that smoking reduced the effect of both non-biological and biological DMARDs in RA treatment [7-9]. Most of our sample were females, as they are more likely to attain autoimmune disorders due to the hormonal changes women experience [10].

To date, no research has studied the prevalence of failure of biological therapy in the Middle East. Nevertheless, further work is valuable to identify the genetic and other constitutional factors, which may be linked to the disease activity that may determine the response of the failure to biological treatment, which is improbable to be a random effect. This study has several limitations, including the fact that it has a single-center, hospital-based, cross-sectional design. However, this is one of few studies that assessed the disease activity by accurate clinical measures using DAS-28. Larger, prospective, multi-centric cohort studies are needed to highlight the advocacy of applying DAS-28 by physicians, consider failure as a complication in all RA patients, and be updated on the next step of management if so.

Conclusions

A high percentage of RA patients had a failure of bDMARDs. A multicentric trial is recommended to look for additional patient-related, drug, genetic, and environmental factors.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Strangfeld A, Hiese F, Rau R, et al.: Risk of incident or recurrent malignancies among patients with rheumatoid arthritis exposed to biologic therapy in the German biologics register RABBIT. *Arthritis Res Ther.* 2010, 12:R5. [10.1186/ar2904](https://doi.org/10.1186/ar2904)
2. Novella-Navarro M, Plasencia C, Tornero C, et al.: Clinical predictors of multiple failure to biological therapy in patients with rheumatoid arthritis. *Arthritis Res Ther.* 2020, 22:284. [10.1186/s13075-020-02354-1](https://doi.org/10.1186/s13075-020-02354-1)
3. Saevarsdottir S, Wallin H, Seddighzadeh M, et al.: Predictors of response to methotrexate in early DMARD naive rheumatoid arthritis: results from the initial open-label phase of the SWEFOT trial. *Ann Rheum Dis.* 2011, 70:469-75. [10.1136/ard.2010.139212](https://doi.org/10.1136/ard.2010.139212)
4. Marchesoni A, Zaccara E, Gorla R, et al.: TNF- α antagonist survival rate in a cohort of rheumatoid arthritis patients observed under conditions of standard clinical practice. *Ann N Y Acad Sci.* 2009, 1173:837-46. [10.1111/j.1749-6632.2009.04621.x](https://doi.org/10.1111/j.1749-6632.2009.04621.x)
5. Soliman MM, Hyrich KL, Lunt M, Watson KD, Symmons DP, Ashcroft DM: Rituximab or a second anti-tumor necrosis factor therapy for rheumatoid arthritis patients who have failed their first anti-tumor necrosis factor therapy? Comparative analysis from the British Society for Rheumatology Biologics Register. *Arthritis Care Res (Hoboken).* 2012, 64:1108-15. [10.1002/acr.21663](https://doi.org/10.1002/acr.21663)
6. Santos-Moreno P, Sánchez G, Castro C: Rheumatoid factor as predictor of response to treatment with anti-TNF alpha drugs in patients with rheumatoid arthritis. Results of a cohort study. *Medicine (Baltimore).* 2019, 98:e14181. [10.1097/MD.00000000000014181](https://doi.org/10.1097/MD.00000000000014181)
7. Bakker MF, Jacobs JW, Welsing PM, et al.: Low-dose prednisone inclusion in a methotrexate-based, tight control strategy for early rheumatoid arthritis: a randomized trial. *Ann Intern Med.* 2012, 156:329-39. [10.7326/0003-4819-156-5-201203060-00004](https://doi.org/10.7326/0003-4819-156-5-201203060-00004)
8. Rojas-Serrano J, Pérez LL, García CG, et al.: Current smoking status is associated to a non-ACR 50 response in early rheumatoid arthritis. A cohort study. *Clin Rheumatol.* 2011, 30:1589-93. [10.1007/s10067-011-1775-5](https://doi.org/10.1007/s10067-011-1775-5)
9. Saevarsdottir S, Wedrén S, Seddighzadeh M, et al.: Patients with early rheumatoid arthritis who smoke are less likely to respond to treatment with methotrexate and tumor necrosis factor inhibitors: observations from the Epidemiological Investigation of Rheumatoid Arthritis and the Swedish Rheumatology Register cohorts. *Arthritis Rheum.* 2011, 63:26-36. [10.1002/art.27758](https://doi.org/10.1002/art.27758)
10. Angum F, Khan T, Kaler J, Siddiqui L, Hussain A: The prevalence of autoimmune disorders in women: a narrative review. *Cureus.* 2020, 12:e8094. [10.7759/cureus.8094](https://doi.org/10.7759/cureus.8094)