Review began 08/18/2021

Review ended 08/24/2021

Published 09/01/2021

© Copyright 2021

Poulouse et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Correlation of Venous Clinical Severity Score With Dermatology Life Quality Index Among Patients With Chronic Venous Insufficiency: A Cross-Sectional Study

Divya Poulouse¹, Kirti Deo¹, Jaya Madhurya Gogineni¹, Aditi Mahajan¹, Samruddhi Lote¹, Roshni Mishra¹, George Jacob², Mariet Zacharias³, Abida Babu⁴

1. Department of Dermatology, Dr. D. Y. Patil Medical College, Hospital and Research Centre, Pune, IND 2. Department of Oral and Maxillofacial Surgery, Dr. D. Y. Patil Dental College and Hospital, Pune, IND 3. Department of Dermatology, Kempegowda Institute of Medical Sciences, Bangalore, IND 4. Internal Medicine, University of South Florida Morsani College of Medicine, Tampa, USA

Corresponding author: Divya Poulouse, divyapoulouse5@gmail.com

Abstract

Introduction

Chronic venous insufficiency (CVI) is characterized by inadequate functioning of venous valves in the lower limb. CVI is associated with a significant reduction in patient's quality of life (QOL). The severity of CVI was determined by CEAP (clinical, etiological, anatomical, pathophysiological) classification and venous clinical severity score (VCSS). The study is aimed to evaluate and correlate Dermatology Life Quality Index (DLQI) with VCSS, CEAP in patients with CVI.

Methods

A cross-sectional study of 57 patients with CVI was conducted over a period of 12 months. A sociographic survey, clinical and severity grading using CEAP classification, and VCSS were done for all venous doppler confirmed patients. QOL was evaluated by validated DLQI questionnaires using English and native languages Hindi and Marathi.

Results

A total of 57 patients with a male to female ratio of 6.1:1 and a mean age of 51.68 years were included in the study. CEAP grading in patients showed 49.12% (C4a), 21.05% (C6), 15.7% (C4b), 7.01% (C3), 3.50% (C2 and C5). Mean VCSS and DLQI were 11.47 and 10.12, respectively; 49.12%, 40.35%, 10.53% of patients had a moderate, very large, and small impact on DLQI respectively, positively correlating to VCSS ($P < 0.001$).

Conclusion

From this study, it was observed that VCSS and CEAP positively correlated with DLQI, and the impact increases in proportion with the seriousness of the disease.

Categories: Dermatology, Internal Medicine, General Surgery

Keywords: chronic venous insufficiency, venous clinical severity score, dermatology life quality index, venous ulcers, varicose veins, lipodermatosclerosis

Introduction

Chronic venous disease is defined as abnormalities of the venous system's morphology and functions, which are presented by various clinical signs and manifestations suggesting the necessity for investigation and management. The word chronic venous insufficiency (CVI) is used for leading signs of venous hypertension that have advanced to edema, various alterations in the skin, and venous ulcers [1]. Many factors like venous stasis, venous hypertension, fibrin cuff, and leukocyte trapping influence chronic venous insufficiency. The incidence of CVI varies from less than 1-40% in women and 1-17% in men whereas for varicose veins it is 1-73% in women and 2-56% in men [2]. Predisposing factors comprise old age, more than two pregnancies, genealogy of venous disease, obesity, female sex, sedentary lifestyle, long-standing at work causing orthostasis [2-3]. CVI in the leg causes multiple dermatological manifestations like varicose veins, venous ulcers, lipodermatosclerosis, induration, edema, venous eczema [4-6]. To standardize these manifestations of CVI, clinical, etiological, anatomical, and pathophysiological (CEAP) classification was developed by the American Venous Forum in 1994 [7-9]. CEAP lacks the ability to assess the disease severity hence a venous clinical severity scoring system (VCSS) was established by the American Venous Forum in March 2000 [9].

VCSS is a dynamic scoring system consisting of 10 components which are pain, varicose veins, edema,

How to cite this article

Poulouse D, Deo K, Gogineni J, et al. (September 01, 2021) Correlation of Venous Clinical Severity Score With Dermatology Life Quality Index Among Patients With Chronic Venous Insufficiency: A Cross-Sectional Study. Cureus 13(9): e17654. DOI 10.7759/cureus.17654

pigmentation, inflammation, induration, ulcer size, number of ulcers, duration of ulcer, and use of stockings which are graded from zero, one, two, three corresponding to absent, mild, moderate and severe respectively to produce a 30-point maximum score [10-11]. Finlay and Khan developed Dermatology Life Quality Index (DLQI) in the year 1994 and it comprises a 10-item questionnaire that covers seven domains: symptoms, feelings, daily activities, leisure items, work and school, personal relationship, treatment. Skin disorders have been acknowledged to have a detrimental effect on the quality of life of patients. This psychosocial aspect of skin disease has major implications which need to be addressed for optimal management of patients. The DLQI is estimated by totaling up the outcomes of each question, giving rise to a maximum score of 30 and a minimum score of 0, with the score being directly correlating with the impairment [12]. The study is aimed to evaluate and correlate DLQI with VCSS, CEAP in patients with chronic venous insufficiency.

Materials And Methods

This cross-sectional study was done over a period of one year from November 2019 to October 2020 among 57 patients with chronic venous insufficiency in a tertiary care center. Venous Doppler ultrasound scanning was performed in all patients after institutional ethics committee approval and informed consent from patients, VCSS was evaluated (Table 1) and clinical grading was performed using CEAP classification (Table 2). The data regarding age, sex, height, weight, BMI, occupation, level of daily physical activity, duration of standing was recorded.

Attributes	Absent = 0	Mild = 1	Moderate = 2	Severe = 3
Pain	None	Occasional	Daily not limiting	Daily limiting
Varicose veins	None	Few	Multiple	Extensive
Venous edema	None	Foot and ankle	Below knee	Knee and above
Skin pigmentation	None	Limited (old)	Diffuse (recent)	Wider (recent)
Inflammation	None	Mild cellulitis	Moderate cellulitis	Severe cellulitis
Induration	None	Focal <5 cm	<1/3 of a gaiter	>1/3 of a gaiter
Number of active ulcers	None	One	Two	More than two
Active ulcer duration (diameter)	None	<3 months	3-12 months	>1 year
Active ulcer size	None	<2 cm	2-6 cm	>6 cm
Compressive therapy	None	Intermittent	Most days	Full compliance

TABLE 1: Venous Clinical Severity Score (VCSS)

C0	No visible or palpable signs of venous disease
C1	Telangiectasias or reticular veins
C2	Varicose veins
C3	Edema
C4a	Pigmentation or eczema
C4b	Lipodermatosclerosis or Atrophie blanche
C5	Healed venous ulcer
C6	Active venous ulcer

TABLE 2: CEAP classification

CEAP: Clinical, Etiological, Anatomical, Pathophysiological Classification

The patients were given DLQI questionnaires comprising 10 open-ended questions in English and the native languages of Hindi and Marathi. Patients were asked to complete the questionnaires based on their experiences from the past week. DLQI has four different responses: 'not relevant', 'not at all', 'a little', 'a lot' or 'very much' which are graded from zero to four which then is evaluated as 0-1, 2-5, 6-10, 11-20, 21-30 which correlates to no effect, small effect, moderate effect, very large effect, extremely large effect, respectively (Table 3).

0-1	No effect at all on the patient's life
2-5	Small effect on patient's life
6-10	Moderate effect on patient's life
11-20	Very large effect on patient's life
21-30	Extremely large effect on patient's life

TABLE 3: Interpretation of DLQI score

DLQI: Dermatology Life Quality Index

All the collected data were analyzed by using SPSS V.18 (SPSS Inc., Chicago, IL). Correlation between VCSS and DLQI was assessed by Spearman's rank correlation coefficient. To investigate the relationship between VCSS and impact on quality of life (QoL), logistic regression was performed. P-value <0.05 is observed 'to indicate a statistically significant difference.

Inclusion criteria: Clinically and Venous Doppler confirmed cases of CVI who have not received treatment in the last six months.

Exclusion criteria: Age <18 years, arterial disease, deep vein thrombosis, congenital venous disease.

Results

A total of 57 patients were included in this study. The age of patients ranged from 26 to 80 years with a mean of 51.68 ± 14.53 years. Most of the patients were between the age of 40-60 years (40.35%). 26.31% and 33.33% of patients were between age 20-40 years and 60-80 years, respectively. The mean duration of the disease was 45.91 ± 9.41 months. Most of the patients were males (86%) with a ratio of male to female being 6.1:1. The average BMI was 27.18 ± 4.33 with 51% being overweight, 19% being obese and 30% being normal weight. Almost all patients gave a history of long-standing hours at work. Half of the female patients (four females) in this study had three or more pregnancies. Occupational diversification among the study group showed predominant involvement of security guards (15.8%), followed by shopkeepers (12.3%) and others (Table 4).

Profession	Number	%
Security guard	9	15.78%
Shop keeper	7	12.28%
Miscellaneous	6	10.5%
Hotel worker	5	8.77%
Housewife	5	8.77%
Driver	4	7.01%
Construction worker	4	7.01%
Conductor	3	5.26%

TABLE 4: Occupation wise distribution of patients

Among 57 patients in our study, there were two patients (3.50%) of CEAP grade C2, four patients (7.01%) of

grade C3, 28 patients (49.12%) of grade C4a, nine patients (15.7%) of grade C4b, two patients (3.50%) of grade C5, 12 patients (21.05%) of grade C6. Among the 10 aspects of VCSS, varicose veins (Figure 1) were present in all patients. Moderate pain and varicosity were the most common findings seen in 27 patients (47.37%), followed by moderate venous edema in 24 patients (42.10%). Twenty-six patients (45.61%) had moderate skin pigmentation. Moderate inflammation and moderate use of compression therapy were seen in 22 patients (38.60%) and 16 patients (28.07%), respectively. Moderate induration was seen in 14 patients (24.56%). Active leg ulcerations (Figure 2) were seen in 21.05% of patients. Single, double, and multiple ulcerations were seen in 5.26%, 10.53%, 5.26% of patients, respectively. Ulcers lasting for more than three months were seen in 10 patients (17.54%), of which 8.77% of patients had ulcers persisting for more than a year. Eight patients (14.04%) had ulcers of size more than 2 cm and four patients (7.02%) had ulcers of size less than 2 cm. Compliance with compression therapy was seen in 38 patients (66.67%), of which only one patient (1.75) showed full compliance. Distribution of different attributes of VCSS is given in Table 5.

FIGURE 1: Varicose veins in a 50-year-old male in the left lower limb

FIGURE 2: Venous ulcer with surrounding hyperpigmentation above the left medial malleolus in a 65-year-old male

Attributes	Absent = 0	Mild = 1	Moderate = 2	Severe = 3
Pain	5 (8.77%)	20 (35.09%)	27 (47.37%)	5 (8.77%)
Varicose vein	0	12 (21.05%)	27 (47.37%)	18 (31.58%)
Venous edema	5 (8.77%)	17 (29.82%)	24 (42.10%)	11 (19.30%)
Inflammation	11 (19.30%)	22 (38.60%)	22 (38.60%)	2 (3.51%)
Skin pigmentation	6 (10.53%)	16 (28.07%)	26 (45.61%)	9 (15.79%)
Induration	34 (59.65%)	2 (3.51%)	14 (24.56%)	7 (12.28%)
Number of active ulcers	45 (78.95%)	3 (5.26%)	6 (10.53%)	3 (5.26%)
Duration of ulcer (months)	45 (78.95%)	2 (3.51%)	5 (8.77%)	5 (8.77%)
Size of ulcer (cm)	45 (78.95%)	4 (7.02%)	8 (14.04%)	0
Compression therapy	19 (33.33%)	21 (36.84%)	16 (28.07%)	1 (1.75%)

TABLE 5: Distribution of different attributes of VCSS (n = 57)

VCSS: Venous Clinical Severity Score

VCSS of the patients in this study ranged from 3-28 with a mean of 11.47 ± 5.63 . DLQI of the patients ranged from 3-18 with a mean of 10.12 ± 4.16 . Six patients (10.53%), 28 patients (49.12%), 23 patients (40.35%) had a small, moderate, and very large effect on their daily lives, respectively. None of the patients had any effect or an extremely large effect on DLQI. A strong positive correlation was found between DLQI and VCSS in which the value of Spearman's rank correlation coefficient ' r ' = 0.840 and the ' p ' value obtained is <0.05 (Figure 3).

FIGURE 3: Scatter plot showing the correlation between DLQI and VCSS

DLQI: Dermatology Life Quality Index; VCSS: Venous Clinical Severity Score.

Discussion

Chronic venous insufficiency is often accompanied by distinguishing (varied)/various dermatological outcomes/features. Identification of these cutaneous findings is extremely advantageous in detecting the most frequently seen chronic vascular diseases and starting early management [1]. The CEAP classification was developed by an international consensus conference for the notification of diagnostic details in chronic venous disease, as well as for routine patient documentation and treatment [3, 13-14]. DLQI is the first dermatology-specific quality of life questionnaire published to measure dermatological distress [15-19]. In the Edinburg vein study, the prevalence of CVI is 9.4% in men and 6.6% in women [20]. In this study, the proportion of males was higher than females (6.1:1). A study done by Mallick et al. [15] in 2020 showed VCSS positively correlating with DLQI and CVI exerting small to moderate effect on patient's QoL, whereas in our study it showed moderate to very large effect. In a study done by Gillet et al. [21] on 60 patients, it was found that increased VCSS values directly correlate with increased CEAP scores. Ricci et al. evaluated the VCSS in 210 patients (420 limbs) where VCSS was 0 in 283 limbs, and VCSS was 1 or greater in the following categories: pain, varicose veins, edema, skin pigmentation, inflammation, induration, and compressive therapy in 63, 70, 51, 17, 2, 8, 9 limbs, respectively. This study was done to estimate the gravity of the disease and to substantiate the VCSS results with the ultrasonographic findings and observed the potentiality of the VCSS to act as a screening tool for CVI which could recognize patients at a higher risk for complications and with severe disease [22].

Epidemiologic studies have shown that varicose veins are more common in patients with increased body mass index (BMI) (especially $>30 \text{ kg/m}^2$) [3], as it was also seen in our study that 51% of patients being overweight and 19% being obese but 50% of patients in our study also had normal BMI indicating the role of multiple predisposing factors in the pathogenesis of CVI. In the study done by Lullove and Alvarez [23], varicosity, skin pigmentation, induration, and inflammation were the frequently noticed findings but in our study pain, varicose vein, venous edema were the most common findings. In our study, the presence of active venous ulcer was found in 21.05% of cases; amongst them, ulcers with the largest dimension of 2-6 cm (14.04%) and nonhealing ulcers persisting for 3-12 and >12 months were found in maximum patients. But the study done by Vasquez et al. found active ulcers only in 8% of cases, and small ulcers (<2 cm) and large ulcers (2-6 cm) were equally common, and nonhealing ulcers for 3-12 months were frequently observed findings. In the study done by Vasquez et al., compliance to compression therapy was higher (98.9%) as compared to our study (66.67%) [24]. This study was done to document various dermatological manifestations of chronic venous insufficiency and grade their severity with CEAP score and correlate Venous Clinical Severity Score (VCSS) with Dermatology Life Quality Index (DLQI) to prevent further complication and early management and to understand the psychosocial impact on patients. Kahn et al. [25] showed in their study that the higher the CEAP class lower was the SF-36 Physical Component Summary scores and VEINES-QOL with no effect in SF-36 mental Component Summary. In our study, it was found that VCSS showed a strong positive correlation to DLQI, and hence it can be elucidated that the higher the VCSS, the higher is the effect on the quality of life. DLQI, therefore, helps in assessing the effectiveness of treatment in the patients and their emotional aspect to the disease addressing the need to give counseling to patients regarding the same. Our study shows a positive correlation and a linear relationship between DLQI and VCSS, thereby helping the overall patient satisfaction once the issues are addressed and properly taken care of.

The main limitation in using the CEAP classification is that it responds poorly to changes over time and many of its components are relatively static and do not change significantly after treatment [25]. The advantage of using VCSS over CEAP is the avoidance of static elements and the use of only those that had the ability to reflect changes over time. It is hence recommended to use both the scoring system while evaluation rather than using one or the other [10]. In our study, we have used both the scoring systems and correlated with the quality of life. Few limitations in this study are the small sample size, which represents data from a specific region. DLQI was only translated to three languages in this study, hence limiting the patients from other linguistic states. The inclusion of control groups and a large sample size would have added more value to the study and its outcomes.

Conclusions

Chronic venous insufficiency adversely affects the quality of life by influencing the psychosocial wellbeing of the patients. In our study, mean VCSS positively correlates with DLQI in patients with CVI. Hence more severe the disease, the more it affects the DLQI. Assessment of DLQI in patients with CVI improves the effectiveness of the treatment and also addresses other psychosocial consequences of the disease which need to be managed giving more accurate insight to the clinicians.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. Institutional Ethics Committee of Dr. D. Y. Patil Medical College, Pimpri, Pune issued approval DPU / R&R (M) / 1124 / 44 / IESC / PGS / 2019 / 45 / 2019. Institutional Ethics Committee of Dr. D.Y. Patil Medical College, Pimpri, Pune has approved this study on 8th November 2019. **Animal subjects:** All authors have confirmed that this study

did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Dean SM: Cutaneous manifestations of chronic vascular disease. *Prog Cardiovasc Dis*. 2018, 60:567-579. [10.1016/j.pcad.2018.03.004](#)
- Beebe-Dimmer JL, Pfeifer JR, Engle JS, Schottenfeld D: The epidemiology of chronic venous insufficiency and varicose veins. *Ann Epidemiol*. 2005, 15:175-184. [10.1016/j.annepidem.2004.05.015](#)
- Gujja K, Sanina C, Wiley JM: Chronic venous insufficiency. In: *Interventional Cardiology: Principles and Practice*. Wiley, 2016. 759-767. [10.1002/9781118983652.ch83](#)
- Lees TA, Lambert D: Patterns of venous reflux in limbs with skin changes associated with chronic venous insufficiency. *Br J Surg*. 1993, 80:725-728. [10.1002/bjs.1800800617](#)
- Patel SK, Surowiec SM: Venous Insufficiency. StatPearls Publishing, Treasure Island, FL; 2020.
- Parsi K: Dermatological manifestations of venous disease: Part I. *Aus New Zealand J Phlebol*. 2007, 10:11-19.
- Eklöf B, Rutherford RB, Bergan JJ, et al.: Revision of the CEAP classification for chronic venous disorders: consensus statement. *J Vasc Surg*. 2004, 40:1248-1252. [10.1016/j.jvs.2004.09.027](#)
- Porter JM, Moneta GL: Reporting standards in venous disease: an update. International Consensus Committee on Chronic Venous Disease. *J Vasc Surg*. 1995, 21:635-645. [10.1016/s0741-5214\(95\)70195-8](#)
- Carpentier PH, Cornu-Thénard A, Uhl JF, Partsch H, Antignani PL: Appraisal of the information content of the C classes of CEAP clinical classification of chronic venous disorders: a multicenter evaluation of 872 patients. *J Vasc Surg*. 2003, 37:827-833. [10.1067/mva.2003.147](#)
- Rutherford RB, Padberg FT Jr, Comerota AJ, Kistner RL, Meissner MH, Moneta GL: Venous severity scoring: an adjunct to venous outcome assessment. *J Vasc Surg*. 2000, 31:1307-1312. [10.1067/mva.2000.107094](#)
- Kakkos SK, Rivera MA, Matsagas MI, Lazarides MK, Robless P, Belcaro G, Geroulakos G: Validation of the new venous severity scoring system in varicose vein surgery. *J Vasc Surg*. 2003, 38:224-228. [10.1016/S0741-5214\(03\)00323-9](#)
- Finlay AY, Khan GK: Dermatology Life Quality Index (DLQI)--a simple practical measure for routine clinical use. *Clin Exp Dermatol*. 1994, 19:210-216. [10.1111/j.1365-2230.1994.tb01167.x](#)
- Lurie F, Passman M, Meisner M, et al.: The 2020 update of the CEAP classification system and reporting standards. *J Vasc Surg Venous Lymphat Disord*. 2020, 8:342-352. [10.1016/j.jvsv.2019.12.075](#)
- Waheed SM, Kudaravalli P, Hotwagner DT: Deep vein thrombosis. In: StatPearls [Internet]. StatPearls Publishing, Treasure Island; 2020.
- Mallick S, Sarkar T, Gayen T, Naskar B, Datta A, Sarkar S: Correlation of venous clinical severity score and venous disability score with dermatology life quality index in chronic venous insufficiency. *Indian J Dermatol*. 2020, 65:489-494. [10.4103/ijd.IJD_485_20](#)
- Lewis V, Finlay AY: 10 years experience of the Dermatology Life Quality Index (DLQI). *J Invest Dermatol Symp Proc*. 2004, 9:169-180. [10.1111/j.1087-0024.2004.09113.x](#)
- Hongbo Y, Thomas CL, Harrison MA, Salek MS, Finlay AY: Translating the science of quality of life into practice: what do dermatology life quality index scores mean?. *J Invest Dermatol*. 2005, 125:659-664. [10.1111/j.0022-202X.2005.23621.x](#)
- Finlay AY: Current severe psoriasis and the rule of tens. *Br J Dermatol*. 2005, 152:861-867. [10.1111/j.1365-2133.2005.06502.x](#)
- Basra MKA, Fenech R, Gatt RM, Salek MS, Finlay AY: The Dermatology Life Quality Index 1994-2007: a comprehensive review of validation data and clinical results. *Br J Dermatol*. 2008, 159:997-1035. [10.1111/j.1365-2133.2008.08832.x](#)
- Ruckley CV, Evans CJ, Allan PL, Lee AJ, Fowkes FG: Chronic venous insufficiency: clinical and duplex correlations. The Edinburgh Vein Study of venous disorders in the general population. *J Vasc Surg*. 2002, 36:520-525. [10.1067/mva.2002.126547](#)
- Gillet JL, Perrin MR, Allaert FA: Clinical presentation and venous severity scoring of patients with extended deep axial venous reflux. *J Vasc Surg*. 2006, 44:588-594. [10.1016/j.jvs.2006.04.056](#)
- Ricci MA, Emmerich J, Callas PW, et al.: Evaluating chronic venous disease with a new venous severity scoring system. *J Vasc Surg*. 2003, 38:909-915. [10.1016/s0741-5214\(03\)00930-3](#)
- Lullove E, Alvarez OM: Improvement in clinical outcomes, physical function, and bodily pain following a 12-week course of intermittent pneumatic compression therapy in patients with chronic venous ulcers: results of an observational longitudinal retrospective study. *Cureus*. 2014, 6:175. [10.7759/cureus.175](#)
- Vasquez MA, Wang J, Mahathanaruk M, Buczkowski G, Sprehe E, Dosluoglu HH: The utility of the venous clinical severity score in 682 limbs treated by radiofrequency saphenous vein ablation. *J Vasc Surg*. 2007, 45:1008-1015. [10.1016/j.jvs.2006.12.061](#)
- Kahn SR, M'lan CE, Lamping DL, Kurz X, Bérard A, Abenhaim LA: Relationship between clinical classification of chronic venous disease and patient-reported quality of life: results from an international cohort study. *J Vasc Surg*. 2004, 39:823-828. [10.1016/j.jvs.2003.12.007](#)