Review began 08/25/2021

Review ended 08/30/2021

Published 09/06/2021

© Copyright 2021

Jaqua et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

A Common Concern With a Rare Diagnosis: Pancreatic Neuroendocrine Neoplasms

Ecler Jaqua¹, Van Nguyen¹, Vincent Pan¹, Carlos Cereser²

1. Family Medicine, Loma Linda University Medical Center, Loma Linda, USA 2. Surgical Oncology, Santa Casa de Misericórdia de Porto Alegre, Porto Alegre, BRA

Corresponding author: Ecler Jaqua, eclerjaqua@gmail.com

Abstract

Pancreatic neuroendocrine neoplasms (pNENs) are rare, representing only a small percentage of all pancreatic tumors. We report the clinical and radiological features of pNENs. Intraoperative pathology confirmed pNENs with clear margins and the patient did not require adjuvant chemoradiation. The patient is currently doing well and being closely monitored due to the high risk of relapse.

Categories: Gastroenterology, General Surgery, Oncology**Keywords:** endocrine oncology, pancreas preserving resection, gastrointestinal oncology, pancreas surgery, pancreatic neuroendocrine neoplasms

Introduction

Many patients may present to the outpatient, primary care office with abdominal pain, but its etiology is not always obvious. Abdominal pain is associated with increased morbidity whereas high body mass index, tobacco and alcohol use are not [1]. Abdominal pain even accounts for approximately 7%-10% of all emergency department concerns [2]. Although a common concern, potentially life-threatening disorders, such as neoplasm, must be excluded in the evaluation of abdominal pain. Pancreatic neuroendocrine neoplasms (pNENs), previously known as islet cell tumor, are relatively rare representing only 1%-2% of all pancreatic tumors, but has had increased prevalence in recent years due to incidental radiologic findings and physician awareness [3,4].

Case Presentation

A 63-year-old male with a history of hypertension, atrial fibrillation, obesity, and gastroesophageal reflux disease, presented with a one-week history of dull, left lower quadrant pain that radiated to the mid-epigastric region, not relieved by acetaminophen. The patient denied any associated weight changes, nausea, blood in stool, changes in bowel patterns, fever, or chills. Physical exam showed an obese abdomen with normoactive bowel sounds and tenderness to palpation diffusely across the abdomen. Due to the acuity of the pain, the patient was started on a proton-pump inhibitor and amoxicillin-clavulanic acid as empiric treatment for dyspepsia and diverticulitis. A computed tomography (CT) of the abdomen and pelvis was also ordered but did not reveal any evidence of diverticulitis. Imaging did not show any lymphadenopathy but did show a 4.6 cm nodular lesion in the body of the pancreas, suggestive of a neoplastic process, and a second 1.1 cm cystic lesion in the head of the pancreas (Figures 1-2).

How to cite this article

Jaqua E, Nguyen V, Pan V, et al. (September 06, 2021) A Common Concern With a Rare Diagnosis: Pancreatic Neuroendocrine Neoplasms. Cureus 13(9): e17767. DOI 10.7759/cureus.17767

FIGURE 1: Multiple phase CT scan abdomen and pelvis with contrast (portal-venous, early-arterial, late-arterial phase). Lesion noted to the body of the pancreas.

FIGURE 2: CT scan abdomen and pelvis with contrast; lesion enhanced. Lesion noted to the body of the pancreas.

These imaging findings were discussed with the patient and urgent laboratory work and further diagnostic imaging were ordered. Labs, including complete blood count, metabolic panel, amylase, lipase, blood sugars, carcinoembryonic antigen (CEA), and carbohydrate antigen (CA) 19-9 were unremarkable. Positron emission tomography (PET) confirmed increased metabolic activity in the body of the pancreas, measuring 3.7 x 3.5 cm.

Although increased metabolic activity was isolated to the pancreas, the patient underwent magnetic resonance imaging (MRI) to further evaluate the liver for metabolic disease. MRI was more suggestive of neuroendocrine neoplasm without obvious metastasis, but pathology would need to be obtained prior to diagnosis (Figure 3).

FIGURE 3: MRI abdomen and pelvis with contrast; lesion marked. Lesion noted to the body of the pancreas.

Approximately four weeks after presentation, the patient underwent an exploratory laparotomy, partial pancreatectomy, and total splenectomy. Pathology confirmed pancreatic neuroendocrine tumor with clear margins (Figures 4-5). The patient had a non-complicated post-operative course with normal blood sugars and was discharged home on postoperative day seven.

FIGURE 4: Surgical resection of neoplasm – partial pancreatectomy with total splenectomy.

FIGURE 5: Surgical resection of neoplasm – partial pancreatectomy with total splenectomy.

Discussion

The differentiation between functional gastrointestinal disorders and organic disease is difficult. Abdominal pain presenting with unintentional weight loss is associated with organic disease [5]. It must be acknowledged, however, that labeling abdominal pain as “chronic” usually limits the improvement of symptoms and hinders further investigation by clinicians. When used appropriately, advanced imaging, such as CT, can be ordered to rule out more sinister diagnoses such as malignancy. It is due to this imaging, usually requested to evaluate for other diagnoses, that rare neoplasms such as pNENs are increasing in incidence [3,4].

Presentation

Since pNENs can be functional or non-functional, the clinical presentation can vary depending on their type. The most common functional pNENs are insulinoma (40%-55%), gastrinoma (25%-50%), glucagonoma, and VIPoma [6]. Non-functional pNENs are asymptomatic and usually incidental findings. Symptoms of non-functional pNENs are due to tumor burden - abdominal pain (35%-78%), anorexia and nausea (45%), and weight loss (22%-35%) [7]. The majority of pNENs are sporadic cases but a small minority are associated with hereditary diseases such as multiple endocrine neoplasia type 1, von Hippel Lindau, neurofibromatosis type 1, and tuberous sclerosis [8]. pNENs tend to have multifocal metastases. The most common site of metastases is to the liver, lungs, bone, peritoneum, adrenal glands, and spleen [9].

Diagnosis

Due to its wide availability and high sensitivity and specificity, CT has become the diagnostic imaging of choice for pNENs. Helical (spiral), multiphasic, or contrast-enhanced CT should be used as pNENs are highly vascular. MRI may be a favorable test to characterize smaller pancreatic lesions and metastasis to the liver [10]. PET can be used to further localize neoplasms [11]. The last modality that can be used, endoscopic ultrasonography (EUS), can be used to provide high-resolution imaging of the pancreas, with higher sensitivity than CT. The other benefit of EUS is that fine-needle aspiration can be done at the same time as imaging [12].

Treatment

Small pNENs, less than 2 cm, can be followed with observation. Surgical resection is still the mainstay of treatment, a potentially curative option for pNENs [13]. Localization of pNENs, as well as the grade and biological activity of the tumor, lymph node involvement, metastasis, surgical margins, tumor size, and macro/microvascular involvement, affect surgical cure rate and overall prognosis. Due to the high relapse rate of pNENs, adjuvant systemic therapy such as chemotherapy, somatostatin analogs, targeted inhibitor therapies, and peptide receptor radionuclide therapy should be considered.

Prognosis

Resection of localized pNENs has a five-year survival rate of 55%. Non-resectable pNENs have a five-year survival rate of 15% [14]. TNM system (tumor, node and metastasis) is the most common and recognized cancer staging system. pNENs can also be staged by their grade (G) and degree of differentiation going from G1 to G3. G1-grade tumors, in comparison to G3-grade tumors, have a median survival of twelve years and ten months, respectively [14]. The higher the tumor grade is, the more poorly differentiated cells are and consequently worsen prognosis.

Conclusions

Patients with abdominal pain and perhaps unexplained weight loss should be suspected of malignancy and undergo diagnostic imaging via the widely available, and highly sensitive and specific CT. In the case of localized pNENs, surgical treatment can be curative but may still require adjuvant therapies due to its high frequency of relapse.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Wallander MA, Johansson S, Ruigómez A, García Rodríguez LA: Unspecified abdominal pain in primary care: the role of gastrointestinal morbidity. *Int J Clin Pract.* 2007, 61:1663-70. [10.1111/j.1742-1241.2007.01529.x](https://doi.org/10.1111/j.1742-1241.2007.01529.x)
2. Powers RD, Guertler AT: Abdominal pain in the ED: stability and change over 20 years. *Am J Emerg Med.* 1995, 13:301-3. [10.1016/0735-6757\(95\)90204-X](https://doi.org/10.1016/0735-6757(95)90204-X)
3. Heitz PU, Komminoth P, Perren A, et al.: Pancreatic endocrine tumours: introduction. *Pathology and Genetics of Tumours of Endocrine Organs.* DeLellis RA, Lloyd RV, Heitz PU, Eng C (ed): IARC Press, Lyon; 2004. 177-182.
4. Hallet J, Law CH, Cukier M, Saskin R, Liu N, Singh S: Exploring the rising incidence of neuroendocrine tumors: a population-based analysis of epidemiology, metastatic presentation, and outcomes. *Cancer.* 2015, 15:589-97. [10.1002/cncr.29099](https://doi.org/10.1002/cncr.29099)
5. Vandvik PO, Kristensen P, Aabakken L, Farup PG: Abdominal complaints in general practice. *Scand J Prim Health Care.* 2004, 22:157-62. [10.1080/02813430410006503](https://doi.org/10.1080/02813430410006503)
6. Perri G, Prakash LR, Katz MH: Pancreatic neuroendocrine tumors. *Curr Opin Gastroenterol.* 2019, 35:468-77. [10.1097/MOG.0000000000000571](https://doi.org/10.1097/MOG.0000000000000571)
7. UpToDate: classification, epidemiology, clinical presentation, localization, and staging of pancreatic neuroendocrine neoplasms. (2021). Accessed: July 10, 2021: <https://www.uptodate.com/contents/classification-epidemiology-clinical-presentation-localization-and-staging-of-pancr....>
8. Vinik A, Perry RR, Casellini C, Hughes MS, Feliberti E: Pathophysiology and Treatment of Pancreatic Neuroendocrine Tumors (PNETs): New Developments. Feingold KR, Anawalt B, Boyce A, et al (ed): Endotext, South Dartmouth (MA); 2000.
9. Zerbi A, Falconi M, Rindi G, et al.: Clinicopathological features of pancreatic endocrine tumors: a prospective multicenter study in Italy of 297 sporadic cases. *Am J Gastroenterol.* 2010, 105:1421-9. [10.1038/ajg.2009.747](https://doi.org/10.1038/ajg.2009.747)
10. Dromain C, de Baere T, Lumbroso J, et al.: Detection of liver metastases from endocrine tumors: a prospective comparison of somatostatin receptor scintigraphy, computed tomography, and magnetic resonance imaging. *J Clin Oncol.* 2005, 23:70-8. [10.1200/JCO.2005.01.013](https://doi.org/10.1200/JCO.2005.01.013)
11. Hofman MS, Kong G, Neels OC, Eu P, Hong E, Hicks RJ: High management impact of Ga-68 DOTATATE (GaTate) PET/CT for imaging neuroendocrine and other somatostatin expressing tumours. *J Med Imaging Radiat Oncol.* 2012, 56:40-47. [10.1111/j.1754-9485.2011.02327.x](https://doi.org/10.1111/j.1754-9485.2011.02327.x)
12. James PD, Tsolakis AV, Zhang M, Bellettrutti PJ, Mohamed R, Roberts DJ, Heitman SJ: Incremental benefit of preoperative EUS for the detection of pancreatic neuroendocrine tumors: a meta-analysis. *Gastrointest Endosc.* 2015, 81:848-56. [10.1016/j.gie.2014.12.031](https://doi.org/10.1016/j.gie.2014.12.031)
13. Falconi M, Eriksson B, Kaltsas G, et al.: ENETS consensus guidelines update for the management of patients with functional pancreatic neuroendocrine tumors and non-functional pancreatic neuroendocrine tumors. *Neuroendocrinology.* 2016, 103:153-71. [10.1159/000443171](https://doi.org/10.1159/000443171)
14. Sobin LH, Gospodarowicz MK, Wittekind C: TNM Classification of Malignant Tumours. Wiley-Blackwell, Hoboken, NJ; 2009.