Upward Migration and Coiling of the Distal Catheter Toward the Valve Site

Review began 09/01/2021
Review ended 09/05/2021
Published 09/15/2021

© Copyright 2021

Alghamdi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Khalid Alghamdi ¹, Luma H. Kutub ², Ahmed G. Qasem ³, Abdulrahman H. Kaneetah ⁴, Sultan G. Alzahrani ⁵, Hussam Y. Kutub ⁶

1. College of Medicine, King Saud bin Abdulaziz University for Health Sciences, King Abdullah International Medical Research Centre, King Abdulaziz Medical City, National Guard Health Affairs, Jeddah, SAU 2. College of Medicine, Batargi Medical College, Jeddah, SAU 3. Neurosurgery, King Fahad Hospital, Medinah, SAU 4. College of Medicine, King Saud bin Abdulaziz University for Health Sciences, King Abdullah International Medical Research Centre, King Abdulaziz Medical City, National Guard Health Affairs, Jeddah, SAU 5. College of medicine, King Saud bin Abdulaziz University for Health Sciences, King Abdullah International Medical Research Centre, King Abdulaziz Medical City, National Guard Health Affairs, Jeddah, SAU 6. Neurosurgery, King Abdulaziz Medical City, Ministry of National Guards Health Affairs, Jeddah, SAU

Corresponding author: Hussam Y. Kutub, hussamkutub@yahoo.com

Abstract

Hydrocephalus, which is caused by the accumulation of cerebrospinal fluid (CSF), is a common condition in children. It is known to be most likely treated by the insertion of a ventriculoperitoneal (VP) shunt. However, a VP shunt can lead to multiple complications. The upward migration of a VP shunt is considered rare.

A newborn male baby with a known case of Chiari malformation type 2 associated with myelomeningocele (MMC) and hydrocephalus had a VP shunt inserted for control of the hydrocephalus. He presented two months after the surgery with occipital swelling at the surgical site. Shunt series followed by Computerized tomography (CT) scan showed that the distal end of the catheter had migrated upward and coiled around the valve. Urgent revision of the VP shunt was performed.

Reabsorption of subgaleal fluid, increased abdominal pressure, repeated abdominal wall contraction, and repeated head motion of the child are the previously suggested theories of upward migration of distal catheter to the site of the valve. However, the combination of multiple theories can be the logical explanation, as they do not oppose each other.

Categories: Neurosurgery

Keywords: migration and coiling of ventriculoperitoneal shunt, ventriculoperitoneal shunt placement, pediatric hydrocephalus, upward migration of ventriculoperitoneal shunt, ventriculoperitoneal shunt complications

Introduction

Hydrocephalus is considered a common but serious condition in children caused by the accumulation of cerebrospinal fluid (CSF) [1]. The management of hydrocephalus is primarily surgical by insertion of a ventricular shunt [2]. The shunt diverts the accumulated CSF to another part of the body where it can be absorbed [2]. The most common type of shunt is a ventriculoperitoneal (VP) shunt [2]. However, several complications of the VP shunt have been reported, such as obstruction, over drainage of CSF, intracranial infection, seizure, abdominal injury, and visceral perforation [3]. The risk of complications from a VP shunt varies between 11% and 25% during the first year after surgery [4]. The subgaleal migration of VP shunt is considered a type of upward migration, and it is quite a rare complication [5]. We present a rare case of a patient with occipital swelling due to subgaleal upward migration of VP shunt, who was successfully treated.

Case Presentation

A newborn male baby with a known case of Chiari malformation type 2 associated with myelomeningocele (MMC) had a repair of MMC after three days of delivery. The patient was then kept in a prone position. On the first day post MMC repair, his head circumference was 33.5 cm. Ten days post MMC repair, the patient was vitally stable and the fontanelle was soft and flat with head circumference increased by 2 cm (35.5), and computerized tomography (CT) scan of the brain showed interval increase in the size of the lateral ventricles. Hence, the patient underwent a VP shunt. During the surgery, a ventricular Bactiseal catheter was used and inserted to a depth of 4.5 cm, and CSF came in high pressure. The ventricular catheter was attached to a burr hole low-pressure valve. Adequate functioning of the shunt was confirmed, and there was good distal flow. The distal catheter was then inserted into the peritoneal region with sufficient length (20 cm). The patient was then transported back to the neonatal intensive care unit (NICU) in a stable condition (Figure 1).

How to cite this article

Alghamdi K, Kutub L H, Qasem A G, et al. (September 15, 2021) Upward Migration and Coiling of the Distal Catheter Toward the Valve Site. Cureus 13(9): e17993. DOI 10.7759/cureus.17993

FIGURE 1: After the first operation, chest X-ray shows the extension of the catheter from the neck toward the peritoneal cavity.

After one month, the patient presented to the neurosurgery clinic with swelling in the occipital region. Shunt series followed by CT brain showed extracranial CSF collection surrounding the valve of the right VP shunt with retraction and coiling of the distal catheter within the collection (Figure 2). Furthermore, an interval increase in the size of the lateral ventricles was noted, and the patient was booked for VP shunt revision the next day.

FIGURE 2: Before the second operation, VP shunt series, lateral (A) and anterior (B) skull view, show upward migration and coiling of the distal catheter. Chest X-ray (C) shows an absence of the distal catheter in the abdomen.

During shunt revision, the cranial incision was opened, and hypertensive flow was observed. The distal catheter was found coiled freely in the subgaleal space (Figure 3). Both the distal and proximal components of the ventriculoperitoneal shunt system were revised, and clear CSF was observed coming out from the distal catheter. The length of the distal catheter was increased and purse-string suture was used to prevent the recurrence of the same complication. Post VP shunt revision, the patient was vitally stable with a sunken anterior fontanelle.

FIGURE 3: Images of the patient before incision (A) and after incision (B) show subgaleal swelling and coiling of the distal catheter at the valve site.

Discussion

The complications of shunt surgery can be divided into infection, functional failure, and mechanical failure which include disconnection, obstruction, and migration of any part of the shunt system [6]. Shunt migration seems to be more common in children than in adults because of the length of the distal catheter and repeated head motion of the child [7]. Studies have shown migration of the distal catheter to multiple sites such as the heart, scrotum, chest, bladder, liver, colon, and anus. However, few studies reported cases of upward migration of the distal catheter of the VP shunt to the subgaleal space [6, 7, 8, 9, 10, 11]. The cases vary in their shunt system and whether they used the valve system or not, so the association between the device type and the development of upward migration was not found. Although subgaleal migration is a rare complication, many hypotheses have been reported, including reabsorption of subgaleal fluid, increased abdominal pressure, the repeated head motion of the child, and repeated abdominal wall contraction [8, 12, 13, 14]. In our case, the patient was kept in a prone position for 28 days after MMC repair, and that can lead to increase abdominal pressure. The increase in abdominal pressure has been explained by a study by Abou el Nasr, as it would lead to pushing force on the distal catheter [12]. Other hypothesis includes reabsorption of

subgaleal fluid, which leads to the development of negative pressure that drags the distal catheter as explained by Pang et al. [13]. Scott et al. demonstrated "the windlass effect", which considers the repeated head motion of the child as the reason behind the upward migration of distal catheter [14]. The prevention of upward migration cannot be guaranteed, so securing a shunt device has been recommended [7]. In our case, a longer distal catheter (30 cm) was secured to the surrounding fascia to prevent the recurrent upward migration.

Conclusions

The upward migration of the distal catheter to the valve site is considered rare. There are many speculated theories that can explain it. However, a combination of these theories may appear to be the logical explanation, as they do not oppose each other. Increased abdominal pressure and repeated head motion were considered to pull the distal catheter to the valve site at the subgaleal space. Further reporting of similar cases is required as it will increase the understanding of this complication.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Rekate HL: A contemporary definition and classification of hydrocephalus. *Semin Pediatr Neurol.* 2009, 16:9-15. [10.1016/j.spen.2009.01.002](https://doi.org/10.1016/j.spen.2009.01.002)
2. Fowler JB, De Jesus O, Mesfin FB: *Ventriculoperitoneal Shunt.* StatPearls (ed): StatPearls Publishing, Treasure Island (FL); 2021.
3. Hung AL, Vivas-Buitrago T, Adam A, et al.: Ventriculoatrial versus ventriculoperitoneal shunt complications in idiopathic normal pressure hydrocephalus. *Clin Neurol Neurosurg.* 2017, 157:1-6. [10.1016/j.clineuro.2017.03.014](https://doi.org/10.1016/j.clineuro.2017.03.014)
4. Hidayat I, Syahputra DA, Isa MM: Unusual migration of distal ventriculoperitoneal shunt to vagina via fallopian tube: a case report. *Ann Med Surg (Lond).* 2021, 63:102158. [10.1016/j.amsu.2021.02.004](https://doi.org/10.1016/j.amsu.2021.02.004)
5. Harischandra LS, Sharma A, Chatterjee S: Shunt migration in ventriculoperitoneal shunting: a comprehensive review of literature. *Neurol India.* 2019, 67:85-99. [10.4103/0028-3886.253968](https://doi.org/10.4103/0028-3886.253968)
6. Al Hinai QS, Pawar SJ, Sharma RR, Devadas RV: Subgaleal migration of a ventriculoperitoneal shunt. *J Clin Neurosci.* 2006, 13:666-9. [10.1016/j.jocn.2005.07.020](https://doi.org/10.1016/j.jocn.2005.07.020)
7. Cho KR, Yeon JY, Shin HJ: Upward migration of a peritoneal catheter following ventriculoperitoneal shunt. *J Korean Neurosurg Soc.* 2013, 53:383-5. [10.3340/jkns.2013.53.6.383](https://doi.org/10.3340/jkns.2013.53.6.383)
8. Yee GT, Han SR, Choi CY: Migration and coiling of peritoneal catheter into the subgaleal space: a very rare complication of subgaleoperitoneal shunt. *J Korean Neurosurg Soc.* 2013, 54:525-7. [10.3340/jkns.2013.54.6.525](https://doi.org/10.3340/jkns.2013.54.6.525)
9. Heim RC, Kaufman BA, Park TS: Complete migration of peritoneal shunt tubing to the scalp. *Childs Nerv Syst.* 1994, 10:399-400. [10.1007/BF00335131](https://doi.org/10.1007/BF00335131)
10. Dominguez CJ, Tyagi A, Hall G, Timothy J, Chumas PD: Sub-galeal coiling of the proximal and distal components of a ventriculo-peritoneal shunt. An unusual complication and proposed mechanism. *Childs Nerv Syst.* 2000, 16:493-5. [10.1007/pl00007294](https://doi.org/10.1007/pl00007294)
11. Kloss BT, Hart DM, Secreti L: Subgaleal coiling of the proximal and distal components of a ventriculoperitoneal shunt. *Int J Emerg Med.* 2012, 5:15. [10.1186/1865-1380-5-15](https://doi.org/10.1186/1865-1380-5-15)
12. Abou el Nasr HT: Modified method for prophylaxis against unishunt system complications with presentation of total intraventricular migration of unisystem ventriculoperitoneal shunt. *Childs Nerv Syst.* 1988, 4:116-8. [10.1007/BF00271393](https://doi.org/10.1007/BF00271393)
13. Pang D, Wilberger JE Jr: Upward migration of peritoneal tubing. *Surg Neurol.* 1988, 14:363-4.
14. SC M, WY HT, MU F, RE V: Observations on ventricular and lumbar subarachnoid peritoneal shunts in hydrocephalus in infants. *J Neurosurg.* 1955, 12:165-75. [10.3171/jns.1955.12.2.0165](https://doi.org/10.3171/jns.1955.12.2.0165)