For Residents, by Residents: Developing a Physician Handoff Tool at a University Affiliated Community Hospital

Received 09/05/2021

Review began 09/10/2021

Review ended 09/28/2021

Published 09/28/2021

© Copyright 2021

Chen et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Lu Chen¹, Uta S. Guo², Siddharth Bhesania^{3,4}, Hardikkumar Shah⁵, Emdad Ali⁶, Parag Mehta⁴

1. Department of Cardiology, State University of New York Downstate Medical Center, Brooklyn, USA 2. Department of Family Medicine and Community Health, Icahn School of Medicine at Mount Sinai, New York, USA 3. Internal Medicine, Overlook Medical Center, Summit, USA 4. Internal Medicine, NewYork-Presbyterian Brooklyn Methodist Hospital, Brooklyn, USA 5. Gastroenterology and Hepatology, Saint Joseph's University Medical Center, Paterson, USA 6. Internal Medicine, OhioHealth Physician Group, Columbus, USA

Corresponding author: Lu Chen, lu.chen@downstate.edu

Abstract

The 80-hour per week work limit resulted in an increased number of patient handoffs. A satisfactory handoff system should optimize the exchange of vital patient information while concisely minimizing error. This project describes our experience and lessons learned in successfully developing and implementing an Electronic Health Record (EHR)-integrated handoff system based on the I-PASS model. The handoff system, termed Physician Handoff, was refined through end-user feedback. End-users were evaluated on the quality of handoff in the following categories: Illness Severity, Patient Summary, Action List, and Situational Awareness. Resulting survey showed high adoption and satisfaction rate with Physician Handoff. Success can be attributed to interdepartmental collaboration, credentialing the users, and recognizing the importance of end-user feedback.

Categories: Internal Medicine, Medical Education, Healthcare Technology

Keywords: physician satisfaction, patient safety, electronic medical records, standardized handoff, graduate medical education (gme)

Introduction

The devil is in the details. No matter how much the importance of communication is emphasized in our training; errors in communication is one of the leading causes of sentinel events reported [1]. This is critical in graduate medical education. Since the 80 duty-hour per week restriction was put into effect [2], there is an increased number of handoffs performed by resident physicians [3]. Compared to 2003, there has been a 40% increase in handoffs performed by physicians. On average, a resident physician may be involved in over 300 handoffs during a month-long rotation. A patient would be handed off 15 times over a five-day hospital course [4]. Unfortunately, the communication during handoff can overly rely on written handoffs, and the effectiveness of communication is often overestimated [5,6]. Boosted confidence level and increased numbers of handoffs can make handoffs more vulnerable to errors.

Previous studies found that four out of five handoff sheets contain at least one error, with the most common error being medication omission [7]. Additionally, half of the patient handoff documents become inaccurate or outdated within six hours on an average night shift, mostly secondary to medication changes. By the following morning, merely 40% of handoff documents were still current [8]. A few studies have noted an association between handoff quality and patient outcomes. In a critical care setting, more structured handoffs were associated with a decreased number of non-routine clinical events [9]. In the surgical arena, the transition from written to electronic handoff resulted in a reduction in length of stay [10]. Another study described residents inappropriately performing resuscitation on multiple patients due to discrepancies found on handoff sheets. There were also patients who were supposed to be resuscitated but were denied resuscitations due to handoff errors [11].

In order to minimize errors and prevent adverse clinical events, it is necessary that the information on handoff documents is accurate, concise, and complete. Ideally, a handoff system should be fully integrated into the Electronic Health Records (EHR). When available, clinical information in the handoff system should be generated and updated automatically to minimize manual input and errors. A previous study has shown that automatically generated handoff increased the completeness of handoff, minimized errors, and streamlined physician workflow [12,13]. Using EHR-integrated handoffs is associated with a reduction in length of stay for the patients and improvement in provider satisfaction [14]. Early adapters of EHR-integrated handoff found it consistent and generalizable across different study sites [15]. To improve patient safety and comply with Accreditation Council for Graduate Medical Education (ACGME) regulation, our hospital has developed and implemented an EHR-integrated handoff system that provides clear, accurate, and concise patient handoff sheets. The purpose of this paper is to describe our experience and the lessons we have learned in designing a satisfactory handoff system for resident physicians in our hospital.

How to cite this article

Chen L, Guo U S, Bhesania S, et al. (September 28, 2021) For Residents, by Residents: Developing a Physician Handoff Tool at a University Affiliated Community Hospital. Cureus 13(9): e18352. DOI 10.7759/cureus.18352

Technical Report

Previous versions of our handoffs

Patient handoff was previously conducted verbally and supplemented with either written index cards or manually inputted handoff into word-processing software per provider preference. Since the implementation of EHR (Cerner Corporation, Kansas City, MO) at our hospital in 2011, efforts were made to integrate electronic handoff into the EHR. The Information Technology (IT) department and administration integrated Physician Worklist, an existing built-in handoff program into our EHR system (Figure 1). Physician Worklist was designed to be comprehensive and automated, which included information such as code status, vital signs, diet, allergy, diagnosis, active problems, medications, and laboratory results. Unfortunately, Physician Worklist did not include physician assessment, plan of care, and patient disposition. The volume of information limited number of patients per printed page, resulting in 40-60-page sign out packets for the night team residents. End-user survey following implementation showed poor adoption rate and low satisfaction with Physician Worklist.

Physician handoff

A decision was made to implement a more user-friendly handoff system. Two senior resident physicians were elected to lead the project. They were designated as resident liaisons between the Internal Medicine (IM) leadership, the IT department, and fellow resident physicians. The newly developed handoff system was structured according to a published handoff system, which has been linked to reduction in medical errors and adverse events [16-18]. The published handoff system is known for its mnemonic I-PASS (Illness Severity, Patient Summary, Action List, Situational Awareness, and Synthesis by Receiver). We adapted the I-PASS model and termed our new handoff Physician Handoff.

The patient summary section in Physician Handoff was limited to 500 characters, allowed more patients to fit per printed page. Action list allowed providers to create an individualized task list. Physician Handoff auto-populated Code Status, Diagnosis, Physician Contact, and Admission Date, further minimizing input errors (Figure 2). Laboratory finding and medications were omitted from the printed version, but can be easily accessed on computerized version. Handoff can be reviewed, verified, and modified by any member of the care team at any secured workstation with EHR access, ensuring secured and real-time patient information sharing.

Printed: John Smith MD

Physician Handoff

Patient Information			
Patient: Jane Doe 86 Years, Female Primary Contact: John Doe	Location: ED-1 Code Status: Full	MRN: 123456 Admit Date: 3-23-2050	Illness Severity: Stable
Patient Summary 86yo F, PMH of Bipolar, depression, intermittent asthma, presented with cough, DOE, admitted for RLL PNA.			
Situational Awareness and Planning Full Code, Home with services			
Action [] Follow up cultures [] Update son of progress			
Patient Information			
Patient: John Doe 66 Years, Male Primary Contact:	Location: 1025A Code Status: DNR/DNI	MRN: 654321 Admit Date: 9-12-2048	Illness Severity: Watcher
Patient Summary 66yo AAM, h/o CAD s/p PCI, HFrEF, s/p AICD, HTN, HLD, DM, presented with progressive DOE x3 days, +trop, +PND, admitted for ADHF.			
Situational Awareness and Planning DNR DNI, From Home with services			
Action [] Extra dose of IV Lasix 40 if needed for goal 24hr UOP of 2000cc			
Patient Information			
Patient: Joan Doe 46 Years, Female Primary Contact:	Location: 855B Code Status: Full	MRN: 662461 Admit Date: 8-14-2041	Illness Severity: Discharging
Patient Summary 46yo F, G2P2, PMH of UTIs, p/w LLQ/suprapubic tenderness, +UA, +dysuria, found to have pan sensitive e coli, on ceftriaxone x1 days, likely DC tmr			
Situational Awareness and Planning Full Code			
Action [] DC summary drafted			

If found, please return to the nearest ward clerk

Page: 1 of 2

FIGURE 2: Sample of Printed Physician Handoff

ED: emergency department; MRN: medical records number; F: female; PMH: past medical history; DOE: dyspnea on exertion; RLL: right lower lobe; PNA: pneumonia; DNR/DNI: do not resuscitate/do not intubate; AAM: African American male; CAD: coronary artery disease; PCI: percutaneous coronary intervention; HFrEF: heart failure reduced ejection fraction; AICD: automatic implantable cardioverter defibrillator; HTN: hypertension; HLD: hyperlipidemia; DM: diabetes mellitus; DOE: dyspnea on exertion; PND: paroxysmal nocturnal dyspnea; ADHF: acute decompensated heart failure; IV: intravenous; UOP: urine output; G2P2: Gravidity 2 Parity 2; UTI: urinary tract infection; LLQ: left lower quadrant; UA: urinalysis; DC: discharge

Implementation of Physician Handoff required multiple rounds of fine-tuning due to the need of different departments. Following the development of Physician Handoff, all residents were trained and credentialed in the new handoff system. Residents were evaluated on both written and verbal handoffs. Anonymous surveys following implementation showed a better adoption rate and higher satisfaction rate among resident physicians.

Discussion

Developing, implementing, and evaluating an efficient EHR-integrated handoff system was a momentous undertaking for all individuals involved (IM leadership, IT department, and the resident physicians). Valuable lessons were learned during this process: 1) interdepartmental collaboration; 2) credential the users, and most importantly, 3) value end-user's opinions and workflows. By applying the knowledge gained from these lessons, we were able to develop and implement Physician Handoff, a widely accepted system that brought many unforeseen benefits.

Over the span of three years, two EHR-integrated handoff systems were introduced in our hospital to replace manually inputted handoff. Physician Worklist focused on completeness but failed to incorporate end-user feedback. In contrast, Physician Handoff was designed by resident physicians with an emphasis on end-user experience. Differences among the three handoff systems are summarized in Table 1.

	Manual Handoff	Physician Worklist	Physician Handoff
Designed by	Residents	IT & IM Leaders	IT, IM Leaders, Resident
Resident Feedback	Incorporated	Not Incorporated	Incorporated
EHR Integration	No	Yes	Yes
Availability of Access	Single Workstation	Hospital Wide	Hospital Wide
HIPAA Compliance	No	Yes	Yes
Content	Varies	Comprehensive	Concise
Free Text Option	Yes	Limited	Yes
Automation	No	Yes	Yes
Formalized Training	No	No	Yes

TABLE 1: Handoffs At a Glance: Comparison of Manual Handoff, Physician Worklist, and Physician Handoff

IM: Internal Medicine; IT: Information Technology; EHR: Electronic Health Record; HIPAA: Health Insurance Portability and Accountability Act

Lesson 1: Interdepartmental collaboration

The successful development of Physician Handoff can be attributed to the perseverance and collaboration of the multidisciplinary team. Through multiple setbacks, we enacted the Plan-Do-Study-Act (PDSA) workflow, which included: working closely with the IT department and EHR vendor, assigning senior residents as liaisons, and holding discussions with IM leadership.

IM Departmental leadership served as an oversight to the project but allowed resident liaisons and IT department to refine the details. This style of leadership maximized incorporation of residents’ ideas within a set framework. Feedbacks from numerous attending physicians were also conveyed through the department leadership. Resident liaisons serve the crucial role of learning and summarizing the opinion of resident physicians. They voiced the concerns of residents in a concise, and non-confrontational manner. The suggestions for modifications from any party were only incorporated into the handoff design upon agreement by all three parties (IM, IT, and liaisons). Our IT department provided unique insight from the design aspect and troubleshoot various technical challenges in incorporating our new handoff system. Their innovative designs were a novice to many clinicians and made them invaluable members of the design team.

Lesson 2: Credential of the users

Once Physician Handoff debuted, the resident liaisons and the IM leadership were certified as super users of the new system. The super users worked together and designed a complete but concise training course for the remaining users to allow a seamless transition. A formal training ensured that the end-users had a uniform and adequate understanding of not only the I-PASS handoff model but also Physician Handoff program and the services it offered. Following the credentialing course, resident liaisons personally observed and assessed individual handoffs based on all categories of the I-PASS model. The success was contributed to properly credentialing of the end-users and perseverance of the liaisons.

Lesson 3: Value end-user’s opinions and workflows

The development of Physician Worklist neither include end-user feedback, nor did it provide the components essential to resident physicians (plan of care, disposition, and task list). Our study is consistent with the previous studies that have demonstrated the importance of collaborating with end-users in both the development and implementation process of handoff systems [19-22]. Building on user feedback, the inclusion of the “Comment” and “Actions” sections allowed for a personalized aspect of health care to be communicated between providers. Physician Handoff displayed information vital to providers caring for the patient. Due to its conciseness, more patients were listed per page. Because the opinions of the end-users were addressed and incorporated, Physician Handoff was better adapted with a high satisfaction rate.

Physician handoff: A step in the right direction

Physician Handoff is a concise and EHR-integrated handoff system that maximized automation while allowed freedom of manual input. By automatically updating code status and prominently displaying it on the handoff sheet, this build-in feature minimizes errors associated with manual input and can potentially limit inappropriate resuscitations. Physicians identified illness severity on handoff helped in better allocation of resources for both nursing and supporting staff. Additional advantages of an EHR-integrated handoff system include enhancing compliance with Health Insurance Portability and Accountability Act (HIPAA) and providing accurate information transfer. EHR-integrated handoff tools can ensure a secure hospital-wide exchange of patient information, preserve the continuity of care, and prevent potential adverse events in our patients. The effect of this implementation on our patient care is yet to be determined.

Our experience with implementing a standardized and EHR-integrated handoff system echoed the experience of many previous studies. Lessons we have learned independently mirrored the recommendation of a recent literature review [23]. In our experience, collaboration, improvisation, adequate credentialing, and the inclusion of end-users in the design and revision efforts were the roots of our success.

Conclusions

In an effort to improve patient safety we developed a satisfying and widely accepted handoff system. Developing an EHR-integrated handoff tool was a monumental undertaking, requiring coordination between IM departmental leadership, IT Department, and end-users. Our mission was to minimize handoff errors in Physician Handoff by automating objective data and limit manual input. We attributed that success to collaborating, improvising, credentialing the users, and recognizing the importance of end-user feedback.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. NewYork-Presbyterian Brooklyn Methodist Hospital Institutional Review Board issued approval n/a. Study was reviewed and exempted by Internal Review Board at our hospital. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

The authors would like to thank the physicians, physicians in training, and staff members at NewYork-Presbyterian Brooklyn Methodist Hospital, Departments of Medicine and Information Technology for their help and support for this project. Authors would like to especially thank Dr. Eric Karu for his contributions to this project.

References

1. Dingley C, Daugherty K, Derieg MK, Persing R: Improving patient safety through provider communication strategy enhancements. *Advances in Patient Safety: New Directions and Alternative Approaches*. Henriksen K, Battles JB, Keyes MA, Grady ML (ed): Agency for Healthcare Research and Quality, Rockville, MD; 2008. 3: 10.1056/NEJMp1011413
2. Drolet BC, Spalluto LB, Fischer SA: Residents' perspectives on ACGME regulation of supervision and duty hours--a national survey. *N Engl J Med*. 2010, 363:e34. 10.1056/NEJMp1011413
3. DeRienzo CM, Frush K, Barfield ME, et al.: Handoffs in the era of duty hours reform: a focused review and strategy to address changes in the Accreditation Council for Graduate Medical Education Common Program Requirements. *Acad Med*. 2012, 87:403-10. 10.1097/ACM.0b013e318248e5c2
4. Okie S: An elusive balance--residents' work hours and the continuity of care. *N Engl J Med*. 2007, 356:2665-7. 10.1056/NEJMp078085
5. Chang VY, Arora VM, Lev-Ari S, D'Arcy M, Keysar B: Interns overestimate the effectiveness of their hand-off communication. *Pediatrics*. 2010, 125:491-6. 10.1542/peds.2009-0351
6. Blondon K, Zotto MD, Roach J, Nendaz MR, Lovis C: A simulation study on handoffs and cross-coverage: results of an error analysis. *AMIA Annu Symp Proc*. 2018, 2017:448-57.
7. Arora V, Kao J, Lovinger D, Seiden SC, Meltzer D: Medication discrepancies in resident sign-outs and their potential to harm. *J Gen Intern Med*. 2007, 22:1751-5. 10.1007/s11606-007-0415-x
8. Rosenbluth G, Jacobia R, Milev D, Auerbach AD: Half-life of a printed handoff document. *BMJ Qual Saf*. 2016, 25:324-8. 10.1136/bmjqs-2015-004585
9. Nanchal R, Aebly B, Graves G, et al.: Controlled trial to improve resident sign-out in a medical intensive care unit. *BMJ Qual Saf*. 2017, 26:987-92. 10.1136/bmjqs-2017-006657
10. Ryan S, O'Riordan JM, Tierney S, Conlon KC, Ridgway PF: Impact of a new electronic handover system in surgery. *Int J Surg*. 2011, 9:217-20. 10.1016/j.ijsu.2010.11.012
11. Vidyarthi AR, Arora V, Schnipper JL, Wall SD, Wachter RM: Managing discontinuity in academic medical

- centers: strategies for a safe and effective resident sign-out. *J Hosp Med*. 2006, 1:257-66. [10.1002/jhm.103](#)
12. Tisdale RL, Eggers Z, Shieh L: EMR-based handoff tool improves completeness of internal medicine residents' handoffs. *BMJ Open Qual*. 2018, 7:e000188. [10.1136/bmjopen-2017-000188](#)
13. Bernstein JA, Imler DL, Sharek P, Longhurst CA: Improved physician work flow after integrating sign-out notes into the electronic medical record. *Jt Comm J Qual Patient Saf Jt Comm Resour*. 2010, 36:72-8. [10.1016/S1553-7250\(10\)36013-2](#)
14. Palma JP, Sharek PJ, Longhurst CA: Impact of electronic medical record integration of a handoff tool on sign-out in a newborn intensive care unit. *J Perinatol*. 2011, 31:311-7. [10.1038/jp.2010.202](#)
15. Stein DM, Vawdrey DK, Stetson PD, Bakken S: An analysis of team checklists in physician signout notes. *AMIA Annu Symp Proc*. 2010, 2010:767-71.
16. Starmer AJ, Spector ND, Srivastava R, Allen AD, Landrigan CP, Sectish TC, the I-PASS Study Group: I-pass, a mnemonic to standardize verbal handoffs. *Pediatrics*. 2012, 129:201-4. [10.1542/peds.2011-2966](#)
17. Starmer AJ, Sectish TC, Simon DW, et al.: Rates of medical errors and preventable adverse events among hospitalized children following implementation of a resident handoff bundle. *JAMA*. 2013, 310:2262-70. [10.1001/jama.2013.281961](#)
18. Starmer AJ, Spector ND, Srivastava R, et al.: Changes in medical errors after implementation of a handoff program. *N Engl J Med*. 2014, 371:1803-12. [10.1056/NEJMsa1405556](#)
19. Graham KL, Marcantonio ER, Huang GC, Yang J, Davis RB, Smith CC: Effect of a systems intervention on the quality and safety of patient handoffs in an internal medicine residency program. *J Gen Intern Med*. 2013, 28:986-93. [10.1007/s11606-013-2391-7](#)
20. Patel VP, Raptis D, Christofi T, et al.: Development of electronic software for the management of trauma patients on the orthopaedic unit. *Injury*. 2009, 40:388-96. [10.1016/j.injury.2008.10.006](#)
21. Rao BS, Lowe GO, Hughes AJ: Reduced emergency calls and improved weekend discharge after introduction of a new electronic handover system. *Med J Aust*. 2012, 197:569-73. [10.5694/mja11.11048](#)
22. Wohlauer MV, Rove KO, Pshak TJ, et al.: The computerized rounding report: implementation of a model system to support transitions of care. *J Surg Res*. 2012, 172:11-7. [10.1016/j.jss.2011.04.015](#)
23. Davis J, Riesenberg LA, Mardis M, Donnelly J, Benningfield B, Youngstrom M, Vetter I: Evaluating outcomes of electronic tools supporting physician shift-to-shift handoffs: a systematic review. *J Grad Med Educ*. 2015, 7:174-80. [10.4300/JGME-D-14-00205.1](#)