Amyand Hernia With Acute Appendicitis: A Rare Type of Hernia

Review began 10/09/2021
Review ended 10/11/2021
Published 10/12/2021
Exp. concern 04/07/2022

© Copyright 2021

Assad et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Majd A. Assad¹, Thamer A. Boushal², Zeyad W. Halawani³, Rayan S. Alruwaili³, Yakeen H. Alkuwaiti¹, Aisha M. Alghamdi⁴, Wedad A. Bahri⁵, Bader M. Albilasi⁶, Shrouq M. Alghamdi⁷, Bushra M. Alghanmi⁷, Manal M. Alghanmi⁷, Abdulaziz J. Alghamdi⁸, Moath M. Alkhalidi⁹, Malak Alshammari¹⁰

1. Surgery, College of Medicine, Imam Abdulrahman Bin Faisal University, Dammam, SAU 2. Surgery, College of Medicine, King Faisal University, Al-Ahsa, SAU 3. Surgery, College of Medicine & Medical Science, Arabian Gulf University, Manama, BHR 4. Surgery, College of Medicine, Al-Baha University, Al-Baha, SAU 5. Surgery, College of Medicine, Batterjee Medical College, Jeddah, SAU 6. Surgery, College of Medicine, Jof University, Al-Jawf, SAU 7. Surgery, College of Medicine, Umm Al-Qura University, Mecca, SAU 8. Surgery, College of Medicine, Ibn Sina National College for Medical Studies, Jeddah, SAU 9. Surgery, College of Medicine, Jazan University, Jizan, SAU 10. Internal Medicine, College of Medicine, Imam Abdulrahman Bin Faisal University, Dammam, SAU

Corresponding author: Malak Alshammari, saudidoctor2020@gmail.com

Expression of Concern

Expression of Concern date: April 07, 2022. Cite this expression of concern as Assad M A, Boushal T A, Halawani Z W, et al. (April 07, 2022) Expression of Concern: Amyand Hernia With Acute Appendicitis: A Rare Type of Hernia. Cureus 14(4): x8. doi:10.7759/cureus.x8.

The concern relates to the provenance of this article as brought to our attention by Faisal Alhawaj, who denies authorship of this article and others published in Cureus. These articles were submitted and subsequently published purportedly as an effort coordinated by Imam Abdulrahman Bin Faisal University to ensure all medical interns publish at least one peer-reviewed article in order to qualify for enrollment in a postgraduate residency program as stipulated by The Saudi Commission for Health Specialties (SCFHS).

The journal has not been presented with enough evidence to warrant the formal retraction of these articles as both Imam Abdulrahman Bin Faisal University and The Saudi Commission for Health Specialties have failed to respond to numerous communications requesting additional information regarding these allegations. While we acknowledge that the provenance of these articles is very much in question, we cannot act until these claims have been investigated by the appropriate institutions with the results of said investigation communicated to Cureus.

The concern and this note will remain appended to the above-mentioned article until Cureus is provided with official confirmation from Imam Abdulrahman Bin Faisal University or The Saudi Commission for Health Specialties.

Abstract

A hernia is an abnormal protrusion of an organ or tissue from its containing cavity. Inguinal hernia is the most common type of abdominal hernias. The presence of the appendix within the inguinal hernial sac is a rare occurrence. We present the case of a 41-year-old man, with no previous surgical history, who presented to the emergency department with a right-sided painful groin swelling. The swelling was associated with nausea, vomiting, anorexia, and fever. Initial laboratory investigation revealed leukocytosis and elevated C-reactive protein and erythrocyte sedimentation rate. A computed tomography (CT) scan demonstrated the presence of an inflamed appendix with an inguinal hernia. The patient underwent laparoscopic repair of the hernia with appendectomy. The patient tolerated the procedure without any complications. Amyand hernia with acute appendicitis is a very rare surgical condition. Clinicians should have a high index of suspicion for Amyand hernia with appendicitis when they encounter a patient with groin mass with laboratory markers indicating an inflammatory process.

Categories: Emergency Medicine, Radiology, General Surgery

Keywords: case report, operative laparoscopy, amyand's hernia, acute appendicitis, inguinal hernia

Introduction

A hernia is an abnormal protrusion of an organ or tissue from its containing cavity. Inguinal hernia repair remains one of the most commonly performed surgeries worldwide. It accounts for 80% of the abdominal wall hernias. The lifetime prevalence of inguinal hernia reaches over 25% [1]. Surgical repair is the only cure for such hernias. The hernia is termed "incarcerated" if the contained organ is non-reducible. Besides, strangulated hernia refers to a hernia with a compromised blood supply, leading to ischemia and necrosis. The hernial sac often contains omental fat or bowel [2]. A significant array of surgical approaches has been

How to cite this article

Assad M A, Boushal T A, Halawani Z W, et al. (October 12, 2021) Amyand Hernia With Acute Appendicitis: A Rare Type of Hernia. Cureus 13(10): e18720. DOI 10.7759/cureus.18720

described for the management of inguinal hernia. Here, we report the case of a middle-aged man with an Amyand hernia, an inguinal hernia containing the appendix, who was managed successfully by laparoscopic repair. Laparoscopic surgery has diagnostic and therapeutic roles in Amyand hernia. A limited number of cases have been reported where the laparoscopic approach was used for Amyand hernia. The first case of laparoscopic repair of Amyand hernia was described by Vermillion et al. in 1999 [3].

Case Presentation

We present the case of a 41-year-old man who presented to the emergency department complaining of swelling in his right groin for two weeks’ duration. He reported that the swelling became painful and non-reducible for the last two days. The pain was associated with decreased appetite and subjective fever. The patient reported having nausea and recurrent episodes of vomiting in the last 24 hours. The patient did not undergo any previous surgeries. His past medical history was remarkable for well-controlled asthma. He was a non-smoker and never drinks alcohol. The patient worked as a school teacher. He has a significant family history of sickle cell disease.

Upon examination, the patient appeared in distress due to the pain. His vital signs were as follows: a temperature of 38°C, a pulse rate of 110 beats per minute, blood pressure of 118/95 mmHg, and respiratory rate of 14 breaths per minute. There was a swelling in the right inguinal region, measuring 4 x 6 cm. It was oval in shape and had a smooth surface with an evident expansile cough impulse. The swelling was non-reducible with slightly erythematous and warm overlying skin.

In view of the clinical features, the diagnosis of incarcerated inguinal hernia was established. The patient underwent a pre-operative laboratory investigation, which revealed leukocytosis with neutrophilia and left shift, elevated erythrocyte sedimentation rate and C-reactive protein, and normal renal and hepatic profiles (Table 1). The patient underwent an abdominal computed tomography (CT) scan to identify the content of the hernia and diagnose any possible complications. CT demonstrated a right-sided inguinal hernia that contained the cecum and a rim-enhancing structure that represented an inflamed appendix with surrounding inflammatory changes (Figure 1). Such radiological findings conferred the diagnosis of Amyand hernia with acute appendicitis.

Laboratory Investigation	Unit	Result	Reference Range
Hemoglobin	g/dL	14.2	13.0–18.0
White blood cell	1,000/mL	13.4	4.0–11.0
Platelet	1,000/mL	370	140–450
Erythrocyte sedimentation rate	mm/hr.	25	0–20
C-reactive protein	mg/dL	14.5	0.3–10.0
Total bilirubin	mg/dL	0.8	0.2–1.2
Albumin	g/dL	4.1	3.4–5.0
Alkaline phosphatase	U/L	51	46–116
Gamma-glutamyltransferase	U/L	46	15–85
Alanine transferase	U/L	60	14–63
Aspartate transferase	U/L	35	15–37
Blood urea nitrogen	mg/dL	16	7–18
Creatinine	mg/dL	0.8	0.7–1.3
Sodium	mEq/L	138	136–145
Potassium	mEq/L	3.8	3.5–5.1
Chloride	mEq/L	104	98–107

TABLE 1: Summary of the results of laboratory findings

FIGURE 1: CT image

Axial image of CT scan of the abdomen demonstrating a right inguinal hernia with a rim-enhancing structure representing the inflamed appendix (arrow).

CT, computed tomography

The patient was prepared for an emergency laparoscopy operation for the reduction of the inguinal hernia. The patient was administered general anesthesia and was placed in a supine position. Pneumoperitoneum was established. The diagnostic evaluation revealed a right inguinal hernia containing an inflamed appendix. An appendectomy was performed, and the appendix was removed in a sterile bag. The hernia was repaired using the totally extraperitoneal approach with mesh repair. No complications were encountered during the operation. The estimated blood loss was 5 mL and the total operative time was 120 minutes. The patient had an uneventful recovery and was discharged on the fifth postoperative day. The histopathological examination of the appendix confirmed the diagnosis of acute appendicitis.

Discussion

We described the case of a middle-aged man with an Amyand hernia and acute appendicitis that was managed successfully via a laparoscopic approach. The occurrence of the appendix within the sac of inguinal hernia is very rare. It was first described by Claudius Amyand in 1736 who successfully performed appendectomy in a young child with a right inguinal hernia [4]. Amyand hernia represents less than 1% of inguinal hernias [2]. Furthermore, it is very rare for patients with Amyand hernia to present with concomitant Amyand hernia and acute appendicitis. In the present case, the patient was found to have neutrophilic leukocytosis indicating an inflammatory process. However, such findings are non-specific.

In the present case, the appendix was found to be inflamed as per the CT findings. The presence of appendicitis within the hernia sac has essential surgical implications. For instance, the surgeon needs to perform an appendectomy instead of a simple hernia repair in the case of Amyand hernia with acute appendicitis. It should be noted that the Amyand hernia may contain a normal or an inflamed appendix. However, the herniated appendix is at a higher risk of developing an infection. It is postulated that the narrow hernial neck causes restricted blood flow, leading to edema and inflammation [2].

In contrast to the present case, Amyand hernia is three times more prevalent among children than adults. However, it may occur at any age and has been reported in patients ranging from neonates to the elderly [5]. It is much more common among male patients. If Amyand hernia develops among female patients, it tends to be found among postmenopausal women [6].

Early diagnosis and management of Amyand hernia are crucial. The reported mortality rate from Amyand hernia may reach up to 30% [7]. However, the mortality is substantially reduced as earlier Amyand hernia was often diagnosed incidentally during surgery. The widespread use of advanced imaging allowed the preoperative diagnosis of Amyand hernia with acute appendicitis. Since inguinal hernia is a clinical diagnosis, patients often undergo surgical repair without the need for prior radiological imaging. In the present case, however, the patient was found to have abnormal laboratory markers suggestive of an

inflammatory process, which mandated further investigations. We performed a CT scan to rule out any intra-abdominal infection. CT remains the imaging modality of choice for patients with groin mass. However, there are several cases in the literature where the Amyand hernia was diagnosed via ultrasound examination [8].

There are controversies about the optimal surgical approach [9]. Several factors should be taken into consideration regarding the use of mesh for the repair of the hernia, including the patient's age, comorbidities, life expectancy, and presence of complicated appendicitis. Many authors did not recommend performing appendectomy in Amyand hernia if the appendix was normal or mildly congested since it may increase the risk of septic complications [10]. In our case, the appendix was grossly inflamed with a significant risk of perforation. We used a tension-free mesh repair as there was no evidence of peritoneal sepsis.

Conclusions

Amyand hernia with acute appendicitis is a very rare surgical condition. Clinicians should have a high index of suspicion for Amyand hernia with appendicitis when they encounter a patient with groin mass with laboratory markers indicating an inflammatory process. A CT scan is essential in making the preoperative diagnosis. The laparoscopic approach is feasible and safe for the repair of the Amyand hernia with acute appendicitis.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. University Institutional Review issued approval N/A. Case reports are waived by the Institutional Review Board at our institution.

Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. The HerniaSurge Group: International guidelines for groin hernia management . *Hernia*. 2018, 22:1-165. [10.1007/s10029-017-1668-x](https://doi.org/10.1007/s10029-017-1668-x)
2. Ash L, Hatem S, Ramirez GA, Veniero J: Amyand's hernia: a case report of prospective ct diagnosis in the emergency department. *Emerg Radiol*. 2005, 11:231-2. [10.1007/s10140-005-0411-6](https://doi.org/10.1007/s10140-005-0411-6)
3. Vermillion J, Abernathy S, Snyder S: Laparoscopic reduction of Amyand's hernia . *Hernia*. 1999, 3:159-60. [10.1007/BF01195318](https://doi.org/10.1007/BF01195318)
4. Ryan WJ: Hernia of the vermiform appendix . *Ann Surg*. 1937, 106:135-8.
5. Anagnostopoulou S, Dimitroulis D, Troupis TG, et al.: Amyand's hernia: a case report . *World J Gastroenterol*. 2006, 12:4761-3. [10.3748/wjg.v12.i29.4761](https://doi.org/10.3748/wjg.v12.i29.4761)
6. Sharma H, Gupta A, Shekhawat NS, Memon B, Memon MA: Amyand's hernia: a report of 18 consecutive patients over a 15-year period. *Hernia*. 2007, 11:31-5. [10.1007/s10029-006-0153-8](https://doi.org/10.1007/s10029-006-0153-8)
7. D'Alia C, Lo Schiavo MG, Tonante A, et al.: Amyand's hernia: case report and review of the literature . *Hernia*. 2003, 7:89-91. [10.1007/s10029-002-0098-5](https://doi.org/10.1007/s10029-002-0098-5)
8. Robinson MB: Amyand hernia diagnosed with sonography . *J Diagn Med Sonogr*. 2011, 27:279-82. [10.1177/8756479311424433](https://doi.org/10.1177/8756479311424433)
9. Shaban Y, Elkbuli A, McKenney M, Boneva D: Amyand's hernia: a case report and review of the literature . *Int J Surg Case Rep*. 2018, 47:92-6. [10.1016/j.ijscr.2018.04.034](https://doi.org/10.1016/j.ijscr.2018.04.034)
10. Hutchinson R: Amyand's hernia. *J R Soc Med*. 1993, 86:104-5.