Ocular Coloboma With Choroidal Neovascular Membrane: A Case Report

Review began 10/27/2021

Review ended 11/07/2021

Published 11/13/2021

© Copyright 2021

Almasaud et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Jluwi Almasaud¹, Sara A. Aledaili², Reem S. Alshammari³

1. Ophthalmology, King Khaled Hospital, Hail, SAU 2. Medicine, University of Hail College of Medicine, Hail, SAU 3. Ophthalmology, University of Hail College of Medicine, Hail, SAU

Corresponding author: Sara A. Aledaili, saraaledaili@gmail.com

Abstract

Ocular coloboma (OC) is a rare congenital anomaly and a product of a defect in embryogenesis. It is the result of fetal fissure closure error that ends with a persistent cleft. Colobomas are generally accompanied by visual loss. In this article, we present a case of bilateral iris, disc, and retina coloboma that was managed with an anti-vascular endothelial growth factor (ranibizumab), and as a result, caused regression of the choroidal neovascular membranes and improved the patient's visual acuity. However, The patient will need lifelong follow-up to catch any retinochoroidal changes or development of cataract or glaucoma.

Categories: Ophthalmology

Keywords: congenital, choroidal neovascular membrane, ophthalmology, iris, ocular coloboma

Introduction

Ocular coloboma (OC) is a congenital anomaly represented by missing pieces of tissue in the eye structure that appear as notches or gaps in the iris, retina, choroid, or optic nerve. Eye development starts on the 20th day of embryo life when a pair of shallow grooves appear on the sides of the invaginating forebrain [1]. Ocular coloboma is the final product of an error occurring in the closure of the fetal fissure. Hence, it ends with a persistent cleft; the normal process takes place from the fifth to sixth weeks of pregnancy [2]. Generally, colobomas are associated with visual loss, which may need management and intervention besides amblyopia management [3]. Therefore, recognition of congenital anomalies of the eye can improve parents' understanding as well as genetic counseling. In this article, we present a case of bilateral iris, disc, and retina coloboma. The article was presented as a poster at the 2nd Global Congress on Medical & Clinical Case Reports in Dubai, United Arab Emirates, on November 19, 2018.

Case Presentation

A 20-year-old male was referred from a remote village to an Ophthalmology clinic with a painless as well as progressive decrease of vision in the right eye for the past two months. The decrease of vision in the right eye was gradual, with no pain nor history of ocular trauma or surgeries. His left eye was amblyopic. The patient had no medical history or any significant family history. On ocular examination, his unassisted visual acuity was 20/200 in his right eye and hand motion in his left eye. The intraocular pressure was 18 mmHg bilaterally. On inspection, there was exotropia of the left eye. Next, the slit lamp evaluation of the anterior segment was within normal limits except for iris coloboma in the inferior nasal quadrant of each eye (Figure 1).

FIGURE 1: (A) Right eye iris coloboma. (B) Left eye iris coloboma.

Dilated fundus examination showed optic disc and retinochoroidal coloboma in both eyes (Figure 2) with

How to cite this article

Almasaud J, Aledaili S A, Alshammari R S (November 13, 2021) Ocular Coloboma With Choroidal Neovascular Membrane: A Case Report. Cureus 13(11): e19521. DOI 10.7759/cureus.19521

juxtapapillary choroidal neovascularization in the right eye. Furthermore, we performed optical coherence tomography (OCT), which confirmed the presence of choroidal neovascular membrane (CNVM) (Figure 3).

FIGURE 2: (A) Right eye optic disc inferonasal retina coloboma. (B) Left eye optic disc retina coloboma.

FIGURE 3: Right eye OCT.

OCT: optical coherence tomography

Discussion

As it is a rare congenital anomaly, ocular coloboma (OC) has a prevalence that is around 0.14% of the general population [4]. It is initiated by imperfect embryogenesis, which leads to failed closure of the embryonic fissure during the first trimester of fetal life [5]. Worldwide, the incidence of the condition is seven for every 10,000 live births [1]. In this report, we presented a patient with bilateral iris, optic disc, retinochoroidal coloboma. Unfortunately, in Saudi Arabia, no records exist of the incidence of OC. The occurrence of sporadic, unilateral, or bilateral coloboma varies with genetic and non-genetic factors; it also could be related to some syndromes such as coloboma, heart anomaly, choanal atresia, retardation, genital and ear anomalies (CHARGE) syndrome [6]. Studies suggest that the usage of some drugs during pregnancy could be linked with OC [7,8].

Here we described a case of OC associated with optic disc, retina, choroid, and iris coloboma that we consider unlikely to be inherited because neither one of the parents or siblings of the patients had the same condition. And it is less often that an autosomal dominant inheritance may occur [9]. Chromosomal analysis is needed for a patient with OC, but it was not possible in our case due to non-availability at our facilities.

Our patient presented with bilateral iris, optic disc, and retinochoroidal coloboma, which is consistent with previous studies that suggest that 50% percent of patients with OC have bilateral disease [10]. He also had right choroidal neovascularization in the inferior nasal area that caused the decrease of vision. In such cases,

the treatment plan will include either anti-vascular endothelial growth factor (anti-VEGF), focal laser photocoagulation, or photodynamic therapy. According to the literature, it is usually preferred to start with focal laser photocoagulation [11], but regarding our patient who had the CNVM in juxtapapillary area, we decided to treat him with anti-VEGF to avoid the potential risk of iatrogenic vision loss. The treatment with anti-VEGF has resulted in regression of the CNVM, hence improving the patient's visual acuity. Furthermore, we plan to continue the patient on more anti-VEGF until reaching complete regression of the CNVM and routine follow-up in the future.

Conclusions

To the best of our knowledge, the presented case is the first case of coloboma reported in the northern region of Saudi Arabia. We recommend early recognition of such cases, which will improve parents' understanding as well as genetic counseling. Furthermore, we also recommend lifelong follow-up for coloboma cases to catch any retinochoroidal changes or development of cataract and glaucoma. Correspondingly, more studies are needed to better understand ocular coloboma's prevalence, causes, and management.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Sadler TW, Langman J: Langman's Medical Embryology. 12th edition / Sadler TW (ed): Wolters Kluwer Health/Lippincott Williams & Wilkins, Philadelphia; 2012.
2. Onwochei BC, Simon JW, Bateman JB, Couture KC, Mir E: Ocular colobomata. *Surv Ophthalmol.* 2000, 45:175-94. [10.1016/s0039-6257\(00\)00151-x](https://doi.org/10.1016/s0039-6257(00)00151-x)
3. Levin AV: Congenital eye anomalies. *Pediatr Clin N Am.* 2003, 50:55-76. [10.1016/S0031-3955\(02\)00113-X](https://doi.org/10.1016/S0031-3955(02)00113-X)
4. Amador-Patarroyo MJ, Pérez-Rueda MA, Tellez CH: Congenital anomalies of the optic nerve. *Saudi J Ophthalmol.* 2015, 29:32-8. [10.1016/j.sjopt.2014.09.011](https://doi.org/10.1016/j.sjopt.2014.09.011)
5. Chang L, Blain D, Bertuzzi S, Brooks BP: Uveal coloboma: clinical and basic science update. *Curr Opin Ophthalmol.* 2006, 17:447-70. [10.1097/01.icu.0000243020.82380.f6](https://doi.org/10.1097/01.icu.0000243020.82380.f6)
6. Russell-Eggitt IM, Blake KD, Taylor DS, Wyse RK: The eye in the CHARGE association. *Br J Ophthalmol.* 1990, 74:421-6. [10.1136/bjo.74.7.421](https://doi.org/10.1136/bjo.74.7.421)
7. Miller MT, Strömland K: Teratogen update: thalidomide: a review, with a focus on ocular findings and new potential uses. *Teratology.* 1999, 60:306-21. [10.1002/\(SICI\)1096-9926\(199911\)60:5<306::AID-TERA11>3.0.CO;2-Y](https://doi.org/10.1002/(SICI)1096-9926(199911)60:5<306::AID-TERA11>3.0.CO;2-Y)
8. Strömland K, Pinazo-Durán MD: Ophthalmic involvement in the fetal alcohol syndrome: clinical and animal model studies. *Alcohol Alcohol.* 2002, 37:2-8. [10.1093/alcalc/37.1.2](https://doi.org/10.1093/alcalc/37.1.2)
9. Savell J, Cook JR: Optic nerve colobomas of autosomal-dominant heredity. *Arch Ophthalmol.* 1976, 94:395-400. [10.1001/archophth.1976.03910030183002](https://doi.org/10.1001/archophth.1976.03910030183002)
10. Biedner B, Klemperer I, Dagan M, Yassur Y: Optic disc coloboma associated with macular hole and retinal detachment. *Ann Ophthalmol.* 1993, 25:350-2.
11. Hussain RM, Abbey AM, Shah AR, Drenser KA, Trese MT, Capone A Jr: Chorioretinal coloboma complications: retinal detachment and choroidal neovascular membrane. *J Ophthalmic Vis Res.* 2017, 12:3-10. [10.4103/2008-322X.200163](https://doi.org/10.4103/2008-322X.200163)