Posterior Percutaneous Screw Fixation to Treat Vertebral Fracture Non-union in Diffuse Idiopathic Skeletal Hyperostosis

Review began 11/16/2021
Review ended 11/22/2021
Published 11/25/2021

© Copyright 2021

Wensley et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Katherine E. Wensley¹, Daniel Rolton²

1. Plastic and Reconstructive Surgery, Aberdeen Royal Infirmary, Aberdeen, GBR 2. Trauma and Orthopaedics, Royal Berkshire National Health Service Foundation Trust, Reading, GBR

Corresponding author: Katherine E. Wensley, kate.wensley@nhs.scot

Abstract

Diffuse idiopathic skeletal hyperostosis (DISH) alters the biomechanical properties of the spine, rendering it highly prone to fracture, following even minor trauma. Risk of delayed diagnosis of vertebral fractures is particularly high in this cohort of patients since radiographs are notoriously difficult to interpret and presentation is late, due to difficulty distinguishing new from pre-existing back pain. Our case describes a gentleman in his late sixties with a six-month delay in presentation to our services with a T12 fracture, secondary to previously undiagnosed DISH, which had progressed to non-union. He underwent T9-L5 thoracolumbar posterior percutaneous stabilisation and fusion. At eighteen months follow-up, there was evidence of union, significant improvement in his pain, no focal neurology signs, and the patient had returned to his activities of daily living (ADLs).

Categories: Radiology, Orthopedics, Trauma

Keywords: spine imaging, minimally invasive and robotic spine surgery, vertebra fracture, ankylosing spinal disorder, diffuse idiopathic skeletal hyperostosis

Introduction

Diffuse idiopathic skeletal hyperostosis (DISH) is an increasingly common non-inflammatory ankylosing spinal condition, often defined by the Resnick and Niwayama criteria, the hallmarks of which are: the bridging of four adjacent vertebral bodies by newly formed bone, without severe loss of the intervertebral disc height and an absence of degeneration of the apophyseal and sacroiliac joints [1].

These features result in progressive spinal fusion with reduced motion and kyphotic spinal column alignment. This alteration of spinal biomechanical properties results in a significantly increased fracture risk, in trauma considered otherwise minor [2]. Whilst fracture risk in DISH has not been quantified, in patients with ankylosing spondylitis—where fractures have been attributed to the same mechanism—risk [3,4] is increased fourfold [5]. In addition to this increased fracture risk, complications with healing are more prevalent and non-union rates higher. If vertebral fractures are unstable or progress to non-union, patients can be left with considerable pain and are at high risk of developing progressive neurological signs [6].

Incidence of delayed diagnosis of fractures in DISH is persistently high [7]. Multiple patient and doctor factors for these delays have been cited. These include late patient presentation due to difficulty distinguishing new from pre-existing back pain and difficulty interpreting radiographs due to fused joints and osteoporotic bone quality [8]. Clinician awareness of the increased fracture risk in DISH is therefore essential to facilitating CT or MRI screen scanning in this patient cohort. This allows timely diagnosis and appropriate management.

Case Presentation

We present the case of a gentleman in his late sixties who was referred to our spinal clinic by his general practitioner (GP). He first attended the clinic six months after a fall from standing height, complaining of ongoing back pain. Plain radiographs arranged by his GP three weeks after his fall revealed a T12 fracture, and over the next several months, his pain failed to settle. He had no radicular symptoms and no change in bowel or bladder habits. His medical history included a radical oesophagectomy for oesophageal carcinoma eight years previously, chronic obstructive pulmonary disease (COPD), asthma, and morbid obesity. He lived alone in a bungalow, normally mobilised without any walking aids and was independent with his activities of daily living (ADLs). Following the fracture, he occasionally used a stick to mobilise and reported that his pain was preventing him from carrying out his ADLs.

Plain radiographs were arranged by his GP three weeks following his fall and revealed a minor wedge fracture of the T12 body with uncertain chronicity. Repeat imaging three months later suggested worsening of the fracture. Upon receipt of referral to our services, six months following his original injury, and prior to clinic attendance, he underwent an MRI scan on which there was evidence of fracture non-union (Figure 1).

How to cite this article

Wensley K E, Rolton D (November 25, 2021) Posterior Percutaneous Screw Fixation to Treat Vertebral Fracture Non-union in Diffuse Idiopathic Skeletal Hyperostosis. Cureus 13(11): e19895. DOI 10.7759/cureus.19895

FIGURE 1: Sagittal T2 MRI image demonstrating T12 fracture non-union (red arrow), note that the image also demonstrates evidence of endplate changes.

Following assessment in the clinic, standing thoracolumbar radiographs and CT scan were arranged (Figure 2). The standing thoracolumbar radiographs were felt to be sub-optimal, but within the limitations of the study, the T12 fracture appeared to have worsened from initial radiographs, and with multi-level degenerative disc disease in the form of prominent anterior osteophyte formation was noted. On the basis of this imaging, it was felt that instrumentation was the best option to achieve adequate analgesia through spinal stabilisation and fracture union.

FIGURE 2: CT image demonstrating the horizontal T12 fracture to be irregular (blue arrow) with the reduction in vertebral body height, and no evidence of bony union. In keeping with the diagnosis of DISH, there is annulus fibrosus and central discal calcification (yellow arrow), associated with smooth flowing right anterolateral vertebral body osteophytosis.

DISH: diffuse idiopathic skeletal hyperostosis.

The procedure was performed under general anaesthetic by consultant spinal surgeon Daniel Rolton, with great care taken over patient positioning due to the risk of iatrogenic injury. Surgical stabilisation was achieved using a minimally invasive technique, through the use of posterior percutaneous pedicle screws, spanning three levels on either side of the fracture (from T9 to L3) and 5.5 mm titanium rods bilaterally. Pedicle screws were 6 mm × 40 mm for thoracic vertebrae and 6 mm × 45 mm for lumbar vertebrae (Viper, DePuy-Synthes). The operation was tolerated well and check imaging revealed satisfactory metalwork placement and satisfactory fracture reduction (Figures [3A](#), [3B](#)).

FIGURE 3: Post-operative radiographs (A) anterior-posterior view, (B) lateral view demonstrating good fracture reduction and satisfactory metalwork placement.

Our patient remained in hospital for two days post-operatively, with no complications. He was followed up in our clinic at four-month intervals from six weeks post-operatively to eighteen months post-operatively, at which point he was discharged back to the care of his GP. His surgical wounds healed well, with no metalwork complications, and there was radiological evidence of union. At four months post-operatively, he required no mobilisation aids and returned to work. When seen twelve months following his operation, he described himself as "85% better" compared to his pre-operative status, with no pain routinely, having discontinued his analgesia. However, he reported some backache when standing and walking for long periods.

Discussion

DISH is a common but poorly understood condition. The aetiology is unknown but close associations with advancing age, diabetes, obesity, and a male sex have been demonstrated. The reported incidence varies widely based on age and population studied, from 2.9% in the Asian population [9] to 25% in the Caucasian population [10]. Given increasing life expectancies and increasing rates of type 2 diabetes mellitus and obesity, the incidence of DISH is likely to continue to rise in the coming years.

Non-operative management was previously favoured for vertebral fractures in patients with ankylosing spinal disease (ASD) but outcomes were generally poor [11,12]. Open pedicle screw fixation has also been widely practiced but more recently, both Yeoh et al. and Bredin et al. have reported good outcomes and lower complication rates with percutaneous pedicle screw fixation in patients with ASD [8,13]. However, to our knowledge, no series exist on the management of missed fractures that have progressed to non-union in patients with DISH.

As in this case, DISH patients are often heavily co-morbid, making minimally invasive surgery (MIS) a particularly attractive option [6,14,15]. A meta-analysis by McAnany et al. demonstrated equivalence in clinical outcomes between open and percutaneous procedures [16]. Additionally, recovery time, pain, and time until return to work have all been demonstrated to be reduced, when comparing percutaneous to open spinal procedures [17,18].

There is no consensus as to the optimal number of levels to fix in patients with ASD. The progressive fusion of the spinal column in DISH produces fractures similar to those seen in long bones, with a torque associated with the lever arms of the spine [14]. This, in combination with the reduced bone mineral density in DISH

patients, necessitates multiple point fixation to stabilise the fracture. The decision to instrument three levels above and below the fracture was based on work by Werner et al. who suggest this as the minimum for effective fixation [19].

Conclusions

With the ageing and increasingly co-morbid population, the prevalence of DISH is rising and should feature in the differential for patients presenting with back pain following minor trauma. Where fractures secondary to DISH are suspected, CT or MRI scanning should be considered as plain radiographs frequently fail to identify DISH. If the diagnosis is delayed and fractures have progressed to non-union, minimally invasive thoracolumbar posterior percutaneous stabilisation offers effective means of providing analgesia through the promotion of union.

Our case demonstrates the successful use of thoracolumbar posterior percutaneous stabilisation to manage a thoracic vertebral fracture, secondary to DISH, which had progressed to non-union. However, increased clinical awareness of DISH is vital to improve recognition rates of affected patients at initial presentation and as such, minimise the number of patients with a delayed diagnosis and fracture non-union.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Resnick D, Niwayama G: Radiographic and pathologic features of spinal involvement in diffuse idiopathic skeletal hyperostosis (DISH). *Radiology*. 1976, 119:559-68. [10.1148/119.3.559](#)
2. Hunter T, Forster B, Dvorak M: Ankylosed spines are prone to fracture. *Can Fam Physician*. 1995, 41:1213-6.
3. Finkelstein JA, Chapman JR, Mirza S: Occult vertebral fractures in ankylosing spondylitis. *Spinal Cord*. 1999, 37:444-7. [10.1038/sj.sc.3100837](#)
4. Paley D, Schwartz M, Cooper P, et al.: Fractures of the spine in diffuse idiopathic skeletal hyperostosis. *Clin Orthop*. 1991, 267:22-32.
5. Young JS, Cheshire DJE, Pierce JA, Vivian JM: Cervical ankylosis with acute spinal cord injury. *Spinal Cord*. 1977, 15:133-46. [10.1038/sc.1977.19](#)
6. Westerveld LA, Verlaan JJ, Oner FC: Spinal fractures in patients with ankylosing spinal disorders: a systematic review of the literature on treatment, neurological status and complications. *Eur Spine J*. 2009, 18:145-56. [10.1007/s00586-008-0764-0](#)
7. Rustagi T, Drazin D, Oner C, et al.: Fractures in spinal ankylosing disorders: a narrative review of disease and injury types, treatment techniques, and outcomes. *J Orthop Trauma*. 2017, 31:57-74. [10.1097/BOT.0000000000000953](#)
8. Yeoh D, Moffatt T, Karmani S: Good outcomes of percutaneous fixation of spinal fractures in ankylosing spinal disorders. *Injury*. 2014, 45:1534-8. [10.1016/j.injury.2014.03.020](#)
9. Kim SK, Choi BR, Kim CG, et al.: The prevalence of diffuse idiopathic skeletal hyperostosis in Korea. *J Rheumatol*. 2004, 31:2032-5.
10. Weinfeld RM, Olson PN, Maki DD, Griffiths HJ: The prevalence of diffuse idiopathic skeletal hyperostosis (DISH) in two large American Midwest metropolitan hospital populations. *Skeletal Radiol*. 1997, 26:222-5. [10.1007/s002560050225](#)
11. Westerveld LA, van Bommel JC, Dhert WJ, Oner FC, Verlaan JJ: Clinical outcome after traumatic spinal fractures in patients with ankylosing spinal disorders compared with control patients. *Spine J*. 2014, 24:729-40. [10.1016/j.spinee.2013.06.038](#)
12. Chaudhary SB, Hullinger H, Vives MJ: Management of acute spinal fractures in ankylosing spondylitis. *ISRN Rheumatol*. 2011, 2011:150484. [10.5402/2011/150484](#)
13. Bredin S, Fabre-Aubrespy M, Blondel B, et al.: Percutaneous surgery for thoraco-lumbar fractures in ankylosing spondylitis: study of 31 patients. *Orthop Traumatol Surg Res*. 2017, 103:1235-9. [10.1016/j.otsr.2017.07.023](#)
14. Caron T, Bransford R, Nguyen Q, Agel J, Chapman J, Bellabarba C: Spine fractures in patients with ankylosing spinal disorders. *Spine*. 2010, 35:458-64. [10.1097/BRS.0b013e3181cc764f](#)
15. Whang PG, Goldberg G, Lawrence JP, et al.: The management of spinal injuries in patients with ankylosing spondylitis or diffuse idiopathic skeletal hyperostosis: a comparison of treatment methods and clinical outcomes. *Clin Spine Surg*. 2009, 22:77-85. [10.1097/BSD.0b013e3181679bcb](#)
16. McAnany SJ, Overley SC, Kim JS, Baird EO, Qureshi SA, Anderson PA: Open versus minimally invasive fixation techniques for thoracolumbar trauma: a meta-analysis. *Global Spine J*. 2016, 26:186-94. [10.1055/s-0035-1554777](#)
17. Hsieh PC, Koski TR, Sciubba DM, et al.: Maximizing the potential of minimally invasive spine surgery in complex spinal disorders. *Neurosurg Focus*. 2008, 25:E19. [10.3171/FOC/2008/25/8/E19](#)

18. Foley KT, Holly LT, Schwender JD: Minimally invasive lumbar fusion . Spine. 2003, 28:26-35. [10.1097/01.BRS.0000076895.52418.5E](https://doi.org/10.1097/01.BRS.0000076895.52418.5E)
19. Werner BC, Samartzis D, Shen FH: Spinal fractures in patients with ankylosing spondylitis: etiology, diagnosis, and management. J Am Acad Orthop Surg. 2016, 24:241-9. [10.5435/JAOS-D-14-00149](https://doi.org/10.5435/JAOS-D-14-00149)