A Clinico-Epidemiological Study of Neurofibromatosis Type 1 and Its Relation to Quality of Life: A Cross-Sectional Study From India

Review began 02/07/2022

Review ended 02/17/2022

Published 02/19/2022

© Copyright 2022

Mahajan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Aditi Mahajan¹, Milind Patvekar¹, Samruddhi Lote¹, Mahendra Singh Deora¹, Divya Poulse¹, Jaya Madhurya Gogineni¹, Krish Panikar², Biswajit Chaklader³

1. Dermatology, Dr. D. Y. Patil Medical College, Hospital and Research Centre, Pune, IND 2. Internal Medicine, A.J. Institute of Medical Sciences, Mangalore, IND 3. Preventive Medicine, Dr. D. Y. Patil Medical College, Hospital and Research Centre, Pune, IND

Corresponding author: Aditi Mahajan, aditiginger@gmail.com

Abstract

Introduction: Neurofibromatosis type 1 (NF1) is an inherited neuroectodermal abnormality with multisystem effects, which can have heavy psychological and physical burdens, especially in countries like India, wherein skin disease is significantly stigmatized. This study was performed to understand the clinical and epidemiological trends of NF1 at a tertiary care center in India and evaluate the association between clinical severity and quality of life in these patients.

Methods: We conducted a cross-sectional study of 40 patients with NF1 over a period of two years at a tertiary hospital in western India. After obtaining consent, demographic and clinical information was collected from the patients and recorded in a pre-designed proforma. Quality of life was assessed by a validated Dermatology Life Quality Index (DLQI) questionnaire in languages understood by the patients and subsequently analyzed.

Results: This study included 40 patients at a mean age of 28.6 years, with a slight male predominance. The most frequently occurring lesions were café-au-lait macules, followed by neurofibromas and intertriginous freckling. The mean DLQI score was 12.35, implying a large effect on most patients' lives. Questions related to self-consciousness, embarrassment, and the influence of skin lesions on clothing choices had the highest scores, indicating a significant effect on social perception.

Conclusion: NF1 has a profound impact on a patient's quality of life, as evidenced by the high DLQI scores in our patient cohort. The early identification and management of such patients can help prevent further deterioration of their quality of life.

Categories: Dermatology, Psychiatry, Epidemiology/Public Health

Keywords: india, dermatology, neurofibromas, dlqi, quality of life (qol), neurofibromatosis 1

Introduction

Neurofibromatosis type 1 (NF1) is an inherited neuroectodermal disease, primarily defined by the presence of six or more café-au-lait macules, intertriginous freckles, two or more neurofibromas, Lisch nodules, bony defects like sphenoid dysplasia, and optic gliomas. It is an autosomal dominant condition with an estimated prevalence of one in 2500-3300 births, with 50% of cases occurring due to sporadic gene mutations [1]. Dermatological manifestations appear early in life and progress with age, but neurological features usually present at a later age [2]. The diagnosis of this condition is usually clinical, with genetic testing rarely being necessary. The cost and accessibility of genetic analysis, which is only available in a few selected tertiary care centers, are major barriers to the research and management of this disease in developing countries like India.

NF1 is a chronic disease that puts a large amount of stress on patients, which adversely impacts their identity, mental well-being, self-confidence, and social relationships. Patients with incurable disorders must cope with multiple impediments to attain a feasible level of health, as well as adequate physical, intellectual, and social functioning [3]. Mental disorders like anxiety, personality disorders, depression, and suicidal tendencies are common in individuals with NF1 [4]. In severe cases, the persistence of cognitive impairment into adulthood hinders a patient's vocational performance, thereby hampering their quality of life (QoL) [5]. In a study of NF1 patients in Iran, Foji et al. found that NF1 affected all aspects of the patient's life. In particular, perceived social alienation and difficulties regarding marriage and childbirth were present due to apprehensions regarding disease transmission to their progeny [6]. In short, dermatological diseases are a major contributor to the burden of non-fatal disease globally and influence different aspects of an individual's life [7]. Since the initial presentation of NF1 is mainly confined to the skin, patients usually

How to cite this article

Mahajan A, Patvekar M, Lote S, et al. (February 19, 2022) A Clinico-Epidemiological Study of Neurofibromatosis Type 1 and Its Relation to Quality of Life: A Cross-Sectional Study From India. Cureus 14(2): e22376. DOI 10.7759/cureus.22376

consult dermatologists first. Thus, dermatologists must especially be aware of the options for screening, genetic counseling, and prenatal diagnosis of the disease. A multidisciplinary approach must be adopted for the management of these patients [8-10].

In 1994, Finlay et al. developed the Dermatology Life Quality Index (DLQI) [11,12]. This is a 10-item questionnaire that covers six domains, specifically regarding symptoms and feelings, daily activities, leisure, work and school, personal relationships, and treatment. In this study, we aimed to evaluate the psychological impact of NF1 on an Indian population, gauge their QoL, and understand the extent of disease burden by using the DLQI questionnaire.

Materials And Methods

Study design

This cross-sectional study enrolled 40 patients with NF1 reporting to the dermatology clinic in a tertiary hospital in western India, across a span of 24 months. The study was approved by the Research and Recognition Committee of Dr. D. Y. Patil Medical College, Hospital and Research Centre (approval number: IESC/PGS/2019/43).

Data collection

Socio-demographic and clinical information was obtained from the patients after due consent and documented in a pre-designed proforma. Clinical history and examination of patients were followed by evaluation of their QoL, using a validated DLQI questionnaire (10-item questionnaire) in languages comprehensible to them.

Inclusion and exclusion criteria

All individuals clinically suspected to have NF1 as per the criteria of the National Institute of Health were included in the study. Patients not consenting to participate were excluded from the study. Age, sex, disease duration, and the presence or absence of any other major illnesses were not considered for inclusion or exclusion.

DLQI questionnaire

Each question of the DLQI questionnaire has five possible responses: “not relevant,” “not at all,” “a little,” “a lot,” or “very much,” which are scored as 0, 0, 1, 2, and 3, respectively. Question 7 is scored differently, with “prevented work or studying” being given a score of 3 [13]. The DLQI questionnaire is presented in the Appendix.

The final DLQI score was calculated as the sum of the scores of all questions, which has a range of 0-30. DLQI scores of 0-1, 2-5, 6-10, 11-20, and 21-30, respectively, correspond to having no, small, moderate, very large, and extremely large effect on the patient's life.

Statistical analysis

Data were entered in Microsoft Excel (Microsoft Corporation, Redmond, WA) and analyzed using the Statistical Package for the Social Sciences (SPSS) software version 20 (IBM Corp., Armonk, NY). Categorical variables were expressed in terms of frequency and percentage, whereas continuous variables were expressed as the mean and standard deviation (SD). For qualitative variables, the percentage and chi-square test were performed. Karl Pearson coefficient of correlation was also performed. A two-tailed probability value of 0.05 or less was considered statistically significant.

Results

This study included 40 patients with an age range of 6-65 (mean: 28.6) years, including 23 (57.5%) males and 17 (42.5%) females (Table 1). A family history of NF1 was positive in 19 (47.5%) patients, with seven (17.5%) patients reporting consanguineous marriage of their parents. Café-au-lait macules and neurofibromas, which were both seen in 38 (95%) patients, were the most commonly occurring lesions. Intertriginous freckling was seen in 28 (70%) patients, Lisch nodules in 19 (47.5%), plexiform neurofibromas in 15 (37.5%), and optic nerve tumors in one (2.5%) patient.

Factors	Variants	n	%
Age	10	2	5
	11-20	9	22.5
	21-30	16	40
	31-40	4	10
	41-50	8	20
	>51	1	2.5
Gender	Male	23	57.5
	Female	17	42.5
Family history of NF1	Positive	19	47.5
	Negative	21	52.5
Consanguineous marriage of parents	Present	7	17.5
	Absent	33	82.5

TABLE 1: Demographic factors of NF1 patients.

NF1, neurofibromatosis type 1.

The mean DLQI score was 12.35 ± 7.73 , which reflects a significant degree of impact on the QoL of most patients. Table 2 demonstrates the various categories of DLQI scores obtained in our cohort.

DLQI categories	n	%
No effect on patient's life	0	0
Small effect on patient's life	10	25
Moderate effect on patient's life	11	27.50
Very large effect on patient's life	10	25
Extremely large effect on patient's life	9	22.50

TABLE 2: DLQI scores in NF1.

DLQI, Dermatology Life Quality Index; NF1, neurofibromatosis type 1.

Table 3 depicts the mean scores of answers to individual questions of the DLQI questionnaire. Impairment of QoL, evaluated via the DLQI questionnaire, revealed that question 2 (i.e., “Over the last week, how embarrassed or self-conscious have you been because of your skin?”) had the highest mean score of 1.9, followed by question 4 (i.e., “Over the last week, how much has your skin influenced the clothes you wear?”), with a score of 1.85. On the other hand, question 6, related to difficulties in playing any sports, had the lowest average score of 0.45.

Question	1	2	3	4	5	6	7	8	9	10	Total
Mean	1	1.9	0.65	1.85	0.95	0.45	1.8	1.43	0.87	1.45	12.35
SD	0.894	0.943	0.823	0.989	1.047	0.705	1.47	0.891	1.053	0.921	7.738

TABLE 3: Mean scores of individual questions of DLQI.

DLQI, Dermatology Life Quality Index.

Tables 4, 5 demonstrate the comparison between the various demographic and clinical factors and DLQI categories. Age, presence of café-au-lait macules, Lisch nodules, freckling, and plexiform neurofibromas had a significant negative correlation with QoL ($p < 0.05$). Gender, family history of the disease, and parental consanguinity did not have a significant correlation with QoL.

		DLQI categories					P-value	
		Small effect	Moderate effect	Very large effect	Extremely large effect	Total		
Gender	Male	7	6	7	3	23	p = 0.35	
		30.43%	26.10%	30.43%	13.04%	100.00%		
	Female	3	5	3	6	17		
		17.65%	29.41%	17.65%	35.29%	100.00%		
Age	DLQI categories					P-value		
		Small effect	Moderate effect	Very large effect	Extremely large effect	Total	p = 0.0015	
	10	2	0	0	0	2		
		18.75%	50.00%	12.50%	18.75%	100.00%		
	31-40	1	0	0	3	4		
		25.00%	0.00%	0.00%	75.00%	100.00%		
	41-50	1	0	4	3	8		
		12.50%	0.00%	50.00%	37.50%	100.00%		
	>51	0	0	1	0	1		
		0.00%	0.00%	100.00%	0.00%	100.00%		
	Family history of NF1	DLQI categories						P-value
			Small effect	Moderate effect	Very large effect	Extremely large effect		Total
Positive		3	4	5	7	19		
		15.79%	21.05%	26.32%	36.84%	100.00%		
Negative		7	7	5	2	21		
		33.33%	33.33%	23.81%	9.53%	100.00%		
DLQI categories					P-value			
		Small effect	Moderate effect	Very large effect	Extremely large effect	Total	p = 0.34	
Present	0	2	2	3	7			
	0.00%	28.57%	28.57%	42.86%	100.00%			
Absent	10	9	8	6	33			
	30.30%	27.28%	24.24%	18.18%	100.00%			

TABLE 4: Comparison between demographic factors and DLQI categories.

DLQI, Dermatology Life Quality Index; NF1, neurofibromatosis type 1.

		DLQI categories					P-value
		Small effect	Moderate effect	Very large effect	Extremely large effect	Total	
CALMS	10	3	2	0	0	5	p < 0.0001
		60.00%	40.00%	0.00%	0.00%	100.00%	
	11-25	6	6	1	2	15	
		40.00%	40.00%	6.67%	13.33%	100.00%	
	26-50	0	3	6	4	13	
		0.00%	23.08%	46.15%	30.77%	100.00%	
	>50	0	0	2	3	5	
		0.00%	0.00%	40.00%	60.00%	100.00%	
	DLQI categories						P-value
NFs	10	6	1	1	0	8	p < 0.01
		75.00%	12.50%	12.50%	0.00%	100.00%	
	11-25	3	8	5	4	20	
		15.00%	40.00%	25.00%	20.00%	100.00%	
	26-50	0	1	2	3	6	
		0.00%	16.67%	33.33%	50.00%	100.00%	
	>50	0	0	2	2	4	
		0.00%	0.00%	50.00%	50.00%	100.00%	
	DLQI categories						P-value
Freckling	Present in	3	8	8	9	28	p < 0.001
		10.71%	28.57%	28.57%	32.15%	100.00%	
	DLQI categories						P-value
Lisch nodules	Present in	3	3	7	6	19	p = 0.007
		15.79%	15.79%	36.84%	31.58%	100.00%	
	DLQI categories						P-value
Plexiform NFs	Present in	0	1	6	8	15	p = 0.006
		0.00%	6.67%	40.00%	53.33%	100.00%	

TABLE 5: Comparison between clinical features and DLQI categories.

DLQI, Dermatology Life Quality Index; CALMS, café-au-lait macules; NFs- neurofibromas.

Discussion

Our study cohort had a slight male predominance, which was also seen in a study by Purkait et al. on patients with neurocutaneous syndromes [2]. The most frequently occurring lesions in our study were café-au-lait macules, seen in 95% of patients, which had a similar prevalence of 93% in a study by Lakshmanan et

al. on NF1 patients in South India. Furthermore, their study demonstrated freckling in 90% of patients, whereas in our group, this occurred in 70%. A family history of NF1 was positive in 47.5% of our patients, in contrast to 33% of patients in their study [14]. Cutaneous neurofibromas were seen in 95% of our patients, similar to the findings of Lakshmanan et al. and Wolkenstein et al., with an occurrence of 96.6% and 91%, respectively [10,14].

The presence of plexiform neurofibromas in our study was comparable to that of a study by Huson et al. (37.5% vs. 32%, respectively) [15]. On the other hand, Lisch nodules were present in 47.5% of patients in our study, which is lower than that in a study from Northwest England and a study by Friedman et al. (63% and 57%, respectively) [16,17].

The mean DLQI was 12.35 ± 7.73 , with higher QoL impairment in females (14 ± 8.66) than in males (11.13 ± 6.72). Higher DLQI scores were observed with more extensive skin involvement, female gender, and the presence of plexiform neurofibromas. Upon analysis of DLQI scores, NF1 had no effect in zero cases but had a small, moderate, very large, and extremely large effect in 10 (25%), 11 (27.5%), 10 (25%), and nine (22.5%) patients, respectively.

Similar studies from Italy and America demonstrated that patients with more visible lesions had a greater impact on the emotional aspect of their QoL [18,19]. A study from Brazil concluded that patients were significantly concerned about transmitting the disease to their progeny and expressed difficulties in obtaining appropriate treatment [20]. Various studies on the QoL of NF1 patients done in a few countries also reported a negative effect on QoL across multiple aspects of life [21,22]. A Canadian study of 176 patients reported that patients' mental well-being and overall QoL were influenced greatly by their perceived physical appearance [23].

The presence of café-au-lait macules, Lisch nodules, plexiform neurofibromas, and freckling had a significant negative correlation with QoL ($P < 0.05$). Individual questions of the DLQI questionnaire demonstrated the highest mean scores for questions 2 and 4 (1.9 and 1.85, respectively), followed by question 10 (1.8), indicating that self-consciousness, embarrassment, choice of clothing, and time taken by treatment were aspects of their lives most affected by the disease. Questions 3 and 6, pertaining to difficulty in playing sports and shopping/gardening, had the lowest mean scores, signifying a lower impact on these spheres of a patient's life.

Study limitations

More extensive questionnaires like the Skindex and 36-Item Short Form Survey (SF-36) were not used to evaluate the QoL due to the poor compliance of patients and the relatively lengthy nature of those questionnaires.

Conclusions

NF1 has a significant effect on patients' QoL due to changes in their appearance and societal stigma. Patients with comparatively fewer lesions can also experience a drastic impairment in their QoL, and their mental health must be considered just as much as a patient with extensive lesions. Early identification of patients with impaired QoL should prompt adequate therapeutic counseling and reference for a multidisciplinary approach.

Appendices

Dermatology Life Quality Index (DLQI) questionnaire

DERMATOLOGY LIFE QUALITY INDEX (DLQI) - INSTRUCTIONS FOR USE

The Dermatology Life Quality Index questionnaire is designed for use in adults, i.e. patients over the age of 16. It is self explanatory and can be simply handed to the patient who is asked to fill it in without the need for detailed explanation. It is usually completed in one or two minutes.

SCORING

The scoring of each question is as follows:

Very much	scored 3
A lot	scored 2
A little	scored 1
Not at all	scored 0
Not relevant	scored 0
Question 7, 'prevented work or studying'	scored 3

The DLQI is calculated by summing the score of each question resulting in a maximum of 30 and a minimum of 0. The higher the score, the more quality of life is impaired.

HOW TO INTERPRET MEANING OF DLQI SCORES

0 – 1	no effect at all on patient's life
2 – 5	small effect on patient's life
6 – 10	moderate effect on patient's life
11 – 20	very large effect on patient's life
21 – 30	extremely large effect on patient's life

REFERENCES

Finlay AY and Khan GK. Dermatology Life Quality Index (DLQI): a simple practical measure for routine clinical use. *Clin Exp Dermatol* 1994; **19**:210-216.

Basra MK, Fenech R, Gatt RM, Salek MS and Finlay AY. The Dermatology Life Quality Index 1994-2007: a comprehensive review of validation data and clinical results. *Br J Dermatol* 2008; **159**:997-1035.

Hongbo Y, Thomas CL, Harrison MA, Salek MS and Finlay AY. Translating the science of quality of life into practice: What do dermatology life quality index scores mean? *J Invest Dermatol* 2005; **125**:659-64.

There is more information about the DLQI, including over 85 translations, at www.dermatology.org.uk. The DLQI is copyright but may be used without seeking permission by clinicians for routine clinical purposes. For other purposes, please contact the copyright owners.

© AY Finlay, GK Khan, April 1992 www.dermatology.org.uk, this must not be copied without the permission of the authors.

FIGURE 1: Dermatology Life Quality Index (DLQI) questionnaire (interpretation).

DERMATOLOGY LIFE QUALITY INDEX (DLQI)

Hospital No: Date:
 Name: Score:
 Address: Diagnosis:

The aim of this questionnaire is to measure how much your skin problem has affected your life
 OVER THE LAST WEEK. Please tick (✓) one box for each question.

1. Over the last week, how itchy, sore, painful or stinging has your skin been?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	
2. Over the last week, how embarrassed or self conscious have you been because of your skin?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	
3. Over the last week, how much has your skin interfered with you going shopping or looking after your home or garden?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐
4. Over the last week, how much has your skin influenced the clothes you wear?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐
5. Over the last week, how much has your skin affected any social or leisure activities?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐
6. Over the last week, how much has your skin made it difficult for you to do any sport?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐
7. Over the last week, has your skin prevented you from working or studying?	Yes ☐	
	No ☐	Not relevant ☐
If "No", over the last week how much has your skin been a problem at work or studying?	A lot ☐	
	A little ☐	
	Not at all ☐	
8. Over the last week, how much has your skin created problems with your partner or any of your close friends or relatives?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐
9. Over the last week, how much has your skin caused any sexual difficulties?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐
10. Over the last week, how much of a problem has the treatment for your skin been, for example by making your home messy, or by taking up time?	Very much ☐	
	A lot ☐	
	A little ☐	
	Not at all ☐	Not relevant ☐

Please check you have answered EVERY question. Thank you.

© AY Finlay, GK Khan, April 1992 www.dermatology.org.uk, this must not be copied without the permission of the authors.

FIGURE 2: Dermatology Life Quality Index (DLQI) questionnaire.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. Research and Recognition Committee, Dr. D. Y. Patil Medical College, Hospital and Research Centre issued approval IESC/PGS/2019/43. The Research and Recognition Committee, under the Faculty of Medicine, at its meeting held on 8th November 2019, has approved the research. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Griffiths CEM, Creamer D, Barker J, Chaimers R, Bleiker T: Rook's Textbook of Dermatology, 9th Edition. Wiley, Hoboken, New Jersey; 2016.
https://www.google.co.in/books/edition/Rook_s_Textbook_of_Dermatology/EyypCwAAQBAJ?hl=en.
- Purkait R, Samanta T, Thakur S, Dhar S: Neurocutaneous syndrome: a prospective study. Indian J Dermatol.

- 2011, 56:375-9. [10.4103/0019-5154.84721](#)
3. Klein-Tasman BP, Colon AM, Brei N, et al.: Adaptive behavior in young children with neurofibromatosis type 1. *Int J Pediatr*. 2013, 2013:690432. [10.1155/2013/690432](#)
4. Wang DL, Smith KB, Esparza S, Leigh FA, Muzikansky A, Park ER, Plotkin SR: Emotional functioning of patients with neurofibromatosis tumor suppressor syndrome. *Genet Med*. 2012, 14:977-82. [10.1038/gim.2012.85](#)
5. Pasini A, Lo-Castro A, Di Carlo L, Pitzianti M, Siracusano M, Rosa C, Galasso C: Detecting anxiety symptoms in children and youths with neurofibromatosis type I. *Am J Med Genet B Neuropsychiatr Genet*. 2012, 159B:869-73. [10.1002/ajmg.b.32095](#)
6. Foji S, Mohammadi E, Sanagoo A, Jouybari L: The patients' experiences of burden of neurofibromatosis: a qualitative study. *Iran J Nurs Midwifery Res*. 2021, 26:342-8. [10.4103/ijnmr.IJNMR_178_20](#)
7. Hay RJ, Johns NE, Williams HC, et al.: The global burden of skin disease in 2010: an analysis of the prevalence and impact of skin conditions. *J Invest Dermatol*. 2014, 134:1527-34. [10.1038/jid.2013.446](#)
8. Ablon J: Gender response to neurofibromatosis 1. *Soc Sci Med*. 1996, 42:99-109. [10.1016/0277-9536\(95\)00076-3](#)
9. Wolkenstein P, Durand-Zaleski I, Moreno JC, Zeller J, Hemery F, Revuz J: Cost evaluation of the medical management of neurofibromatosis 1: a prospective study on 201 patients. *Br J Dermatol*. 2000, 142:1166-70. [10.1046/j.1365-2133.2000.03543.x](#)
10. Wolkenstein P, Zeller J, Revuz J, Ecosse E, Leplège A: Quality-of-life impairment in neurofibromatosis type 1: a cross-sectional study of 128 cases. *Arch Dermatol*. 2001, 137:1421-5. [10.1001/archderm.137.11.1421](#)
11. Finlay AY, Khan GK: Dermatology Life Quality Index (DLQI)—a simple practical measure for routine clinical use. *Clin Exp Dermatol*. 1994, 19:210-6. [10.1111/j.1365-2230.1994.tb01167.x](#)
12. Basra MK, Fenech R, Gatt RM, Salek MS, Finlay AY: The Dermatology Life Quality Index 1994-2007: a comprehensive review of validation data and clinical results. *Br J Dermatol*. 2008, 159:997-1035. [10.1111/j.1365-2133.2008.08832.x](#)
13. Hongbo Y, Thomas CL, Harrison MA, Salek MS, Finlay AY: Translating the science of quality of life into practice: what do dermatology life quality index scores mean?. *J Invest Dermatol*. 2005, 125:659-64. [10.1111/j.0022-202X.2005.23621.x](#)
14. Lakshmanan A, Bubna AK, Sankarasubramaniam A, Veeraraghavan M, Rangarajan S, Sundaram M: A clinical study of neurofibromatosis-1 at a tertiary health care centre in south India. *Pigment Int*. 2016, 3:102-7. [10.4103/2349-5847.196302](#)
15. Huson SM, Harper PS, Compston DA: Von Recklinghausen neurofibromatosis. A clinical and population study in south-east Wales. *Brain*. 1988, 111:1355-81. [10.1093/brain/111.6.1355](#)
16. McGaughran J, Harris D, Donnai D, et al.: A clinical study of type 1 neurofibromatosis in north west England. *J Med Genet*. 1999, 36:197-203.
17. Friedman JM, Birch PH: Type 1 neurofibromatosis: a descriptive analysis of the disorder in 1,728 patients. *Am J Med Genet*. 1997, 70:138-43. [10.1002/\(sici\)1096-8628\(19970516\)70:2<138::aid-ajmg7>3.0.co;2-u](#)
18. Kodra Y, Giustini S, Divona L, Porciello R, Calvieri S, Wolkenstein P, Taruscio D: Health-related quality of life in patients with neurofibromatosis type 1. A survey of 129 Italian patients. *Dermatology*. 2009, 218:215-20. [10.1159/000187594](#)
19. Page PZ, Page GP, Ecosse E, Korf BR, Leplege A, Wolkenstein P: Impact of neurofibromatosis 1 on quality of life: a cross-sectional study of 176 American cases. *Am J Med Genet A*. 2006, 140A:1893-8. [10.1002/ajmg.a.31422](#)
20. Bicudo NP, de Menezes Neto BF, da Silva de Avó LR, Germano CM, Melo DG: Quality of life in adults with neurofibromatosis 1 in Brazil. *J Genet Couns*. 2016, 25:1063-74. [10.1007/s10897-016-9939-8](#)
21. Sanagoo A, Jouybari L, Koochi F, Sayehmiri F: Evaluation of QoL in neurofibromatosis patients: a systematic review and meta-analysis study. *BMC Neurol*. 2019, 19:123. [10.1186/s12883-019-1338-y](#)
22. Lai J, Beaumont JL, Jensen S: Quality of life for patients with neurofibromatosis type 1 associated plexiform neurofibromas (PNF). *Value Health*. 2017, 20:727. [10.1016/j.jval.2017.08.1970](#)
23. Hamoy-Jimenez G, Kim R, Suppiah S, Zadeh G, Bril V, Barnett C: Quality of life in patients with neurofibromatosis type 1 and 2 in Canada. *Neurooncol Adv*. 2020, 2:i141-9. [10.1093/oaajnl/vdaa003](#)