Review began 12/10/2021
Review ended 12/20/2021
Published 12/22/2021

© Copyright 2021
Barros et al. This is an open access article
distributed under the terms of the Creative
Commons Attribution License CC-BY 4.0.,
which permits unrestricted use, distribution,
and reproduction in any medium, provided
the original author and source are credited.

Middle Meningeal Artery Embolization as an Adjuvant Treatment for Bilateral Chronic Subdural Hematomas From Spontaneous Intracranial Hypotension: A Case Report

Guilherme Barros¹, Robert H. Bonow¹, Michael R. Levitt¹

¹. Department of Neurological Surgery, University of Washington, Seattle, USA

Corresponding author: Michael R. Levitt, respub2@uw.edu

Abstract

Middle meningeal artery embolization (MMAE) is an effective initial or adjuvant treatment for chronic subdural hematomas. However, its efficacy in the setting of spontaneous intracranial hypotension has not yet been reported. We present a case of progressive bilateral chronic subdural hematomas secondary to spontaneous intracranial hypotension that failed the initial lumbar epidural blood patch. The patient underwent surgical hematoma evacuation, with adjuvant MMAE and additional epidural blood patch, with good clinical and radiographic results. This case demonstrates the potential value of MMAE for chronic subdural hematomas secondary to spontaneous intracranial hypotension.

Categories: Neurosurgery

Keywords: case report, subdural hematoma, intracranial hypotension, middle meningeal artery, embolization

Introduction

Chronic subdural hematomas (cSDH) secondary to spontaneous intracranial hypotension (SIH) can be challenging to manage. They are present in approximately 50% of SIH patients [1]. Despite epidural blood patches (EBP) to treat the underlying SIH and/or surgical cSDH evacuation, they have a significant rate of recurrence or progression [2-4]. Middle meningeal artery embolization (MMAE) has emerged as a standalone therapy or adjunct to surgical evacuation for cSDH. However, MMAE has not been previously reported as a treatment for cSDH specifically due to SIH. We describe a patient presenting with SIH and progressive bilateral cSDH, ultimately treated successfully with a combination of surgical evacuation, MMAE, and EBP.

Case Presentation

An otherwise healthy male in his 50s presented with several weeks of 8/10 intensity postural headaches. There was no history of trauma, spine surgery, lumbar puncture, pharmacologic anticoagulation, connective tissue disorder, or hereditary/acquired coagulopathy. Computed tomography (CT) of the head demonstrated bilateral cSDH, which mildly enlarged on follow-up.

Following the initial CT, the patient underwent magnetic resonance imaging (MRI) of the brain and cervical, thoracic, and lumbar spine, including MRI myelography. MRI of the brain (Figure 1) redemonstrated bilateral cSDH, in addition to flattening of the pontomesencephalic angle.

How to cite this article

Barros G, Bonow R H, Levitt M R (December 22, 2021) Middle Meningeal Artery Embolization as an Adjuvant Treatment for Bilateral Chronic Subdural Hematomas From Spontaneous Intracranial Hypotension: A Case Report. Cureus 13(12): e20609. DOI 10.7759/cureus.20609

FIGURE 1: Initial consultation, non-contrast magnetic resonance imaging of the brain.

(A) T2 axial plane demonstrating bilateral subdural hematomas (black arrows). (B) T1 sagittal plane demonstrating flattened pontomesencephalic angle (white arrow).

The MRI myelogram showed several right-sided thoracic neuroforaminal cysts, shown in Figure 2, with no definitive site of cerebrospinal fluid leak. Given the patient's postural headaches, lack of history of head trauma, and imaging findings, there was high clinical suspicion for SIH.

FIGURE 2: Axial reformatting MRI thoracic spine myelogram.

Arrows show right-sided perineural cysts within the neuroforamina of (A) T4-T5, (B) T7-T8, and (C) T9-T10.

The patient underwent a lumbar EBP as the initial treatment. This resulted in subjective improvement in headaches. A follow-up CT one week after the EBP showed mildly reduced cSDH size. Three weeks after EBP, CT showed interval growth of the cSDH. The patient presented to the emergency room several days later with new-onset transient perioral numbness and speech arrest, concerning seizure. Additional CT (Figure 3) showed right-to-left midline shift and further progression of the bilateral cSDH.

FIGURE 3: Emergency department presentation, non-contrast computed tomography of the head.

(A) Axial plane. (B) Coronal plane. Enlarging bilateral subdural hematomas (white arrows) and right-to-left midline shift (black arrow).

Levetiracetam monotherapy was initiated. The patient underwent simultaneous right-sided craniotomy and left-sided burr holes for cSDH evacuation. A small craniotomy was required on the right side after the initial attempt with burr holes was unsuccessful in evacuating the hematoma due to extensive membranes. Postoperative CT showed small residual cSDH bilaterally, shown in Figure 4.

FIGURE 4: Immediate postoperative non-contrast CT of the head after right-side craniotomy and left-side burr holes for bilateral subdural hematoma evacuation.

(A) Axial section and (B) coronal section showing small residual subdural hematoma bilaterally (arrows).

The next day, the patient underwent bilateral MMAE with Onyx, and on postoperative day two, he underwent a second lumbar EBP. These additional procedures were done following surgery to attempt a multimodal approach at treating his refractory subdural hematomas secondary to intracranial hypotension. The patient was discharged home on postoperative day three, with minimal headache and no further seizures. At six-week follow-up, the patient reported no headache and occasional seizure auras, with no actual seizures confirmed on an outpatient electroencephalogram (EEG). Head CT (Figure 5) showed significantly decreased bilateral cSDH with minimal residual volume.

FIGURE 5: Six-week postoperative follow-up, non-contrast computed tomography of the head.

(A) Axial plane. (B) Coronal plane. Minimal residual bilateral subdural hematomas (arrows) following surgical evacuation and middle meningeal artery embolization.

Discussion

Management of cSDH due to SIH often requires a multimodal approach. Initial conservative measures include bed rest, hydration, caffeine, and close follow-up brain imaging to ensure there is no progression [1]. However, this has limited clinical efficacy, with reports of only 18%-23% experiencing favorable outcomes [2,3]. Patients may undergo myelography to localize a potential source of cerebrospinal fluid leak, followed by first-line intervention with a blind or targeted EBP [4]. The effectiveness of single or multiple EBP alone for the treatment of cSDH due to SIH without surgery ranges from 63% to 72% [2-4]. Surgical evacuation of the cSDH is usually reserved for progressive enlargement, neurologic deterioration, and/or lack of improvement of SIH symptoms despite first-line therapy, as in the current case. In cases where surgical evacuation is performed as the initial treatment for the cSDH for SIH, the recurrence rate is especially high, with up to two-thirds of patients requiring additional evacuations with or without perioperative EBP [2]. However, surgical evacuation success is significantly improved if EBP is completed perioperatively, with up to 77%-85% exhibiting no cSDH recurrence and symptomatic improvement of SIH [3,4].

Further treatment modalities are needed to treat difficult-to-treat progressive or recurrent cSDH in the SIH setting, especially for those patients failing traditional first-line treatments. The use of MMAE for cSDH has not been reported specifically in the setting of SIH. Among a generalized cSDH patient cohort containing surgically naïve, recurrent after prior surgery, and adjuvant immediately following surgery, Link et al. reported 91.1% stable or decreased hematoma size without the need for further surgery after MMAE [5]. In a multi-center, prospective study with 138 patients undergoing MMAE as initial therapy or for recurrence after surgical evacuation, there was a 70.8% radiographic improvement of at least 50% volume reduction, with only 6.5% of patients requiring further cSDH treatments [6]. Comparing surgical evacuation of cSDH alone to surgical evacuation with adjuvant MMAE, MMAE improved mean volume reduction of the hematoma by 17.5 mL [7].

Conclusions

In our patient with neurologic deterioration and radiographic progression of cSDH despite initial EBP, a multimodal treatment approach was employed. Surgical evacuation of the hematomas was indicated, and adjuvant MMAE with additional EBP likely assisted in preventing recurrence of the cSDH. Furthermore, the patient achieved immediate symptomatic improvement of the chronic headache, which was sustained at follow-up. This report highlights the novel use of adjunct MMAE for treating cSDH in the setting of SIH.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** Michael Levitt declare(s) stock/stock options from Cerebrotech. Michael Levitt declare(s) stock/stock options from Proprio. Michael Levitt declare(s) stock/stock options

from Hyperion Surgical. Michael Levitt declare(s) stock/stock options from Synchron. Michael Levitt declare(s) personal fees from Medtronic. Michael Levitt declare(s) personal fees from Metis Innovative. Michael Levitt declare(s) Unrestricted educational grant from Medtronic. Michael Levitt declare(s) Unrestricted educational grant from Stryker. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

The authors appreciate the assistance of Sharon Durfy, PhD, with manuscript preparation.

References

1. Schievink WI: Spontaneous spinal cerebrospinal fluid leaks and intracranial hypotension . JAMA. 2006, 295:2286-96. [10.1001/jama.295.19.2286](https://doi.org/10.1001/jama.295.19.2286)
2. Takahashi K, Mima T, Akiba Y: Chronic subdural hematoma associated with spontaneous intracranial hypotension: therapeutic strategies and outcomes of 55 cases. Neurol Med Chir (Tokyo). 2016, 56:69-76. [10.2176/nmc.oa.2015-0032](https://doi.org/10.2176/nmc.oa.2015-0032)
3. Chen YC, Wang YF, Li JY, et al.: Treatment and prognosis of subdural hematoma in patients with spontaneous intracranial hypotension. Cephalalgia. 2016, 36:225-31. [10.1177/0333102415585095](https://doi.org/10.1177/0333102415585095)
4. Rettenmaier LA, Park BJ, Holland MT, Hamade YJ, Garg S, Rastogi R, Reddy CG: Value of targeted epidural blood patch and management of subdural hematoma in spontaneous intracranial hypotension: case report and review of the literature. World Neurosurg. 2017, 97:27-38. [10.1016/j.wneu.2016.09.076](https://doi.org/10.1016/j.wneu.2016.09.076)
5. Link TW, Boddu S, Paine SM, Kamel H, Knopman J: Middle meningeal artery embolization for chronic subdural hematoma: a series of 60 cases. Neurosurgery. 2019, 85:801-7. [10.1093/neuros/nyy521](https://doi.org/10.1093/neuros/nyy521)
6. Kan P, Maragkos GA, Srivatsan A, et al.: Middle meningeal artery embolization for chronic subdural hematoma: a multi-center experience of 154 consecutive embolizations. Neurosurgery. 2021, 88:268-77. [10.1093/neuros/nyaa379](https://doi.org/10.1093/neuros/nyaa379)
7. Ng S, Derraz I, Boetto J, et al.: Middle meningeal artery embolization as an adjuvant treatment to surgery for symptomatic chronic subdural hematoma: a pilot study assessing hematoma volume resorption. J Neurointerv Surg. 2020, 12:695-9. [10.1136/neurintsurg-2019-015421](https://doi.org/10.1136/neurintsurg-2019-015421)