Review began 01/21/2022
Review ended 01/31/2022
Published 02/04/2022

© Copyright 2022
Kassan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Gut Microbiota Regulates the Sympathetic Nerve Activity and Peripheral Serotonin Through Hypothalamic MicroRNA-204 in Order to Increase the Browning of White Adipose Tissue in Obesity

Adam Kassan¹, Karima Ait-Aissa², Modar Kassan³

1. Pharmacy, West Coast University, Los Angeles, USA 2. Internal Medicine, University of Iowa, Iowa City, USA 3. Physiology, University of Tennessee Health Science Center, Memphis, USA

Corresponding author: Modar Kassan, modarokassan@gmail.com

Abstract

The prevalence of obesity is increasing worldwide, and novel therapeutic strategies such as enhancement of thermogenic pathways in white adipose tissue (WAT) are gaining more attention. The gut/brain axis plays an essential role in promoting the browning of WAT. However, the mechanism by which this axis regulates WAT function is not fully understood. On the other hand, the role of microRNAs (miRNAs) in the control of WAT browning has already been established. Therefore, understanding the communication pathways linking the gut/brain axis and miRNAs might establish a promising intervention for obesity.

Our published data showed that microRNA-204 (miR-204), a microRNA that plays an important role in the control of the central nervous system (CNS) and the pathogenesis of obesity, is affected by gut dysbiosis. Therefore, miR-204 could be a key element that controls the browning of WAT by acting as a potential link between the gut microbiota and the brain. In this review, we summarized the current knowledge about communication pathways between the brain, gut, and miR-204 and examined the literature to discuss potential research directions that might lead to a better understanding of the mechanisms underlying the browning of WAT in obesity.

Categories: Endocrinology/Diabetes/Metabolism

Keywords: enterochromaffin cells, 5-HT, browning of white adipose tissue, hypothalamic miR-204, obesity

Introduction And Background

The prevalence of obesity and associated metabolic disorders is increasing worldwide, and novel therapeutic strategies are needed. Recently, increasing energy expenditure through thermogenesis of white adipose tissue (WAT), known as browning/beiging of WAT, has been recognized as a therapeutic strategy for weight loss in humans [1]. Over the last decade, extensive studies in the area of metabolic disorders pointed toward gut microbiota as an important player in regulating the browning of WAT [2]. Another component that plays an important role in regulating WAT thermogenesis is the brain [3]. The gut/brain axis is a bidirectional communication between the central nervous system (CNS) and the gastrointestinal tract [4]. Several peripheral signals relaying information about energy status act in the brain, particularly the hypothalamus, to regulate the browning of WAT [5]. It has been shown that these signals, responsible for the control of systemic energy balance, are regulated by hypothalamic miRNAs, key regulators of gene expression [6,7]. Additionally, gut microbiota can affect the expression of central miRNAs [8,9]. Thus, any miscommunication between the gut and the brain could affect these peripheral signals and might lead to metabolic disorders.

Interestingly, our published data showed that microRNA-204 (miR-204), a microRNA that plays an important role in the CNS function [10] and the pathophysiology of obesity [11], is indeed affected by gut dysbiosis [11]. Therefore, we speculated that miR-204 could be a key element connecting the gut microbiota to the brain in order to control the WAT thermogenesis during obesity. In this review, we will summarize what has been published about the communication pathways between the brain, gut, and miR-204. Based on the current knowledge, we will also examine potential research directions that might lead to a better understanding of the mechanisms underlying the browning of WAT in obesity. Based on the different findings in the literature, we will suggest a potential hypothesis by which hypothalamic miR-204 could emerge as a key player in regulating obesity through the browning of WAT.

Review

Hypothalamic miR-204 and sympathetic nerve activity

Hypothalamus is one of the most important areas in the CNS that regulate energy homeostasis [12], but the exact mechanism by which the hypothalamus regulates energy expenditure is not well established. Lately, it has been shown that the sympathetic nervous system (SNS) plays a key role in promoting the browning of

How to cite this article

Kassan A, Ait-Aissa K, Kassan M (February 04, 2022) Gut Microbiota Regulates the Sympathetic Nerve Activity and Peripheral Serotonin Through Hypothalamic MicroRNA-204 in Order to Increase the Browning of White Adipose Tissue in Obesity. Cureus 14(2): e21913. DOI 10.7759/cureus.21913

WAT [13]. Recent scientific publications have reported that many key elements in the hypothalamus could affect the SNS [14,15]. One important element is the nicotinamide adenine dinucleotide (NAD⁺)-dependent histone/protein deacetylase or simply sirtuin 1 (SIRT1), which is widely expressed in many tissues including the hypothalamus [16]. A study by Liu et al. showed that inhibition of hypothalamic SIRT1 led to a decrease in the sympathetic nerve activity (SNA) [14]. Brain-derived neurotrophic factor (BDNF), another element that is expressed in the hypothalamus, has also been shown to regulate SNA [15,17]. Recent studies showed that increased levels of BDNF in the paraventricular nucleus of the hypothalamus (PVN) led to increased sympathetic activity [15,17].

Furthermore, we and others have already demonstrated that miR-204 can target and downregulate SIRT1 [11,18], while other studies showed that miR-204 can target and downregulate BDNF [19-23]. We and others showed that miR-204 increases during obesity [11,24-26], whereas SIRT1 and BDNF are reduced in obesity [27-30]. Additionally, it has already been shown that miR-204 is expressed in the hypothalamus [31]. Based on this literature evidence, we can speculate that increased hypothalamic miR-204 targets and downregulates SIRT1 and BDNF, which in turn could lead to a decrease in the SNA.

SNA, enterochromaffin cells (ECs), peripheral serotonin (5-HT), and thermogenesis

Peripheral serotonin (5-HT) (80%-90% of the total body content of serotonin) is synthesized in the gut predominately by the enterochromaffin cells (ECs) from tryptophan-by-tryptophan hydroxylase 1 (Tph1) [32-35]. Interestingly, the release of 5-HT from the ECs, the most abundant enteroendocrine cell subtype of the colon [36], is under the control of the autonomous nervous system [37]. ECs are known to respond well to several physiological compounds such as norepinephrine (NE) [38], which is released by sympathetic nerve fibers in the gut [39], most specifically, the colon [40]. ECs are known to express the alpha 2 (α 2) adrenergic receptor [38,41]. Once NE binds to its α 2 receptor, it leads to a reduction in cyclic adenosine monophosphate (cAMP) [42]. It is important to mention that cAMP is critical for Tph1 function [43,44] and stability [45]; so, a decrease in cAMP will affect the Tph1 function [43,44], leading to less peripheral 5-HT release [45]. Additionally, studies performed in vivo indicate that α adrenergic agonists inhibit the release of peripheral 5-HT from ECs [41]. It is well known that during obesity, peripheral 5-HT is increased [46] and blunts the thermogenesis by negatively regulating the sensitivity of adipose tissues to β -adrenergic stimulation [47] that is known to stimulate peroxisome proliferator-activated receptors gamma (PPAR γ) and peroxisome proliferator-activated receptor-gamma coactivator 1 α (PGC1 α), which are master regulators of uncoupling protein 1 (UCP1)-mediated thermogenesis [48,49]. Interestingly, downregulation of peripheral 5-HT reduced obesity and promoted the browning of adipose tissue [47,50-51].

We already explained in the previous section, and based on literature findings, that an increase in miR-204 in obesity could lead to downregulation of SIRT1 and BDNF and, therefore, to a decrease in the SNA. Taken all together, we can hypothesize that increased hypothalamic miR-204 in obesity will lead to less NE binding to α 2 receptors in ECs. This eventually will increase cAMP that can activate and stabilize Tph1 leading to more peripheral 5-HT release and, therefore, will destabilize the thermogenesis pathway in WAT. Thus, the hypothalamic miR-204/ peripheral 5-HT axis might be effective for treating obesity and increasing thermogenesis.

Microbiota, miR-204, and peripheral serotonin

Manipulating bacterial composition using antibiotics or germ-free mice (GFM) showed that depletion of bacteria significantly increased the browning of WAT [52,53]. Additionally, mice on antibiotics or GFM showed a decreased level of peripheral 5-HT [54,55], and GFM modulated the sympathetic activity [56]. For instance, it has been shown that antibiotics increased the BDNF [57]. Moreover, GFM or treatment with antibiotics [11] showed decreased levels of miR-204 and increased levels of SIRT1 [11]. Interestingly, recolonization of these mice or fecal transplant from obese mice reversed all these parameters (Figure 1) [11].

FIGURE 1: Scheme summarizing the hypothesis in this review

miR-204, MicroRNA-204; NAD⁺, nicotinamide adenine dinucleotide; SIRT1, sirtuin 1; BDNF, brain-derived neurotrophic factor; NE, norepinephrine; α_2 , alpha 2; Tph1, tryptophan hydroxylase 1; 5-HT, serotonin; PPAR γ , peroxisome proliferator-activated receptors gamma; PGC1 α , peroxisome proliferator-activated receptor-gamma coactivator 1 α ; UCP1, uncoupling protein 1; WAT, white adipose tissue.

Conclusions

Taken all together and based on the literature discussed above, we can hypothesize that obesity could increase hypothalamic miR-204 leading to decreased levels of SIRT1 and BDNF, which in turn will lead to a decreased SNA. A decrease in SNA means that less NE is binding to α_2 receptors in ECs, leading to more cAMP that stabilizes and activates Tph1 to produce more peripheral 5-HTs that blunt the thermogenesis by negatively regulating the sensitivity of adipose tissues to β -adrenergic stimulation. Decreasing the β -adrenergic stimulation will lead to less PPAR γ and PGC1 α that are master regulators of UCP1-mediated thermogenesis, thus producing less thermogenesis.

Additional Information

Disclosures

Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Thyagarajan B, Foster MT: Beiging of white adipose tissue as a therapeutic strategy for weight loss in humans. *Horm Mol Biol Clin Investig.* 2017, 31:1-13. [10.1515/hmbci-2017-0016](https://doi.org/10.1515/hmbci-2017-0016)
- Moreno-Navarrete JM, Fernandez-Real JM: The gut microbiota modulates both browning of white adipose tissue and the activity of brown adipose tissue. *Rev Endocr Metab Disord.* 2019, 20:387-97. [10.1007/s11154-019-09523-x](https://doi.org/10.1007/s11154-019-09523-x)
- Nedergaard J, Cannon B: The browning of white adipose tissue: some burning issues. *Cell Metab.* 2014, 20:396-407. [10.1016/j.cmet.2014.07.005](https://doi.org/10.1016/j.cmet.2014.07.005)
- Carabotti M, Scirocco A, Maselli MA, Severi C: The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. *Ann Gastroenterol.* 2015, 28:203-209.
- Contreras C, Nogueiras R, Diéguez C, Rahmouni K, López M: Traveling from the hypothalamus to the adipose tissue: the thermogenic pathway. *Redox Biol.* 2017, 12:854-63. [10.1016/j.redox.2017.04.019](https://doi.org/10.1016/j.redox.2017.04.019)
- Cao L, Choi EY, Liu X, Martin A, Wang C, Xu X, During MJ: White to brown fat phenotypic switch induced by genetic and environmental activation of a hypothalamic-adipocyte axis. *Cell Metab.* 2011, 14:324-38. [10.1016/j.cmet.2011.06.020](https://doi.org/10.1016/j.cmet.2011.06.020)
- Schneeberger M, Gomez-Valadés AG, Ramirez S, Gomis R, Claret M: Hypothalamic miRNAs: emerging roles

- in energy balance control. *Front Neurosci.* 2015, 9:41. [10.3389/fnins.2015.00041](#)
8. Hoban AE, Stilling RM, Moloney G, et al.: Microbial regulation of microRNA expression in the amygdala and prefrontal cortex. *Microbiome.* 2017, 5:102. [10.1186/s40168-017-0321-3](#)
9. Chen JJ, Zeng BH, Li WW, et al.: Effects of gut microbiota on the microRNA and mRNA expression in the hippocampus of mice. *Behav Brain Res.* 2017, 322:34-41. [10.1016/j.bbr.2017.01.021](#)
10. Bharambe HS, Paul R, Panwalkar P, et al.: Downregulation of miR-204 expression defines a highly aggressive subset of group 3/group 4 medulloblastomas. *Acta Neuropathol Commun.* 2019, 7:52. [10.1186/s40478-019-0697-3](#)
11. Vikram A, Kim YR, Kumar S, Li Q, Kassan M, Jacobs JS, Irani K: Vascular microRNA-204 is remotely governed by the microbiome and impairs endothelium-dependent vasorelaxation by downregulating Sirtuin1. *Nat Commun.* 2016, 7:12565. [10.1038/ncomms12565](#)
12. Schneeberger M, Gomis R, Claret M: Hypothalamic and brainstem neuronal circuits controlling homeostatic energy balance. *J Endocrinol.* 2014, 220:T25-46. [10.1530/JOE-13-0398](#)
13. Cao Q, Jing J, Cui X, Shi H, Xue B: Sympathetic nerve innervation is required for beigeing in white fat. *Physiol Rep.* 2019, 7:e14031. [10.14814/phy2.14031](#)
14. Liu X, Zheng H: Modulation of Sirt1 and FoxO1 on hypothalamic leptin-mediated sympathetic activation and inflammation in diet-induced obese rats. *J Am Heart Assoc.* 2021, 10:e020667. [10.1161/JAHA.120.020667](#)
15. Schaich CL, Wellman TL, Koi B, Erdos B: BDNF acting in the hypothalamus induces acute pressor responses under permissive control of angiotensin II. *Auton Neurosci.* 2016, 197:1-8. [10.1016/j.autneu.2016.02.011](#)
16. Nillni EA: The metabolic sensor Sirt1 and the hypothalamus: Interplay between peptide hormones and pro-hormone convertases. *Mol Cell Endocrinol.* 2016, 438:77-88. [10.1016/j.mce.2016.09.002](#)
17. Erdos B, Backes I, McCowan ML, Hayward LF, Scheuer DA: Brain-derived neurotrophic factor modulates angiotensin signaling in the hypothalamus to increase blood pressure in rats. *Am J Physiol Heart Circ Physiol.* 2015, 308:H612-22. [10.1152/ajpheart.00776.2014](#)
18. Zhang L, Wang X, Chen P: MiR-204 down regulates SIRT1 and reverts SIRT1-induced epithelial-mesenchymal transition, anoikis resistance and invasion in gastric cancer cells. *BMC Cancer.* 2013, 13:290. [10.1186/1471-2407-13-290](#)
19. Ge QD, Tan Y, Luo Y, Wang WJ, Zhang H, Xie C: MiR-132, miR-204 and BDNF-TrkB signaling pathway may be involved in spatial learning and memory impairment of the offspring rats caused by fluorine and aluminum exposure during the embryonic stage and into adulthood. *Environ Toxicol Pharmacol.* 2018, 63:60-8. [10.1016/j.etap.2018.08.011](#)
20. Liu H, Wang J, Yan R, et al.: MicroRNA-204-5p mediates sevoflurane-induced cytotoxicity in HT22 cells by targeting brain-derived neurotrophic factor. *Histol Histopathol.* 2020, 35:1353-61. [10.14670/HH-18-266](#)
21. Luo Y, Li C, Xie C, Tan Y: Effects of fluorine exposure on miR-204 and BDNF-TrkB pathways in HT22 hippocampal neurons. *Chin J Endem.* 2020, 12:172-7.
22. Imam JS, Plyler JR, Bansal H, et al.: Genomic loss of tumor suppressor miRNA-204 promotes cancer cell migration and invasion by activating AKT/mTOR/Rac1 signaling and actin reorganization. *PLoS One.* 2012, 7:e52397. [10.1371/journal.pone.0052397](#)
23. Zhang Q, Su J, Kong W, et al.: Roles of miR-10a-5p and miR-103a-3p, regulators of BDNF expression in follicular fluid, in the outcomes of IVF-ET. *Front Endocrinol (Lausanne).* 2021, 12:637384. [10.3389/fendo.2021.637384](#)
24. Kurylowicz A: microRNAs in human adipose tissue physiology and dysfunction. *Cells.* 2021, 10:3342. [10.3390/cells10123342](#)
25. Zhang Y, Gu M, Ma Y, Peng Y: LncRNA TUG1 reduces inflammation and enhances insulin sensitivity in white adipose tissue by regulating miR-204/SIRT1 axis in obesity mice. *Mol Cell Biochem.* 2020, 475:171-83. [10.1007/s11010-020-03869-6](#)
26. Wang H, Chen Y, Mao X, Du M: Maternal obesity impairs fetal mitochondriogenesis and brown adipose tissue development partially via upregulation of miR-204-5p. *Biochim Biophys Acta Mol Basis Dis.* 2019, 1865:2706-15. [10.1016/j.bbadis.2019.07.012](#)
27. Sasaki T, Kikuchi O, Shimpuku M, et al.: Hypothalamic SIRT1 prevents age-associated weight gain by improving leptin sensitivity in mice. *Diabetologia.* 2014, 57:819-31. [10.1007/s00125-013-3140-5](#)
28. Sasaki T: Age-associated weight gain, leptin, and SIRT1: a possible role for hypothalamic SIRT1 in the prevention of weight gain and aging through modulation of leptin sensitivity. *Front Endocrinol (Lausanne).* 2015, 6:109. [10.3389/fendo.2015.00109](#)
29. Abdel-Maksoud SM, Hassanein SI, Gohar NA, Attia SM, Gad MZ: Investigation of brain-derived neurotrophic factor (BDNF) gene expression in hypothalamus of obese rats: modulation by omega-3 fatty acids. *Nutr Neurosci.* 2017, 20:443-8. [10.1080/1028415X.2016.1180859](#)
30. Sandrini L, Di Minno A, Amadio P, Ieraci A, Tremoli E, Barbieri SS: Association between obesity and circulating brain-derived neurotrophic factor (BDNF) levels: systematic review of literature and meta-analysis. *Int J Mol Sci.* 2018, 19:2281. [10.3390/ijms19082281](#)
31. Cui Z, Liu L, Kwame Amedvor F, et al.: High expression of miR-204 in chicken atrophic ovaries promotes granulosa cell apoptosis and inhibits autophagy. *Front Cell Dev Biol.* 2020, 8:580072. [10.3389/fcell.2020.580072](#)
32. Bruta K, Bhasin K: The role of serotonin and diet in the prevalence of irritable bowel syndrome: a systematic review. *Transl Med Commun.* 2021, 6:1-9. [10.1186/s41231-020-00081-y](#)
33. Roth W, Zadeh K, Vekariya R, Ge Y, Mohamadzadeh M: Tryptophan metabolism and gut-brain homeostasis. *Int J Mol Sci.* 2021, 22:2973. [10.3390/ijms22062973](#)
34. Mawe GM, Hoffman JM: Serotonin signalling in the gut--functions, dysfunctions and therapeutic targets. *Nat Rev Gastroenterol Hepatol.* 2013, 10:473-86. [10.1038/nrgastro.2013.105](#)
35. Erspamer V: Pharmacology of indole-alkylamines. *Pharmacol Rev.* 1954, 6:425-87.
36. Gunawardene AR, Corfe BM, Staton CA: Classification and functions of enteroendocrine cells of the lower gastrointestinal tract. *Int J Exp Pathol.* 2011, 92:219-31. [10.1111/j.1365-2613.2011.00767.x](#)
37. Rhee SH, Pothoulakis C, Mayer EA: Principles and clinical implications of the brain-gut-enteric microbiota

- axis. *Nat Rev Gastroenterol Hepatol*. 2009, 6:306-14. [10.1038/nrgastro.2009.35](https://doi.org/10.1038/nrgastro.2009.35)
38. Bellono NW, Bayrer JR, Leitch DB, et al.: Enterochromaffin cells are gut chemosensors that couple to sensory neural pathways. *Cell*. 2017, 170:185-198.e16. [10.1016/j.cell.2017.05.034](https://doi.org/10.1016/j.cell.2017.05.034)
39. Whalley K: Gut-brain communication: making contact . *Nat Rev Neurosci*. 2017, 18:452. [10.1038/nrn.2017.89](https://doi.org/10.1038/nrn.2017.89)
40. Luckensmeyer GB, Keast JR: Activation of alpha- and beta-adrenoceptors by sympathetic nerve stimulation in the large intestine of the rat. *J Physiol*. 1998, 510:549-61. [10.1111/j.1469-7793.1998.549bk.x](https://doi.org/10.1111/j.1469-7793.1998.549bk.x)
41. Lundgren O: Sympathetic input into the enteric nervous system. *Gut*. 2000, 47:33-5. [10.1136/gut.47.suppl_4.iv33](https://doi.org/10.1136/gut.47.suppl_4.iv33)
42. Shivachar AC, Eikenburg DC: Differential effects of epinephrine and norepinephrine on cAMP response and g(i3)alpha protein expression in cultured sympathetic neurons. *J Pharmacol Exp Ther*. 1999, 291:258-64.
43. Makita Y, Okuno S, Fujisawa H: Involvement of activator protein in the activation of tryptophan hydroxylase by cAMP-dependent protein kinase. *FEBS Lett*. 1990, 268:185-8. [10.1016/0014-5793\(90\)81004-8](https://doi.org/10.1016/0014-5793(90)81004-8)
44. McKinney J, Knappskog PM, Haavik J: Different properties of the central and peripheral forms of human tryptophan hydroxylase. *J Neurochem*. 2005, 92:311-20. [10.1111/j.1471-4159.2004.02850.x](https://doi.org/10.1111/j.1471-4159.2004.02850.x)
45. Huang Z, Liu T, Chattoraj A, et al.: Posttranslational regulation of TPH1 is responsible for the nightly surge of 5-HT output in the rat pineal gland. *J Pineal Res*. 2008, 45:506-14. [10.1111/j.1600-079X.2008.00627.x](https://doi.org/10.1111/j.1600-079X.2008.00627.x)
46. Young RL, Lumsden AL, Martin AM, et al.: Augmented capacity for peripheral serotonin release in human obesity. *Int J Obes (Lond)*. 2018, 42:1880-9. [10.1038/s41366-018-0047-8](https://doi.org/10.1038/s41366-018-0047-8)
47. Carey AL, Kingwell BA: Reducing peripheral serotonin turns up the heat in brown fat . *Nat Med*. 2015, 21:114-6. [10.1038/nm.3797](https://doi.org/10.1038/nm.3797)
48. Sharma BK, Patil M, Satyanarayana A: Negative regulators of brown adipose tissue (BAT)-mediated thermogenesis. *J Cell Physiol*. 2014, 229:1901-7. [10.1002/jcp.24664](https://doi.org/10.1002/jcp.24664)
49. Crane JD, Palanivel R, Mottillo EP, et al.: Inhibiting peripheral serotonin synthesis reduces obesity and metabolic dysfunction by promoting brown adipose tissue thermogenesis. *Nat Med*. 2015, 21:166-72. [10.1038/nm.3766](https://doi.org/10.1038/nm.3766)
50. Lasar D, Rosenwald M, Kiehlmann E, et al.: Peroxisome proliferator activated receptor gamma controls mature brown adipocyte inducibility through glycerol kinase. *Cell Rep*. 2018, 22:760-73. [10.1016/j.celrep.2017.12.067](https://doi.org/10.1016/j.celrep.2017.12.067)
51. Yabut JM, Crane JD, Green AE, Keating DJ, Khan WI, Steinberg GR: Emerging roles for serotonin in regulating metabolism: new implications for an ancient molecule. *Endocr Rev*. 2019, 40:1092-107. [10.1210/er.2018-00283](https://doi.org/10.1210/er.2018-00283)
52. Xiao H, Kang S: The role of the gut microbiome in energy balance with a focus on the gut-adipose tissue axis. *Front Genet*. 2020, 11:297. [10.3389/fgene.2020.00297](https://doi.org/10.3389/fgene.2020.00297)
53. Suárez-Zamorano N, Fabbiano S, Chevalier C, et al.: Microbiota depletion promotes browning of white adipose tissue and reduces obesity. *Nat Med*. 2015, 21:1497-501. [10.1038/nm.3994](https://doi.org/10.1038/nm.3994)
54. Yano JM, Yu K, Donaldson GP, et al.: Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell*. 2015, 161:264-76. [10.1016/j.cell.2015.02.047](https://doi.org/10.1016/j.cell.2015.02.047)
55. Martin AM, Yabut JM, Choo JM, et al.: The gut microbiome regulates host glucose homeostasis via peripheral serotonin. *Proc Natl Acad Sci U S A*. 2019, 116:19802-4. [10.1073/pnas.1909311116](https://doi.org/10.1073/pnas.1909311116)
56. Muller PA, Schneeberger M, Matheis F, et al.: Microbiota modulate sympathetic neurons via a gut-brain circuit. *Nature*. 2020, 583:441-6. [10.1038/s41586-020-2474-7](https://doi.org/10.1038/s41586-020-2474-7)
57. Bercik P, Denou E, Collins J, et al.: The intestinal microbiota affect central levels of brain-derived neurotrophic factor and behavior in mice. *Gastroenterology*. 2011, 141:599-609, 609.e1-3. [10.1053/j.gastro.2011.04.052](https://doi.org/10.1053/j.gastro.2011.04.052)