Eruptive Angiomatosis Triggered by COVID-19 Vaccination

Mohammed Shanshal ¹

1. Dermatology, Basildon University Hospital, Basildon, GBR

Corresponding author: Mohammed Shanshal, doctor.dermatology30@yahoo.com

Review began 02/26/2022

Review ended 03/04/2022

Published 03/07/2022

© Copyright 2022

Shanshal. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Despite meeting strict standards of quality, safety, and effectiveness, rare systemic and cutaneous side effects of coronavirus disease 2019 (COVID-19) vaccinations continue to be reported throughout the world. We report a case of eruptive cherry angiomatosis in a female following her first dose of COVID-19 vaccination with subsequent crops appearing after the second dose. The biopsy revealed dilated capillaries within the superficial dermis consistent with the clinical diagnosis.

Categories: Dermatology

Keywords: benign vascular lesions, eruptive angiomatosis, covid-19, covid-19 vaccine adverse effects, covid-19 vaccination

Introduction

Since the World Health Organization declared it a global pandemic in March 2020, coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has been spreading all across the world. In an effort to control its spread and associated fatality rates, COVID-19 has spurred the biggest vaccination campaign in history. Despite the well-documented safety profile of these vaccines, rare incidences of cutaneous and systemic adverse effects have been reported as millions of people receive their vaccinations worldwide [1,2].

Case Presentation

A 55-year-old female presented to the Ambulatory Care Unit at the Basildon University Hospital with a rash affecting her chest area as well as the abdominal wall, breast, and arms. She had first noticed these lesions developing in February 2021, 10 days after receiving her first dose of the Oxford-AstraZeneca COVID-19 vaccine. The rash had consisted of multiple red-purplish papules associated with flu-like illness. She had subsequently noticed more crops of these lesions appear after she had received her second dose in the middle of April. The lesions were asymptomatic and had not faded. The patient had been under the care of the Rheumatology team for psoriatic arthritis. She was on sulfasalazine and Enbrel injection. Her skin was normal and she only experienced nail changes associated with her psoriasis.

On examination, multiple erythematous papules of 2-3 mm were detected predominantly on her chest and abdominal wall; scattered papules on her arms and legs were also observed. Her blood exam on admission showed normal liver function, full blood count, clotting screen, renal function, and a C-reactive protein of 1 mg/L. She had a low positive antinuclear antibody level; however, this was of uncertain significance since she had normal myeloperoxidase and low proteinase 3 antibodies. Punch biopsy of the upper abdomen revealed dilated interconnecting capillaries within the superficial dermis consistent with the clinical picture of cherry angioma (Figures 1A-1D).

FIGURE 1: COVID-19-triggered cherry angiomas

A, B: erythematous papules involving the chest, breasts, and abdomen; C, D: hematoxylin and eosin stain (20x and 40x) showing dilated interconnecting capillaries within the superficial dermis

COVID-19: coronavirus disease 2019

Discussion

Based on the clinical appearance, the histopathological findings, and the temporal relationship between the first dose of vaccination and the flare-up following the second dose, a diagnosis of eruptive angiomas triggered by COVID-19 vaccination was made. Cherry angioma is a benign vascular lesion of unknown etiology more commonly associated with old age, pregnancy, and somatic mutations in GNAQ and GNA11 genes [3]. Its eruptive form may be seen in association with lymphoproliferative disorders, certain medications, and the human herpesvirus-8 [4].

Vascular lesions including eruptive cherry angiomas have been reported secondary to COVID-19, which could be mediated by angiotensin-converting enzyme 2 (ACE2) receptors; subsequent endothelial damage has also been reported [5,6]. Mohta et al. have reported 12 cases of eruptive pseudoangiomatosis, a similar vascular lesion that commonly occurs in the pediatric age group and is associated with ChAdOx1 nCoV-19 coronavirus vaccine (recombinant) in India [7]. The possible vascular injury [8,9] or immune dysregulation [10] caused by the COVID-19 vaccination could play a role in the development of these eruptive vascular lesions.

Conclusions

Cherry angiomas, also known as Campbell de Morgan spots, are harmless and generally asymptomatic vascular lesions comprising dilated capillaries. They typically present as bright red or purplish papules of varying sizes, and they are more common in older adults and pregnant women. Eruptive cherry angiomas can occur in association with certain medications, viral infections, and lymphoproliferative disorders. We reported a case of new-onset eruptive angiomas following the COVID-19 vaccination in a female patient. The report highlights one of the cutaneous side effects of COVID-19 vaccines. Even though these rare adverse events are relevant, they should not undervalue the importance of immunization against COVID-19 and its overall safety profile.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Bogdanov G, Bogdanov I, Kazandjieva J, Tsankov N: Cutaneous adverse effects of the available COVID-19 vaccines. *Clin Dermatol*. 2021, 39:523-31. [10.1016/j.clindermatol.2021.04.001](https://doi.org/10.1016/j.clindermatol.2021.04.001)
2. Al Khames Aga QA, Alkhaffaf WH, Hatem TH, et al.: Safety of COVID-19 vaccines. *J Med Virol*. 2021, 93:6588-94. [10.1002/jmv.27214](https://doi.org/10.1002/jmv.27214)
3. Klebanov N, Lin WM, Artomov M, et al.: Use of targeted next-generation sequencing to identify activating hot spot mutations in cherry angiomas. *JAMA Dermatol*. 2019, 155:211-5. [10.1001/jamadermatol.2018.4231](https://doi.org/10.1001/jamadermatol.2018.4231)
4. Borghi A, Benedetti S, Corazza M, et al.: Detection of human herpesvirus 8 sequences in cutaneous cherry angiomas. *Arch Dermatol Res*. 2013, 305:659-64. [10.1007/s00403-013-1346-5](https://doi.org/10.1007/s00403-013-1346-5)
5. Bouaziz JD, Duong TA, Jachiet M, et al.: Vascular skin symptoms in COVID-19: a French observational study. *J Eur Acad Dermatol Venereol*. 2020, 34:e451-2. [10.1111/jdv.16544](https://doi.org/10.1111/jdv.16544)
6. Agirgol S, Çaytemel C, Kolan AŞ, Vural H: A case of COVID-19 with papulovesicular rash that progressed to retiform purpura, accompanied by cherry angiomas. *Sao Paulo Med J*. 2021, 139:186-9. [10.1590/1516-3180.2020.0620.R1.0212020](https://doi.org/10.1590/1516-3180.2020.0620.R1.0212020)
7. Mohta A, Jain SK, Mehta RD, Arora A: Development of eruptive pseudoangiomatosis following COVID-19 immunization - apropos of 5 cases. *J Eur Acad Dermatol Venereol*. 2021, 35:e722-5. [10.1111/jdv.17499](https://doi.org/10.1111/jdv.17499)
8. Islam A, Bashir MS, Joyce K, Rashid H, Laher I, Elshazly S: An update on COVID-19 vaccine induced thrombotic thrombocytopenia syndrome and some management recommendations. *Molecules*. 2021, 26:5004. [10.3390/molecules26165004](https://doi.org/10.3390/molecules26165004)
9. Shakoor MT, Birkenbach MP, Lynch M: ANCA-associated vasculitis following Pfizer-BioNTech COVID-19 vaccine. *Am J Kidney Dis*. 2021, 78:611-3. [10.1053/j.ajkd.2021.06.016](https://doi.org/10.1053/j.ajkd.2021.06.016)
10. Akinosoglou K, Tzivaki I, Marangos M: Covid-19 vaccine and autoimmunity: awakening the sleeping dragon. *Clin Immunol*. 2021, 226:108721. [10.1016/j.clim.2021.108721](https://doi.org/10.1016/j.clim.2021.108721)